



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>







ON
DISEASES OF THE ABDOMEN.



PATHOLOGICAL AND PRACTICAL OBSERVATIONS
ON
DISEASES OF THE ABDOMEN,
COMPRISING THOSE OF THE
STOMACH, AND OTHER PARTS OF THE ALIMENTARY
CANAL, ŒSOPHAGUS, CÆCUM, INTESTINES,
AND PERITONEUM.

BY
S. O. HABERSHON, M.D. LOND.,
FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS; SENIOR ASSISTANT PHYSICIAN TO GUY'S HOSPITAL,
AND LECTURER ON MATERIA MEDICA AND THERAPEUTICS; LATE DEMONSTRATOR OF MORBID
ANATOMY, AND CURATOR OF THE MUSEUM AT GUY'S; LATE PHYSICIAN TO THE
CITY DISPENSARY, ETC., ETC.

SECOND EDITION,
CONSIDERABLY ENLARGED AND REVISED.



LONDON:
JOHN CHURCHILL, NEW BURLINGTON STREET.
MDCCCLXII.

~~150. a. 61.~~
151. a. 63.



PRINTED BY J. E. ADLARD, BARTHOLOMEW CLOSE.

ADVERTISEMENT.

IN bringing out a second edition of my work on Diseases of the Alimentary Canal, I have been encouraged by the favorable manner in which the first edition has been received by the profession. Much care has been given in thoroughly revising each portion of the work; and important additions have been made, not only in numerous interesting cases still further illustrating the phases of disease, but large sections have been added to the text. My former suggestion of opening the stomach as a means of sustaining life in obstruction of the œsophagus has been attempted; and the result is recorded, as warranting the repetition of the operation at an earlier period of the complaint. The functional, as well as the organic diseases of the stomach have received fresh elucidation, and prominent symptoms have been presented in their relative significance. The morbid affections of the cæcum, and obstructive disease of the intestines have been enlarged upon; and a chapter on Peritonitis has also been appended to the work.

ON
DISEASES OF THE ABDOMEN.

development, organization, nature, &c., each new observer considering himself more clear-sighted than his predecessor, although he has merely substituted one term for another. This living force is in close correlative relation with other physical forces, and the fuller investigations of physiological science show that the same forces are in operation, namely, gravitation, chemical action, &c., in the living organism, as external to it; modified, it is true, by another force, namely, life. And as in the science of physics generally, so in the study of living phenomena, we must ever bear in mind that a like cause always produces a like effect. Vague observation, and the superficial remarks of some writers, would lead us to suppose, that, in living phenomena, the same cause is followed, at first by one effect, then by another; interpreting fixed realities by prejudice rather than by reason.

Some phraseology is necessary to express our meaning and ideas, and one great difficulty is overcome, if we can understand that the same words convey to each the same thought. It may be convenient, as we have mentioned, to regard life as the resultant of certain forces, and disease as a deviation from the normal direction. If any of the forces which are in natural operation be modified in intensity, a deviation is the result, and diseased action is produced, the resultant being necessarily changed; still the tendency is such that, on the withdrawal of the modifying force, the normal course is reassumed. Not only may it be natural force which has led to this departure from the healthy state, but new force may be added, as much as when the earth in its orbit is disturbed by the attraction of some other celestial body.

In diseases, many sources of change arise—modifying forces—thus syphilitic or miasmatic poison, small-pox, &c., alter the character of every function; new substance is added as much as in the voltaic battery, when the fluid in one or other cell may be changed by the admixture of any substance; this may be merely of the kind already existing, or a foreign body; in any case, the phenomena—the same in general development—are modified.

Such, to some extent, is the case in pathological changes. These changes, produced by perverted nutrition, or altered vital forces, are in many instances of such a character, that no examination of the structure itself could discern the state which had been produced; as fruitless would it be to search in the nerve of a limb for the altered force which had led to spasm, as to expect to find a telegraphic message by a microscopical examination of the wire, although the structure of both had been transiently modified by the disturbance of the forces they transmitted. If the character of the disturbance in disease is one, which, like a polar force, reverts to its former condition, no trace can be found on inspection, but, in many instances, obvious structural modifications are the result.

Diseased action, however, as generally manifested, is the resultant, not of one, but of several changes in the normal condition, and very few persons are literally in perfect health. The living forces are modified by hereditary tendency, as struma; to this, perhaps, is added syphilis, to that miasm; still further sudden changes of temperature, improper supply of nourishment, of heat, and light; each of these may act as fresh sources of deviation from the normal healthy direction of living action, superadded to the resultant produced by the previous combination.

Some have supposed that acute disease quickly passes off, and that with the subsidence of the more marked symptoms no trace is left behind, but very generally this is not the case; the attentive study of pathology will soon convince us of the contrary; new exciting causes of disease arise, perhaps of a different character, but the resultant (to revert to the previous phraseology of forces) is not precisely the same, the former diseased state acts as a modifying force. This course may be often repeated, and if the changes have been such as to entail discomfort or weakness, chronic disease is said to exist; but if without these, the patient generally passes for one in sound health. It is the acumen of the practical physician that detects the traces of previous morbid action, and he alone knows how

rightly to estimate the course likely to be assumed by any new addition to a state widely diverse from that of health; hence, also, the variety of diseases by which the same organ is affected, the causes are different and so of necessity their effect.

It is the province of medical science, rightly to estimate the effects produced on vital action by any disturbing causes, and to study their almost endless varieties; several general characters lead us to group these effects into classes; and although in this volume we have spoken of diseased action as manifesting itself especially in the alimentary canal in the changes described, and the symptoms depicted, it will be found that these parts are in many cases only affected in common with the whole economy, whilst in others that the special manifestation of morbid action is in the abdomen.

Numerous means are available for checking and modifying diseased action, and we must protest against the ignorance of those who regard the draught of medicine as the only important agent. The skill of the physician is often most manifest in the suggestion of hygienic measures which by many may be assumed as of trifling moment; and whilst it is perfectly true that many morbid conditions cease after a time, or that the frame becomes so accustomed to perverted action that the balance of functions is apparently maintained without cognizance on the part of the patient, still the aid of medical science is most important. The first agents to which we must refer are those which are in constant operation in the maintenance of life—in one state preservative of health, in another the cause of disease—as the character of the air breathed, whether saturated with moisture, poisoned with miasm, or with the decomposing effluvia of crowded cities, as compared with that found in more elevated situations, on the sea or its coasts, &c.; so also with reference to diet, to light, to clothing, to temperature, to habits of mental or bodily training or exercise; the right use of all these is not less important in the restoration to health, than in the maintenance of it, and in both cases alike, is within the province of medical science.

|

Too frequently medicine is taxed to obviate the disease whilst its cause is pertinaciously adhered to. It is in vain to recommend to the dyspeptic patient remedies which would certainly mitigate his disease, whilst intense anxiety remains, and hurried, half masticated meals, are taken at irregular hours and without due moderation—or, again, it is useless to direct means to relieve a disturbed brain, whilst excitement is added to excitement, the senses stimulated by light, noise, animated conversation, and active thought—or to give opium to check peritonitis, to quiet muscular movement, when the patient is allowed to move from the recumbent position.

The consideration of the fundamental conditions of life demands strict attention, not only as indicating the tendency vital action has to assume its natural direction, but also in enabling us to use and apply effective means for the removal of morbid processes; and we are at no great loss to understand the intellectual power and acumen of those who, because everything is not known in the pathology and treatment of disease, would contemptuously discard that which is known and established. We deplore the ignorance of those who know not the value of opium in peritonitis, &c., of iodide of potassium in secondary syphilis, of purgatives and mercurials in hepatic engorgement, of preparations of steel in many forms of anæmia, of quinine and arsenic in intermittents; our object is not to confute errors which arise from wilfully closing the eye to light already attained, but to seek to add facts upon which science may safely advance.

We have generally divided the chapters according to the anatomical divisions, rather than in a strictly pathological manner. The first chapter contains many interesting cases of disease of the *Œsophagus*, some of them obscure in their pathology, and very insidious in their origin; some instances of ulceration, perforating the trachea or bronchi, which we have described, have generally, and we believe incorrectly, been considered as instances of cancerous disease. The diseases of the stomach constitute the subject of the next section, and have obtained

from authors very considerable attention ; we believe that there are forms of ulceration, superficial and evanescent, which leave scarcely more trace in the mucous membrane of the stomach than the aphthous ulceration of the mucous membrane of the mouth, whilst others are permanent, and are manifest after death ; and indeed we find the same forms of diseased action in the mouth as in the stomach ; thus both are affected by inflammatory congestion, perverted epithelial growth and secretion, sluggish condition of the circulation, acute inflammatory disease, as well as by fibroid and cancerous disease. Ulceration of the stomach is probably a more common condition than is generally supposed, and in many instances yields to judicious treatment ; the instances we have adduced show that there are several distinguishing marks by which it may be known from cancer. Fibroid degeneration of the pylorus has generally been considered as of a cancerous nature ; and whilst we are unable to remove this almost certainly fatal form of disease, we may, as in cancer, do much to mitigate the symptoms and to prolong life.

In the so-called functional disease of the stomach, chemical research has removed much that was obscure, and will do still more to clear up the pathological changes induced ; and the investigation of the physiological connections of the sympathetic nerve and the branches of the semilunar ganglia will enable us more correctly to estimate the very diverse symptoms produced in dyspepsia, many of which have their origin in this source. The right estimate of such symptoms as pain in the region of the stomach and vomiting are most important in the diagnosis, and equally so in the treatment of gastric affections.

The chapter on the Duodenum presents us with instances of disease which closely simulate maladies of the pyloric extremity of the stomach.

The next chapter is on Gastro-Enteritis and Enteritis, diseases in which correct diagnosis is very important ; in the latter class of diseases especially, life may be easily sacrificed by time being thrown away, and by improper treatment ; in the former,

with judicious diet, warmth, demulcents, &c., recovery generally takes place. Whilst we strongly recommend, in many of these instances of gastro-enteritis, the avoidance of mercurials, the value of salines, of bicarbonate and chlorate of potash, and carbonate of soda, are well known to those who have carefully watched the effect that has followed their administration.

In the chapter on Strumous Disease of the Intestine and Peritoneum we have sought to show that these diseases are only part of a general perverted nutrition, and that, in many instances, disease in other organs is entirely obscured by the more marked affection of the abdominal viscera; here, also, we should strongly urge the avoidance of mercurial medicines and of drastic purgatives; the lives of many delicate children are sacrificed by worm powders and quack nostrums administered in these diseases.

The next class of cases in Chapter VIII is on diseases of the Cæcum and its Appendix. We have shown that unusual freedom of the cæcal attachments may determine intestinal obstruction from rotatory movements of the intestine; the symptoms and treatment of cæcal distension and of local enteritis are described, as well as the more serious consequences of perforation of the appendix; we have given numerous instances of these forms of affection, and their perusal will show the great similarity in the symptoms and their general course. Dr. Burns, in a valuable paper in the 'Medico-Chirurgical Transactions,' described, several years ago, some of these affections. It would appear that the symptoms of cancerous disease of the cæcum are different from simple cæcal enteritis and perforation; and that in many cases we may discriminate the character of the complaint. It will be found that treatment may do much to relieve and to assist the cure of cæcal disease; the pain demands rest, and it is well after the acute pain has subsided, still to maintain absolute repose for several days. The bowels are often confined, but the use of purgatives generally aggravates the disease without effecting the desired operation; this is better obtained by the application of leeches and by opium; those remedies, which

diminish the enteric inflammation lead most speedily to the subsidence of the morbid symptoms.

In the ninth chapter we have pointed out the characters of the several forms of Diarrhœa, but we are well aware that diarrhœa is merely a symptom of very different conditions, and that in many instances it passes almost imperceptibly into dysentery.

Dysentery and Colitis are the subjects of the tenth chapter; and the instances we have adduced show that inflammation of the colon, of a most severe form, arises in our own country. Most of the writers on this subject are those who have observed it in its worst forms abroad or in Ireland. In some of the cases typhoid fever was simulated; in others, perforation of the colon had taken place; in one there was pyæmia and commencing suppuration in the liver; in several chronic cases the secondary effects were shown in producing contraction of the intestine, perforation, and artificial anus, &c.; as regards abscess in the liver, in one case the abscess had dried and contracted; in another, fresh diseased action was set up around it, and abscess in the brain was the result. We must confess, that in some of the most severe forms all treatment is ineffectual to cure, whilst it partially soothes and relieves; but in the great majority of instances means may be used which effectually combat the symptoms of disease.

We had intended to give some observations on Asiatic cholera, but for several reasons have not done so; first, because although the disease manifests itself more apparently in the disturbed functions of the alimentary canal, it has not been clearly shown that the disease is really one affecting alone or even principally the abdomen; and secondly, all the facts known in reference to this disease are better and more clearly given in the report drawn up by Dr. Baly and Dr. Gull.

In the chapter on Typhoid Fever, we have merely described the condition of the abdominal affection, without entering into the general question of fever, and its treatment;—in the latter, it is well to guard against the danger of so freely administering

opiates to check diarrhoea as to lead to cerebral oppression, and excessive engorgement of the lungs, from imperfect performance of the respiratory function.

In the chapter on Colic, we have cursorily spoken of the simpler forms of the disease, and have separated the more severe varieties of ileus—internal strangulation, intussusception, and cancerous disease of the colon—in Chapter XIV. It will be found that whilst the latter conditions bear a strong general resemblance in producing fatal constipation, they may, in many instances, be distinguished, the one from the other; intussusception closely simulates simple colic, but, in not a few instances, is accompanied by discharge of bloody mucus, or with actual diarrhoea; and this latter symptom sometimes arises even with cancerous disease of the sigmoid flexure. We have very strongly urged the avoidance of drastic purgatives, calomel, colocynth, and even milder purgatives, and as strongly recommended the free administration of opium; the cases detailed almost uniformly show that, where purgatives were given, vomiting, pain, and distress, were increased, whilst these and other symptoms were, on the contrary, relieved by opium. Opium, in such cases, appears to be the best means of procuring relief to the bowels, if an action be possible. In the chapter on Worms, we have designedly been very brief in our remarks.

The cases of Suppuration in the Abdominal Parietes, and of Perforation of the Intestine from Without, are an interesting series; many of them are obscure in their diagnosis, and different in their course. Great care is required in watching the symptoms as they become fully developed, and in avoiding the aggravation of them by too active treatment.

The serous membrane of the intestines, the peritoneum, is so intimately connected with the morbid conditions of the alimentary canal, that we have appended some remarks on its diseases. In very many instances the serous membrane is implicated by direct extension of disease; in other cases, the peritoneal change is the expression of a general morbid condition of the whole system.

The cases we have recorded might have been given at greater length, and on each one fuller remarks might have been made ; but the design of the work has been, in a very few words to point out the peculiarities of each instance, embodying in more general deductions the apparent conclusions derived from the whole. Our desire is to shed some light on the difficulties which often present themselves in the daily practice of the profession, and to suggest means whereby the alleviation of disease may be promoted.

22, WIMPOLE STREET, CAVENDISH SQUARE.

CONTENTS.

CHAPTER	PAGE
I. INTRODUCTION	1
II. ON DISEASES OF THE ESOPHAGUS	13
III. ON ORGANIC DISEASES OF THE STOMACH	83
IV. ON FUNCTIONAL DISEASES OF THE STOMACH—HEMATEMESIS— PAIN—VOMITING	185
V. ON DISEASES OF THE DUODENUM	242
VI. ON MUCO-ENTERITIS AND ENTERITIS	268
VII. ON STRUMOUS DISEASE OF THE ALIMENTARY CANAL	284
VIII. ON DISEASES OF THE CÆCUM AND APPENDIX CÆCI	313
IX. ON DIARRHŒA	362
X. ON COLITIS AND DYSENTERY	381
XI. ON TYPHOID DISEASE OF INTESTINE	422
XII. ON COLIC	429
XIII. ON CONSTIPATION	441
XIV. ON INTERNAL STRANGULATION—INTUSSUSCEPTION—CARCINOMA OF THE INTESTINE	459
XV. ON INTESTINAL WORMS	522
XVI. ON PERFORATION OF INTESTINE FROM WITHOUT—ABSCESS IN THE ABDOMINAL PARIETES, EXTENDING INTO THE INTESTINE— FECAL ABSCESS	529
XVII. ON PERITONITIS	547






Fig. 2

Fig. 1



Fig. 1

PLATE III.



Author del. Luffen West co.

W West 1879.

DESCRIPTION OF PLATES.

PLATE I.

Stomach presenting a chronic ulcer ; at its upper margin the pneumogastric nerve is shown extending into dense fibrous tissue. The pancreas and the left lobe of the liver formed the base of the ulcer ; the latter presented fibroid degeneration of its structure.—Case LIV.

PLATE II.

Stomach exceedingly contracted from chronic ulceration, with villous growth. Simulating cancer.—Case LXX.

1. External view, resembling the colon in appearance.
2. Internal surface, showing ulceration near the pylorus, and villous growth near the centre of the stomach.

PLATE III.

FIG. I. Section of the mucous membrane of a portion of the ileum in acute inflammation of the colon and ileum, showing the surface covered with false membrane (diphtherite) and continuous with the mucous follicles.—Case CLXV.

FIG. II. Section of a solitary gland from the cæcum, from the same Case, CLXV, showing (a) the edges of a raised portion intensely injected from distended capillaries ; (b) surface of mucous membrane covered with diphtheritic granular membrane ; (c) opening into the gland ; (d) small phosphatic crystals.

FIG. III. Appearance of inflamed colon in Case CLXXVI ; (a) false membrane composed of granule cells ; (b) surface of a portion of the colon beneath the false membrane ; (c) follicle or crypt containing cells similar to those composing the false membrane.

FIG. IV. Colloid cancer of the sigmoid flexure $\times 400$ diam. ; (a b) columnar epithelium ; (c) nuclei with granular blastema ; (d) large cells, with large nuclei, and some with several nuclei in them ; (e) intervening delicate tissue ; (f) elongated fibre cells.—Case CCVIII.

FIG. V. Obstruction of sigmoid flexure by cancerous growth; (a) columnar epithelium and nuclei; (b) fibrous tissue beneath the mucous membrane; (c) interspaces filled with nuclei; (d) surface of mucous membrane composed of dense fibre tissue.—Case CC.

PLATE IV.

FIG. I. Cæcum inverted, appendix towards the pelvis, where it was adherent; ascending colon commencing opposite the ileum.—Case CXXXIX.

FIG. II. Cæcum inverted and twisted on its own axis into the left hypochondriac region, appendix close to the spleen; ascending colon constricted; constriction increased by band of adhesion to the sigmoid flexure, which appeared to have been the primary cause of the fatal twist and obstruction.—Case CXXX.


FIG. III. Position of intestines in case of intussusception of cæcum and ascending colon into descending colon and sigmoid flexure; the commencement of the rectum is drawn from its position, to show the strangulated bowel within. Dr. Hughes' case, No. CXCH.

PLATE V.

Dissection showing the distribution of the pneumogastric nerve on the anterior surface of the stomach, its extension to the pancreas and pylorus, and its connection with the semilunar ganglia, &c.; (a) oesophageal extremity of the stomach; (b) pylorus; (cc) pneumogastric nerves; (ee) branch of the pneumogastric to the pancreas, connected also with the sympathetic, and then passing onwards to the pylorus; (ff) other branches to the pylorus; (ggg) branches of the pneumogastric nerve distributed on the anterior surface of the stomach, presenting a peculiar dichotomous division, and repeated union of its branches; (h h h h h) splanchnic nerves; (i) aorta; (j) diaphragmatic artery, with a filament of nerve upon it; (kk) coronary artery; (l) splenic artery; (m) hepatic artery turned aside from its position in front of the aorta, and from its origin at the celiac axis; and thus it appears to be behind the aorta; the large branches of the sympathetic nerve upon it are continuous with the portion of ganglion (r*) close to the coronary artery; (n) vena portæ; (oo) supra-renal capsules receiving numerous nerve filaments; (rr) semilunar ganglia, and descending branches to the mesenteric artery and renal plexus, &c.; (s) mesenteric artery drawn aside.

In this distribution of nerves, the close sympathy of the stomach with the parts supplied by the semilunar ganglion is explained; thus, it is brought into connection with the liver by its hepatic branches, with the pancreas, with the diaphragm and phrenic nerve, with the supra-renal capsules, and by its descending branches with other abdominal viscera.

PLATE IV




Auctor del. Tuffen West sc.



A. West del.





ON

DISEASES OF THE STOMACH.

CHAPTER I.

INTRODUCTION.

THE function of digestion is essentially connected with life and health; and slight deviations from its normal performance produce suffering in a greater or less degree. He is, indeed, fortunate who can pass through his daily duties without having the thoughts and attention directed to these operations for the solution, absorption, and assimilation of nourishment, which in health are completed unconsciously, without attention, or sense of pain. If there be severe derangement of the digestive functions, not only is the attention directed to them, and discomfort entailed, but there is reaction upon the higher capabilities of man's nature: the energies of the brain are enfeebled, the memory is defective, the will vacillates, and the intellectual powers are less free to guide in daily duty, avocation, and research. The strength and muscular movements are diminished, and the pleasure of life changes to daily suffering and anxiety. Contrast the vigour of mind and body during health, with the enfeebled energy of the dyspeptic and hypochondriac. In the former state there is no impediment to the exercise of deep thought and labour, in any sphere to which the mind may be directed; the whole attention in the latter is absorbed by those functions, which are at best only subservient to the mainly exercise of mind and will.

If the digestive process be altogether checked, and no new supply of nourishment be absorbed and assimilated; if no restoration be made to the waste entailed by the exercise of every function, life must sooner or later cease; and disease, in its ravages, presents few spectacles more distressing to witness, than the gradual wasting of the frame, and cessation of life itself, from the non-supply of food. Thus the whole system sympathises with disorder of the alimentary canal.

A knowledge of the structure and functions of each part of the digestive apparatus, is necessary for the right comprehension of its diseases. The structures of the alimentary canal are various, and their sympathies universal; but in health these are so combined as to form a beautiful and harmonious whole: thus, 1st, we find a mucous membrane richly supplied with glands, lining the alimentary canal throughout its course; these glands are for secretion and excretion; the secretions from these act physically or chemically in the digestive process, whilst the excretory glands separate noxious or effete principles from the blood. 2nd. Beneath the mucous is the muscular coat, essential for the execution of the required movements, and for the propulsion of the contents of the canal. 3rd. The peritoneal or serous covering, which by its smoothness permits of the movement of one portion of the intestine upon another, and allows distension to take place. 4th. The binding tissues, which are found between these previously mentioned tunics, and which support other equally essential parts, namely, 5th. The blood-vessels and lymphatics; and, 6th, the nervous supply from the sympathetic and cerebro-spinal nerve. As Abercrombie has remarked, in reference to diseases of the stomach, so also, it may be added, in reference to every part of the alimentary canal; for the proper performance of the function of digestion, the mucous membrane must be in health, the secretions normal, the supply of blood and nervous energy such as required, and the movements free. It must, however, be borne in mind, that the alimentary canal contains substances which are, strictly speaking, external to the living agency and control of animal life; and that those chemical forces, which we find in operation external to the body, act in the same manner within the stomach and small and large intestines: the food becomes dissolved when the same solvents

are provided, and other circumstances adapted, as to temperature, movements, &c., equally in a phial as in the stomach. The fermentation of its contents takes place in the stomach and canal, as well as in any chemical receiver; and these facts have to be remembered in the study, as well as in the treatment, of disease. Chemical force is in operation throughout the whole animal economy; it is modified and controlled by the living power, or it is free to act alone.

Each of the parts which have been mentioned, may be alone diseased, or all conjointly; the symptoms arising from each are in some cases distinct, in others we cannot separate the one from the other.

1st. Mucous membrane and its secretions. The derangement of these constitute, perhaps, the greater part of the milder ailments of the alimentary canal. The symptoms vary according to the part affected: in the stomach, producing some of the forms of *dyspepsia*; in the intestines, constipation, diarrhoea, &c. But, when the mucous membrane alone is affected, it appears probable that pain is not produced, and this circumstance we must regard as a merciful arrangement. The lining membrane is exposed to varied causes of irritation, but we do not experience pain; if such were the case, every portion of undigested food might produce suffering; in some cases severe pain is found in indigestion, but this arises from an extreme sensibility of the sympathetic and other nervous supply of the stomach, &c., and, is not due to the mucous membrane alone.

Dr. Beaumont, in his observations on the stomach of Alexis, sometimes observed the mucous membrane dry, injected, and much irritated without the production of pain; so, also, I have observed actual inflammation of the stomach, as found in cases of poisoning by oxalic acid, by chloride of zinc, and even by arsenic, without pain from first to last.

2nd. The muscular coat we find so stimulated, that it rapidly contracts, and impels onwards its contents; or, it is so enfeebled as to retain them; sometimes it is spasmodically contracted, or again, dilated, as in the forms of *colic* and flatulent distension, &c. These conditions appear to be productive of pain, sometimes of a very intense form, as we find in the griping of *colic*, and in *enteritis*, &c. As long as the peristaltic action is uniform,

regular, and healthy, we are unconscious of the movement; but as soon as it becomes irregular or tumultuous, retarded or spasmodic, we are sensible of uneasiness, or even of severe pain; the muscular coat of the intestine is probably excited to contraction by the direct stimulus of its contents, but the harmony of its movements is due to the supply of nervous influence which it receives.

3rd. The peritoneal or serous investment also manifests its derangement by pain; and here, again, is a wise provision, for as its disorders require rest, or rather an absence of movement of the coils of intestine one upon another, and the pain of peritoneal disease is increased by muscular exertion, so the patient becomes prompted to assume that position, and to retain that state, which is the best suited for the restoration from disease. The observant pathologist and physician know, practically, the importance of rest in the recumbent position, and they follow the teaching of nature in their stringent directions: by this means inflammation is localized, and when perforations of the intestine have taken place, the injury is limited and life may be prolonged.

4th. The state of the investing or binding tissues; and, 5th, the supply of blood, are important considerations in the study of these diseases of the intestine. The former, in some cases, appears to be the seat of fatal malady, as in the forms of constriction of the pylorus, and in cancer. Still more does the latter, the supply of blood, call for attention: it may be in excess, as in active or passive hyperæmia; in pulmonary, cardiac, and hepatic diseases the engorgement of the mucous membrane leads to peculiar and characteristic symptoms; the rupture of vessels, or ulceration into them, causes hæmorrhage into the canal; and again, a scanty supply or depraved condition of blood prevents the proper performance of digestion, as after great hæmorrhage, in over-lactation, in purpura, scurvy, or starvation. The appearance of the blood discharged into the several portions of the canal greatly differs; thus, in disease of the stomach, it is generally ejected as dark semi-coagulated blood; sometimes, as in the latter stages of cancer, as coffee ground substance; or, in the acute disease of the stomach and duodenum, according to Dr. Fraser and others, as a green fluid. If, however, blood be passed

into the duodenum, and discharged per rectum, it has a black and tarry character ; and in proportion as the source of the discharge is near or more distant from the rectum, a sanguineous appearance is retained.

6th. The state of the nervous supply is often lost sight of ; this is a most complicate system of nervous fibrils and ganglia, which are intimately connected with the cerebro-spinal centres, and with the ganglionic centres of other parts, as of the lungs, the heart, and the urino-genital organs. Many of the signs of intestinal disease arise from this cause, and they have been dwelt upon by various authors. In the Guy's Hospital Reports of 1856, I have described some dissections and observations on this supply of nerves ; they surround the vessels, are distributed with them, and reach every part of the intestine. The sympathy of abdominal disease with other organs is due to this supply. In indigestion, we find cephalalgia, depression of spirits, impaired mental energy, disordered sensations of general or special sense ; and all these arise from the connexion of the sympathetic and the cerebro-spinal nerve. So again, the throbbing of the vessels, the excited or irregular action of the heart in dyspepsia, proceed from the union of the cardiac ganglia with the solar plexus of nerves. With the lungs, the kidneys, the uterus, we notice similar sympathetic disturbance ; and oftentimes, in a most marked manner, the skin is observed in close connexion with the internal mucous membrane ; this disorder of the alimentary canal induces many forms of cutaneous eruption, as urticaria from partaking of mussels ; or the more chronic diseases of lepra, eczema, &c. These sympathies may, however, be due to the vascular condition, as well as to the nervous relation, of one structure with another. This relationship of parts, however, sometimes acts in a reverse direction ; the alimentary canal is affected, secondarily, from disease of other structures, for example, vomiting is a symptom of disease of the brain, of the kidney, and of the uterus. But beside these, there appear to be symptoms of primary disease of the alimentary canal, which are due directly to the sympathetic nerve. 1. A remarkable depression of the pulse, which we often find in these diseases of the abdomen, when the pulse becomes soft and compressible, and often irregular. 2. A sense

of sinking and exhaustion is one of the most marked signs of abdominal disease; and in some cases this exhaustion leads to sudden death, not only in cases when a person may have died from a blow on the epigastrium, but in other instances. A short time ago I observed a man suffering from aneurism of the descending aorta; he endured very severe pain, and the pulse became much enfeebled; in a few days he died with comparative suddenness. On examination, an aneurism of the aorta was found at the diaphragm; it had led to absorption of the bodies of the vertebræ, but there had been no extravasation of blood into the peritoneum, the cellular tissue, nor into other parts. The aneurismal sac was about four inches in length, and one and a half in height; it had pushed aside the pillars of the diaphragm, which were white and degenerated; the splanchnic nerves were stretched across the sac, and the semi-lunar ganglion pushed considerably forward and pressed upon. I think we were justified in believing that in this case the depression, and comparatively sudden death, were in great measure due to the pressure on the great sympathetic nerve centre of the abdomen. We have, also, often observed in cases of gastritis from poisons, as arsenic, dilute sulphuric acid, chloride of zinc, oxalic acid, that the pulse becomes remarkably depressed; and sometimes, where we might have been led, from the absence of pain and other symptoms, to have given a favorable prognosis, the patient has suddenly died.

The pneumo-gastric nerve has an important influence on the stomach. This was shown, in a marked degree, by the experiments of Dr. Wilson Philip, who demonstrated the effect of section of the pneumo-gastric nerve on digestion, in checking its progress: it does not, however, completely prevent it, but only for a time checks the secretion of gastric juice. The irritation of this part of the pneumo-gastric sometimes leads to symptoms indicative of disturbance of the pulmonary branches of the same nerve: cough may be set up; and it is probable that the converse takes place; the pulmonary branches may cause reflex influence on the stomachic branches, and produce violent vomiting. In the 'Medical Times and Gazette,' is a very interesting paper by Mr. J. Hutchinson on the Dyspepsia of Phthisis; and many have found in the early stage of phthisis

that the power to digest food is impaired, the diminished nutrition tending greatly to promote the deposition of low organized product in the lungs.

Not only do the signs of abdominal disease arise from the derangement of the structures of the canal, and from changes in the secretions and contents of the same, but the administration of remedies is guided by similar considerations. Many may be led to the use of means by mere empiricism, but the observations of Chambers, Turnbull, Budd, Handfield Jones, &c., lead to a more scientific and correct treatment, by directing to the physiological chemistry of digestion or by enlightened pathology.

Remedies may be classified in the following manner:—

1. Those agents which are used to check fermentative and chemical action.
2. To remove offending or injurious materials, and excreta.
3. To correct or improve the secretions from the mucous membrane, or those poured into the canal.
4. To affect the muscular coat, and its movements.
5. To alter the state of the circulation and vessels or absorbents.
6. To act on the abdominal sympathetic nerve.

Dr. Headland has directed attention, in his valuable essay on the action of medicines, to their mode of operation, considering them chemically or mechanically, prior to absorption; then, after entering the blood, as influencing either its constituents or the muscular or nervous structures to which it is supplied; and lastly, in their elimination from the body. These remarks forcibly apply to the action of agents in the treatment of abdominal disease. It is, however, often lost sight of, that whilst the alimentary canal is the structure by which remedies can be most easily made to enter the blood, and there exert their curative influence, it may be in such a condition from morbid changes, that no absorption can take place; and the administration of remedies by the stomach may become almost useless, as far as regards their ultimate action after absorption.

1. Agents which are used to check fermentative action or chemical decomposition.

Chemical science has done much, and will do still more, to

suggest means of counteracting changes of this character. Dr. Turnbull has dwelt, in his work, on the varied forms of fermentative action, and has shown that some agents possess in this manner considerable power: as creosote, sulphurous acid and sulphites, charcoal, so also alcohol, &c. We have much to learn on this subject; and it would well repay the labour of some one well versed in chemical science to extend these researches.

2. As agents for the expulsion of injurious matters, or excreta, we have the whole class of emetics, of laxatives, of purgatives, and the different forms of enemata.

3. To correct the secretions from the mucous membrane; this term is, I believe, strictly correct, for the mucus in the canal is sometimes of an irritating character, and we may do much to change its state after secretion, at the same time that we use means to prevent such abnormal character of secretion from taking place: demulcents of the following kind, as milk, arrow-root, gum acacia, linseed, sheath the mucous membrane; whilst lime-water, chalk, solution of potash, carbonate of soda, lessen the irritating character of the secretion. At the same time, to diminish inflammatory congestion, other agents are called for, as ipecacuanha, potash, soda, magnesia, and some of their salts, or mercurials, antimony, &c. If the object is to correct secretions arising from enfeebled or relaxed state of the membrane, we have vegetable and mineral astringents and tonics, mineral acids, &c.; others stimulate to greater secretion, when there is deficiency, as some irritants, ipecacuanha, salt, capsicum, pepper, &c.

4. Among remedies which act on the muscular movements of the intestine, I may enumerate the class of purgatives, magnesia, and strychnia, as increasing peristaltic action; and conium, opium, henbane, as diminishing these intestinal movements.

5. Those which act on the sympathetic nerve, diminishing its sensibility, are chloroform, hydrocyanic acid, opium, bismuth, oxide and nitrate of silver; as tending to increase its sensibility, steel, quinine, vegetable and mineral tonics, alcohol, &c.

These remedies are variously combined in the treatment of abdominal disease; and by combination their action is modified,

or their efficiency is increased; their presence may thus be better tolerated, and their absorption be facilitated.

The dietetic regimen is one of the most important subjects in diseases of the stomach and intestine; as in other visceral diseases we cannot obtain rest of the affected organ, but we can shield it from unnecessary irritation and fresh excitement.

It will, often, however, be found, that the state of the nervous system modifies the effect of remedies. If a highly sensitive patient, hysterical or hypochondriacal, be led to suppose that a medicine will produce a certain effect, the mind is so directed and influenced, that a powerful action may be produced; or, if a patient firmly believe that a particular medicine or plan of treatment will do him injury, we shall, in all probability, find that the symptoms are described as greatly aggravated thereby, and no argument will remove this persuasion. Thus, in a patient who had suffered from hemiplegia, and was in a nervous condition, but who could not be persuaded to discontinue medicine, two table-spoonfuls of spring water were followed by violent purging, and when changed for a pill of bread the same effect was produced; and nothing could induce her to take a second pill. She believed them to be powerfully aperient, and purging took place. Hence a wide field is opened for the charlatan and the quack; while the experienced practitioner often finds, that in many ailments he will in vain prescribe remedial agents, unless he acquire the confidence of the patient.

The connexion of one disease with another is a subject of great importance, and of much interest to the practical physician. We rarely find that a patient has died free from all disease, except the one which has been the immediate cause of death; such a case, is, indeed, exceptional. It may be that an acute inflammation of the lungs has led to fatal results, whilst chronic disease may have been going on in the abdomen, the heart, or the brain, perhaps quite independently, but having an important influence on the curative or non-curative condition of the disease: chronic disease creeps along with unobserved step, till some acute affection proves fatal. This relation of disease is worthy of our consideration, in studying the affections of the alimentary canal; and we may find that the diseased conditions arrange themselves in the following manner:—

1. They take place simultaneously in the same body, without any connexion, as mere coincidents.

2. The connexion may be that of different manifestations of the same disease in its progressive action, rather than a really different diseased condition.

3. One disease may have important modifying or predisposing influence upon another.

4. Several organs may be affected simultaneously by one exciting cause.

5. One disease may be antagonistic of another ; and,

6. Diseases or abnormal conditions are conservative, the one from the other.

I might enumerate many instances of these associations, in diseases of the nervous system, or of the thoracic viscera, but must content myself with a few illustrations from disease of the abdomen.

1. As instances of coincident disease we may mention the following :—A patient who had been employed in working lead, and was affected with severe colic, was partially relieved ; but he suddenly had intense collapse, and died with all the symptoms of perforated intestine. We found, on inspection, that there was in the stomach a large chronic ulcer, and at its base a minute perforation, which had extended into the peritoneal cavity. A child affected with chorea was relieved by sulphate of zinc, and was about to go home, when it was seized with severe dysentery during the time that cholera prevailed ; and the little patient died in three days. In another patient who lately died under my care, affected with phthisis, we found a large hydatid cyst close to the kidney, in addition to advanced degeneration of the lungs. These diseases could not be looked upon as cause and effect, and are probably correctly regarded as coincidents.

2. As manifestations of the same disease in progressive action, and which ought not to be considered as two but as one disease, we may enumerate the sympathetic vomiting which we find in hydrocephalus, and in diseases of the kidney and uterus ; the diarrhœa in albuminuria, which follows an anasarous condition of the mucous membrane, and the constipation in diseased spine ; and the extension of strumous disease affecting one

organ or viscus after another, as of the intestine in phthisis. So, again, the severe neuralgic pain in the parietes of the abdomen, simulating colic, but arising from disease of the spine, may be only the early manifestations of the spinal disease, though preceding its more marked indications by several months or years. Numerous instances might be adduced in the progressive symptoms of spinal disease, or of valvular disease of the heart, or of chronic disease of the lungs.

3. One disease predisposes to, or modifies another disease. Thus, in affections of the lungs, of the heart, and of the liver, the circulation through the vena portæ may become exceedingly impeded, and the whole of the mucous membrane engorged and turgid with blood; in this state, a slight exciting cause will set up distressing flatulence and distension of the abdomen; chronic catarrh of the mucous membrane is produced, or hæmorrhage, or ulceration. A patient in incipient phthisis, with tubercles or slight ulceration in the mucous membrane of the intestine, is exposed to cold and wet, to hardship and miasm, and very severe diarrhœa or dysentery is set up; whilst his friend, who has had no such predisposing cause, escapes, though exposed to the same exciting influence.

The instances which Dr. Budd has deduced, of abscess in the liver following ulceration of some part of the tract of the canal which supplies the vena portæ, are also illustrations of one diseased state exciting another; here it does not follow as a necessary sequence, or a continuance of the same diseased action, but a new disease is produced.

Again: a strumous subject, after recovering from typhoid fever, may become affected with tubercular disease of the intestine. The previous exhaustion has rendered the patient, already of feeble power, subject to another disease; and the typhoid ulceration of the intestine is sufficient to excite the manifestation of its action: these are by no means rare occurrences.

A sailor was admitted into Guy's, a few years ago, with Asiatic cholera. He died, and in his colon a large circumscribed ulcer was found, about the size of a crown piece, and covered by a slough, with adherent cherry-stones; the presence of such irritation and inflammation in the colon would render him more amenable to an attack of the disease, although it would not produce it.

A young man fell into the Thames, and afterwards was seized with diarrhœa; he was shortly attacked with typhoid fever, and admitted into Guy's. He quickly died; the dysenteric diarrhœa rendered the fever more severe in its character; and perhaps was the immediate cause of the fatal termination.

4. Two diseases sometimes arise from the same exciting cause, or rather two organs become affected: thus, acute inflammation of the colon sometimes comes on with pneumonia. Of these cases we shall speak more fully in our remarks on dysentery.

5. Diseases are in some instances antagonistic. Cancerous disease and struma appear to be in this relation; or it may be that they are so diverse in their mode of operations that they cannot exist together; an excess of formative action, although quite abnormal, being incapable of existing with a deficiency. We sometimes find in cancerous disease of the abdomen, chronic disease of some of the mesenteric glands,—contracted, and calcareous, with old strumous change; such evidently indicates a mode of action which has given place to another of a different kind; and the same kind of deposit is occasionally found in the lung in cancerous disease, which has proved fatal.

6. Disease may be conservative in its character. We have many instances of this in the abdomen. A chronic ulcer of the stomach is prevented oftentimes by adhesions from perforating the peritoneal sac, so that the liver, or the pancreas, forms the base of the ulcer. So, again, in ulceration of the ileum and colon, in disease of the cæcum, and in gall-stone, adhesions prevent extravasation, or limit it after it has taken place. Many instances of this kind might be adduced in which life has been prolonged by these means.

These associations of disease have an important bearing on the correct diagnosis, and still more on the prognosis, of disease; they may oftentimes serve to explain its intractable character, as well as to account for the different effect of remedies under apparently similar circumstances. This complication of disease should place us on our guard in making close observation of every sign which presents itself to us, and should lead to a strict inquiry into the history of the patient, and the previous ailments to which he may have been subject.

CHAPTER II.

ON DISEASE OF THE ŒSOPHAGUS.

THE simple functional character of the Œsophagus, as the communication between the mouth and the stomach, and the rapid transit of food over its mucous surface, render it less subject to organic disease than some other parts of the alimentary tract. The contiguity of the Œsophagus in the posterior mediastinum, with important thoracic viscera and vessels, occasionally leads to secondary Œsophageal disease; more frequently, however, primary Œsophageal disease extends to the surrounding structures in the chest.

The pharynx is the organ of deglutition; and dysphagia, as applied to the difficult performance of its function, arises not only from disease of the pharynx, but also from derangement of the several cavities and canals communicating with it; thus, the pharynx is continuous by the fauces with the mouth, by the posterior nares with the cavities of the nose, by the Eustachian tubes with the ears, by the superior opening of the larynx with the respiratory organs, and by the commencement of the Œsophagus with the rest of the alimentary tract; and disease of any of these contiguous parts may interfere with the freedom of pharyngeal action. The process of swallowing can scarcely be said, however, to be fully completed till the food is lodged in the stomach; and hence, dysphagia becomes one of the most prominent symptoms of disease, not only of the pharynx, but also of the Œsophagus.

The causes of dysphagia are, therefore, various, and may be divided as follows:—

I. Disease of the structures connected with the first part of the act of deglutition, affecting the tongue, the tonsils, the fauces, &c.; as in cynanche tonsillaris, scarlet fever, syphilitic,

strumous and cancerous ulceration, acute and diphtheritic inflammation.

II. Diseases of the pharynx, as diffused inflammation of the cellular tissue, local suppuration, sometimes in connexion with disease of the spine, ulceration, whether strumous, syphilitic, or cancerous, acute and diphtheritic inflammation.

III. Diseases of the larynx, as in laryngitis and croup, ulcerations of the epiglottis or laryngeal cartilages, whether strumous, syphilitic, or cancerous.

IV. Functional and spasmodic stricture of the Œsophagus and pharynx, as in hysteria, hydrophobia, &c.

V. Paralysis of the muscles of deglutition.

VI. Inflammation of the mucous membrane of the Œsophagus.

VII. Mechanical injury and poisons.

VIII. Structural obstruction of the Œsophagus.

1. Constrictions.

2. Ulcerations, sometimes communicating with the larynx.

3. Cancerous disease.

4. Pressure of aneurismal or other tumours.

I. *Dysphagia arising from disease of the tongue, tonsils, and fauces*, as we find in glossitis, in cynanche tonsillaris, in acute and diphtheritic inflammation, in scarlet fever, in strumous, syphilitic, and cancerous ulcerations, needs only to be mentioned in connexion with disease of the Œsophagus.

II. *Pharyngeal Dysphagia*.—Acute and diphtheritic inflammation of the pharynx, and ulcerations of a specific and non-specific character have dysphagia as an almost constant symptom.

In the acute stage of diphtheria deglutition is generally difficult; but, after the subsidence of the more urgent symptoms, a peculiar form of paralysis is occasionally developed, and dysphagia results from the loss of power of the pharyngeal muscles. This is referred to in cases described by Dr. Gull and Dr. Kingsford, and is very fully dwelt upon by Maingault, in his valuable treatise on 'Diphtheric Paralysis.' The latter author states that the symptoms of paralysis of the soft palate are "a nasal voice, pain in deglutition, difficulty or impossibility which

the patients experience in exercising suction, in distending the cheeks, in blowing with the mouth, and in gargling ; and, on examining the throat, we find immobility of the soft palate, a lengthened condition of it, numbness, absence of pain, and a diminution or loss of special sensibility. These various phenomena supervene a longer or shorter time after the acute affection of the throat has been cured ; and when the pain having already completely disappeared, and the deglutition having become easy, the convalescence is apparently established." Maingault gives the following table to show the relative frequency of the different forms of paralysis which come on after diphtheria :—

Paralysis of the lower limbs	13
General paralysis	64
Paralysis of the soft palate	70
Disordered sensibility without muscular weakness	8
Amaurosis	39
Strabismus	10
Paralysis of the muscles of the neck and trunk .	9
Anaphrodisia	8
Paralysis of the bladder	4
Paralysis of the rectum	6

He also adds, that "dysphagia never fails to come on whenever diphtheria is followed by general paralysis." In the treatment of these cases a very generous diet is necessary ; and, if deglutition be impossible, an œsophageal tube should be used ; electricity has been found of great service, and tonic remedies, as quinine, steel, &c. ; others speak very favorably of the efficacy of strychnia, and of nux vomica. In very many instances recovery slowly takes place, but a fatal result occasionally ensues from this complication to a disease terribly fatal in its primary effect. Dr. Gull* has recorded an instance of this kind in a child aged 12 ; and he attributes these forms of secondary paralysis to the extension of disease to the spinal nerve centres. In the young patient just referred to, drooping of the head came on five weeks after the attack of diphtheria. The child could utter no sound, and the diaphragm was unmoved in respiration,

* 'Lancet,' July, 1858.

indicating a loss of power in the phrenic nerves. Fearful attacks of suffocative dyspnœa were produced when the head was moved in particular situations, and in one of these paroxysms he died.

Dysphagia is also the result of a local suppuration behind the pharynx. Bleuland* mentions, fatal dysphagia "a collectione puris inter spinæ vertebrarum corpora atque inferiorem pharyngis superioremque œsophagi partem;" and Dr. Fleming† narrates several interesting cases of this kind. One of them took place in a strumous child aged 11, in whom there was tenderness at the central part of the cervical vertebræ; the tonsils were enlarged, the voice was nasal and muffled, and the muscles spasmodically affected; those at the back of the neck were rigid, and in speaking, the contraction of the labial and nasal muscles produced a sort of tetanic expression; the jaws could be slightly separated, and the tongue could be suddenly protruded. The abscess was opened by a bistoury, and the child recovered. A still more remarkable instance occurred in an infant aged two months; a peculiar snuffle, difficulty in deglutition, so that the child could scarcely suckle, with occasional dyspnœa, were followed by convulsion, and a semi-comatose state; by pressure of the finger against the swollen part in the pharynx the abscess ruptured; its contents were discharged through the nose, and the infant recovered.

Local suppuration of the pharynx produces sudden and urgent dyspnœa, with dysphagia, accompanied also with febrile disturbance; the respiration, however, is less impeded in these cases than in primary disease of the larynx. If suppuration have taken place below the level of the soft palate, a projecting tumour is observed on examination of the throat. The diagnosis is then sufficiently evident; and, when it is possible, puncturing the abscess relieves the urgent symptoms. Abscesses of this kind arise from disease of the vertebræ, or from local sources of irritation.

Diffused suppuration is generally the result of erysipelas, of pyæmia, sometimes of diphtheria and of scarlet fever. The patient rapidly passes into a typhoid condition, the dysphagia

* 'Bleuland de sanâ et Morbosâ œsophagi structurâ.'

† 'Dublin Quarterly Review.'

- becomes extreme, the respiration impeded, and, on examining the neck, we find either erysipelatous redness of the skin, or a fulness and tenseness among the infra-hyoid muscles, impeding the free movement of the parts concerned in deglutition. The examination of the neck will generally enable us to distinguish the dyspnoea arising from this cause, from that produced by disease of the larynx, or trachea, as well as from that consequent on pressure or injury of the nerves of respiration.

In my notes I find the following case, a very interesting one of the kind, occurring in 1847 :—

CASE I. *Diffused Inflammation of the Throat.*—Abraham S—, æt. 36, a sailor, of intemperate habits, was admitted into Guy's Hospital, October 13th, 1847. On the 5th, whilst unloading coals, he received a blow on the back of the hand, and on the following day rigors ensued, with pain in the axilla, but the skin of the arm did not become inflamed. On admission, on the 13th, he presented the appearance of a man suffering from typhoid fever; there was delirium at night; the pulse was very soft, 106; the tongue was moist, and the respiration was much oppressed; no fluctuation could be found under the pectoral muscle, nor any suppuration detected in the neck; the wound on the hand was dry. Stimulants and opium were administered. On the 15th, the respiration was difficult and laboured, 42 per minute; there was evident obstruction of the larynx, and some tenderness about it, but scarcely any swelling; and neither fluctuation nor suppuration could be detected on very careful examination; there was also great difficulty in swallowing. On the 16th, the respiration and deglutition were somewhat easier, but the skin was clammy, and the tongue dry. He died on the following day, after vomiting some blood. On inspection, the whole of the cellular tissue surrounding the muscles of the neck was found infiltrated with pus, but there was no suppuration below the pectoral muscles.

CASE II. *Diffused Inflammation of the Throat. Ulceration of the Pharynx.*—In this case, which was admitted in May, 1847, a woman, aged 66, had sore throat, with pyrexia, quickly followed by typhoid symptoms, and death on the fifth day. On inspection, suppuration was found among the muscles of the neck, which extended round the Œsophagus, as low as the root of the lung. In the pharynx there were several superficial ulcers; and one opposite the arytenoid cartilage had extended into the cellular tissue.

These diseases appear to be an erysipelatous form of inflammation, and are generally of such an aggravated kind as to be quite beyond the reach of remedial measures. Hot fomentations should be used, and ammonia with stimulants administered; it is rare, however, that the suppuration in these cases is sufficiently localized to admit of relief by incisions.

III. *Laryngeal Dysphagia*.—In acute laryngitis and in croup, it is the exception to find deglutition performed in the normal manner; and sometimes, especially in laryngitis, dysphagia is an urgent symptom. In these cases, however, the dyspnoea and cough are the earlier and the more marked indications of disease. The epiglottis is found to be injected and tumid; and the branches of the superior laryngeal nerve are rendered intensely sensitive. This abnormal sensibility is an explanation of the dysphagia which is generally present, even if the mischief do not spread directly to the pharynx.

Diseases of the laryngeal cartilages rarely extend to the pharynx, unless the malady be of a cancerous character; more frequently, as in necrosis, suppuration takes place among the muscles of the neck, and chronic laryngitis of a most intractable form is produced. But although dysphagia is not a prominent symptom of disease when only the cartilages of the larynx are affected, the reverse is the case when the fibro-cartilage, the epiglottis, is also implicated, whether in syphilitic, in strumous, or in the cancerous diseases.

In syphilis, both the glossal and laryngeal surfaces of the epiglottis become involved, and sometimes nearly the whole of the fibro-cartilage is destroyed leading to distressing dysphagia; and in phthisis ulceration of the epiglottis is one of the most trying complications of the complaint, the ulceration extending on its inner surface as far as the margin, which becomes eroded and gradually destroyed, and the contact of food with this irritated surface sometimes leads to the instant rejection of it through the nares. In chronic phthisis I have seen this condition attributed to organic disease of the Œsophagus itself, on account of the extreme urgency of the dysphagia, and because the food appeared to have passed below the pharynx before it was forcibly ejected. It sometimes happens that solids are more easily swallowed than fluids; and this is the case in some instances where the dysphagia arises from disease of the larynx; a solid will pass over the diseased epiglottis and fall beyond it, whilst a fluid comes in close contact with it.

Laryngeal dysphagia is often greatly mitigated by the inhalation of steam, or of the fumes from conium or stramonium. In less severe cases astringent gargles, and in syphilitic ulcera-

tion the application of a strong solution of nitrate of silver, afford relief by diminishing the extreme sensibility of the diseased surface. Counter irritation may occasionally be applied with advantage, as the tincture of iodine, hot fomentations, cantharides, &c. The improval of health by residence at the sea coast, by cod-liver oil, by preparations of iodine, and of steel, and by the use of such forms of nutritious diet as can be taken by the patient, are the means most likely to be followed by enduring benefit.

IV. *Spasmodic Stricture of the Œsophagus*.—The few cases of this kind which have come under my own observation have been in young women of an excitable character, who were suffering from leucorrhœa, or painful menstruation, and impaired digestion. The strongest language was used by these patients to express their inability to swallow, and they showed the greatest unwillingness even to make the attempt. One of them was a young woman about twenty-three years of age, thin and imperfectly nourished. On passing an œsophageal bougie, no obstruction whatever was found; she afterwards swallowed food in small quantities, which was increased day by day until she took the usual amount. Fright, terror and cold are found to produce spasmodic dysphagia, and instances of this kind we have seen after violent storms of thunder and lightning; but in lesser degrees it is not unfrequent in hysterical subjects, and as the symptoms of hysteria are well marked there is no danger of mistaking the complaint for cancerous obstruction, although there may be some difficulty in distinguishing it from perforating ulcer extending into the trachea.

Abercrombie * gives the case of a lady, æt. 40, in whom stricture of the œsophagus was supposed to exist. The symptoms continued for a year, and were entirely relieved by passing "an egg-shaped silver ball attached to a handle of silver wire."

This state, however, is not limited to one sex. Mayo† mentions a remarkable instance of spasmodic stricture of the œsophagus in a gentleman, æt. 60, who had sudden obstruction of the œsophagus whilst at dinner. It was relieved by the passage of a

* Abercrombie on 'Diseases of the Stomach.'

† Mayo, 'Outlines of Pathology.'

bougie. The brother of the patient, who had suffered from gout, had had a similar seizure. The same author narrates a case of spasmodic stricture in a young man, produced by ulceration of the interior of the larynx. It must be borne in mind also, that spasmodic contraction of the œsophagus tends to increase the obstruction arising from organic causes, so that the degrees of dysphagia in the same case differ exceedingly.

The general symptoms and history aid us in the diagnosis of these cases: thus, there is an absence of emaciation; the attack comes on suddenly after a slight cause, as from nervous shock or slight catarrh; there is freedom from pain, but nervous excitement is always present.

Hot fomentations, the use of fluid, instead of solid food for a short time, aperient or antispasmodic enemata as of turpentine or rue, will afford relief in these cases. Tonics are often of service, as the compound iron mixture with decoction of aloes, or the compound steel pill, with aloes and myrrh; so also quinine, zinc, valerian, vegetable tonics may be used; and the shower bath, good air and exercise, and cheerful occupation of the mind will greatly assist in the restoration to health.

Bougies are often employed, but their use is not generally beneficial, and may be detrimental by tending to perpetuate and aggravate a state of spasmodic irritation and contraction; but in cases where the muscles of the pharynx have lost their contractile power, the direct introduction of food is absolutely required. In some hysterical patients, the refusal to swallow arises from a disordered will, rather than from any disease in the œsophagus itself.

In *Hydrophobia* true spasm of the pharynx and œsophagus is present in a most marked degree. A few years ago a case of this terrible disease occurred at Guy's Hospital, and on inspection the appearance of the pharynx was very peculiar, its cavity appeared more than twice its natural size; the constrictor muscles were retracted to the utmost, and the fauces were exceedingly large from the rigid contraction of the soft palate; the spasmodic condition of the muscles rendered every part of the pharynx as dilated as possible; whilst from similar muscular spasm the œsophagus was contracted. The mucous membrane of these parts was injected and covered with

tenacious mucus. There was great congestion of the membranes of the brain and spinal cord; and the lungs were in a similar state of engorgement. The other viscera were healthy; but there was emphysema of the cellular tissue of the neck. The symptoms during life indicated extreme irritability of the nerves supplying the pharynx, as, indeed, of all the branches of the fifth and pneumogastric nerves.

CASE III. *Hydrophobia*.—The patient was a young man, æt. 23, who was said to have been bitten by a dog nine years previously. On the day of admission into Guy's, May 15th, 1854, difficulty of swallowing came on, with great mental excitement. He was removed to one of the adjoining work-houses, and afterwards brought to the hospital, about nine o'clock in the evening. He was a strong muscular man, and at first sight appeared to be affected with acute mania, or delirium tremens; but there was a sudden starting, especially when a draught of cold air came in contact with his face, which clearly indicated the character of the disease. This starting evidently resulted from spasmodic action of the muscles of the face and pharynx; his countenance had a wild and excited aspect; he thought that he was being murdered, that boiling water was dropping upon his face, and he said, that he felt "choked." The pulse at nine o'clock was 90, and at eleven, it was 120; the tongue was clean; the pupils were widely dilated; the face was bathed with perspiration; and the hands clammy; he would not attempt to drink, but dashed the cup away from him with a violent spasmodic action, but he ate a small portion of bread; and he was frequently spitting out saliva. Restraint was required, for in his terror, which was fearful to witness, he rushed at the window, and would have seriously injured himself. I remained with Dr. Gull for several hours during the night with this patient, a witness of one of the most fearful spectacles of misery and disease I have ever seen. About 12:30, an injection, containing 10 grains of cannabis indica, was administered; but the whole of the enema was at once returned. At one o'clock a longer tube was passed, and the same quantity again injected; the paroxysms had then become very violent and frequent, and the pulse exceedingly small, irregular, and occasionally intermittent, 120 to 130 per minute. At 2:15, he was still more violent, calling out as loudly as his strength would permit. It was then determined to administer chloroform. During the inhalation, intense congestion of the eyes and face came on; the pupils became much smaller, and the pulse a little more perceptible; the respiration, which had been catching and accompanied with gasping and sighing, became more regular. In four or five minutes after leaving off the chloroform the paroxysms began to return; the face, however, did not at once become sensible to impressions. Chloroform was administered three times during the hour, and on leaving it off the same return of paroxysm took place; the pulse became almost imperceptible, and the respiration more stertorous. About 3 a.m., whilst under the influence of chloroform, 10 grains of cannabis indica were placed in his mouth; it became

mixed with saliva, but was entirely ejected. He died at 3.30 a.m., from exhaustion and apnoea.

V. *Paralysis of the muscles of deglutition* is generally observed immediately to precede those of respiration; and is looked upon, correctly, as a common sign of approaching death. The nervous centre of the function of swallowing is close to that of respiration, and there is an intimate connexion between them. Where there is loss of the power of deglutition, the placing of fluid in the mouth will be followed by its entrance into the larynx, or by violent cough, or it may even hasten death.

We not unfrequently, however, observe, in cases of cerebral disease, when the muscles of the tongue are paralysed, that the swallowing, especially of solids, becomes exceedingly difficult. This difficulty arises from the movements of the tongue being restrained, for the bolus of food cannot be formed, and pushed back into the fauces; fluids are more easily swallowed, because more readily brought within the action of the true muscles of deglutition. The introduction of nourishment by an elastic tube has been the means of prolonging life; it is probable also that the muscular fibres of the Œsophagus have diminished contractile power in these cases.*

Another class of cases are those connected with mental disease, some of which may easily be mistaken for true paralysis. With great feebleness of muscular power, we may find that the will is unable to excite muscular action; that the muscles of the pharynx appear paralysed, because they are not stimulated to healthy contraction, and hence deglutition cannot be performed. The following interesting case, admitted under my care into Guy's Hospital in July, 1856, is of this kind.

CASE IV. *Dysphagia. Mania.*—A. B— was an emaciated man, æt. 60, of a dingy and sallow appearance, a gas-fitter, who had resided at Deptford. His wife stated, that for several years he had suffered from attacks of extreme irritability; but that his only complaint was of pain in the region of the transverse colon. On the 18th July, he appeared to lose the power both of speaking and of swallowing, having previously said, that "he did not know what was coming over him." On the 23rd, he was brought to Guy's

* Abercrombie, 'On Diseases of Brain,' and 'On the Stomach.'

Hospital; he was then prostrate, and unable to stand, but could slowly move his legs and arms; his countenance was not without intelligence, and he appeared slightly to understand questions; he could not protrude his tongue, which remained almost motionless at the floor of the mouth, and was dry on its surface; fluids put into the mouth were retained for a short time, and then ran out again at the angles of the lips, but he could not be induced to attempt to swallow; placing a teaspoon at the back of the mouth excited some action of the muscles; the pupils were active, the right was rather larger than the left; the pulse was 56, and compressible; the heart's action was very feeble; the respiration was normal, 20 per minute, but the air could scarcely be heard to enter the chest. The abdominal muscles were exceedingly rigid, but the abdomen was not distended. Half a drop of croton oil was placed on the back of the tongue, and afterwards, a nutrient enema was administered. On the 24th, my colleague, Mr. Cooper Forster, passed an œsophageal tube into the stomach without any difficulty, and without meeting any obstruction; some beef tea thickened with arrow-root was in this way administered; the patient afterwards swallowed milk and beef tea, &c., with less and less difficulty, and on the third day began to speak; the bowels were acted on by castor oil, and by enemata; ammonia and calumba, with a small quantity of wine, were given on account of his prostrate condition; his tongue lost its brown and furred appearance; and he rapidly improved. His mind, however, was not in a clear state, for as soon as he was able to eat, he had the idea that no other patient in the ward had any food. This case closely resembles some of those found in lunatic asylums; but this patient was unable to make the attempt to swallow; a condition which might easily have been mistaken for paralysis of the muscles themselves. I have since learned that after leaving the hospital he became violently maniacal.

VI. *Inflammation of the mucous membrane of the œsophagus* is described by some authors,* and dysphagia is mentioned amongst its symptoms; but excepting those cases in which irritant substances have been swallowed, and excepting, also, instances of gastro-enteric inflammation, as in children, we have not seen any patient in whom acute œsophagitis could be said to exist; nor, in the examinations after death have we been more successful in finding the indications of ordinary acute disease affecting this part of the alimentary tract. A case is recorded by Abercrombie in which a soft adventitious membrane could be traced from the pharynx to the cardia, and the pharynx and epiglottis were covered with a similar membrane. The patient, a gentleman æt. 26, had swelling of the side of

* Galen—Bleuland 'de sanâ et morbosâ structurâ œsophagi;' Mondière, 'Arch. Gén. de Méd.,' tom. xxx.; 'Dict. de Méd. et de Chir. Pratique;' Copland, 'Med. Dict.'

the neck, hoarseness, some dyspnœa, dysphagia, and febrile excitement, followed by typhoid prostration. The throat was red, and aphthous crusts were observed on the mucous membrane. Death took place in three weeks.

In a case of acute inflammation of both the small and large intestines of a diphtheritic character, admitted under my care into Guy's Hospital, during 1855, the mouth was inflamed, and the pharynx and tonsils were covered with a white film, spread upon an injected mucous membrane. This white film consisted of a beautiful torula, interlacing in all directions, constituting the '*muguet*'; it extended downwards to the commencement of the œsophagus, and some traces of it were found in that canal. In this instance, the symptoms were those of dysenteric diarrhœa, which had come on several months before her admission into Guy's, and had persisted without any intermission for seven weeks. The disease was attributed to her removal into a damp house. The patient was exceedingly prostrate; she had severe vomiting, and whenever she attempted to take food, retching and severe pains were produced. No medicines nor injections had any effect in checking the diarrhœa, and she died on the third day after admission. The mucous membrane of the œsophagus was, perhaps, equally affected with that of the mouth and pharynx; and, indeed, it appeared that the whole tract of the alimentary canal from the mouth to the rectum was inflamed.*

It is probable, that in some of the cases of severe gastro-enteritis in children, in whom the mouth as well as the intestine is evidently inflamed, the whole of the alimentary tract is affected, and would present before death a condition quite abnormal. At the close of chronic disease, we find a similar condition of the pharynx, rendering deglutition both painful and difficult; aphthous inflammation of the mouth having extended into this part. In these conditions, I have not seen any remedy followed by such beneficial effects as the chlorate of potash, associated sometimes with borax and honey; but alone it often acts apparently in a most marked manner. This remedy in stomatitis, introduced I believe by Hunt, was very extensively used by the late Dr. Golding Bird, and subsequent observers

* (See *dysentery*, for a more full account of this case.)

have confirmed the opinion which he entertained. It appears to act partly by its local effect, and also as a saline after its absorption into the system.

Local suppuration in the submucous tissue of the œsophagus sometimes takes place, and produces dysphagia, or rather the regurgitation of food, as in organic obstruction. These instances are rare and obscure in their character, and the febrile disturbance is not sufficiently distinctive of the nature of the affection; diseased glands may set up this local abscess, as well as the annular stricture, to which we have presently to refer.

VII. *Dysphagia from destruction of the mucous membrane by mechanical or chemical agents.*

Every year the lives of many children are destroyed by drinking *boiling water*; vesication of the mouth, fauces, and pharynx is at once produced; but generally, the instantaneous rejection of the scalding fluid prevents any portion from being swallowed. The margin of the epiglottis and the entrance of the larynx are, however, often injured; and the sudden swelling of the mucous membrane of the glottis preventing the entrance of air into the lungs, leads to a rapidly fatal result.

The prognosis in infants after accidents of this kind is very unfavorable; for, although the respiration may be apparently unimpeded several hours after the injury, a slight increase of effusion of serum into the swollen mucous membrane of the larynx leads to complete occlusion of the already diminished rima, and causes death from apnœa. The operation of tracheotomy has in not a few instances saved life, but in many the supervention of acute disease of the trachea and bronchi has proved quickly fatal. In several inspections after death, the mucous membrane of the lower part of the œsophagus and stomach have presented considerable congestion, shewing that some of the hot water had reached those parts; and in a case of poisoning by corrosive sublimate, lymph was found in the œsophagus.

The action of *corrosive poisons* may be divided into that which is immediate or primary, and that which is remote or secondary. The mucous membrane of the mouth, fauces, pharynx, and œsophagus becomes at once discoloured by the

chemical action of the poison, and it assumes a yellowish white or brown colour, according to the strength and character of the agent; if the poison be in a very concentrated state, the membrane is charred and destroyed. The effusion of a sero-albuminous fluid into the mucous membrane, or into the cellular tissue, leads to considerable swelling; but, as with boiling water, the epiglottis rarely escapes injury; its margin often presents an eroded and serrate appearance. The longitudinal rugæ of the œsophagus, especially near the stomach, are discoloured or destroyed; shreds of œsophageal membrane are sometimes ejected, and in one instance,* after strong sulphuric acid had been taken, a complete cast of the œsophagus was thrown off. The stomach also is sometimes extensively injured, its mucous membrane and coats being charred, or even perforated, and the adjoining viscera similarly acted upon wherever the poison comes in contact with them. If the patient survive the immediate action of the poison, a fibro-plastic product is effused into the submucous tissue; thickening and contraction of the new product takes place, and in this manner an annular constriction may arise.

Symptoms.—The first symptom produced by taking corrosive poisons is severe pain of a burning character in the mouth and throat, and along the whole tract of the œsophagus. When any portion has been swallowed it is succeeded by vomiting of dark coloured fluid, containing blood; extreme dysphagia quickly follows, and speaking is exceedingly painful; there is also sometimes urgent dyspnœa. The lips are often injured by the contact of the poison; the tongue is found to be swollen and injected, and the mucous membrane of the whole mouth is discoloured, becoming yellowish white or brown, or completely charred; and the throat is in a similar condition. Cough and dyspnœa are distressing symptoms, if the epiglottis and larynx have been injured. Diarrhœa is occasionally present. The patient has an anxious and dejected countenance; the pulse is compressible and feeble; and sensibility continues till death, which may take place in a few hours. But, in other cases, it is often remarkable how soon the injured mucous membrane of the mouth and throat recover themselves; the vomiting sub-

* Mayo's 'Outlines of Pathology.'

sides; the extreme pain in the throat produced by swallowing or coughing disappears; and in a few days the patient appears almost convalescent, and gives hope of recovery. If only the mucous membrane of the stomach be injured, there may be entire immunity from pain; but the freedom from pain is a very deceptive symptom, the patient often unexpectedly dies, from syncope, or asthenia, although more generally the subsequent contraction of the fibro-plastic product in the œsophagus, stomach, or pylorus leads to obstruction, slow emaciation, the regurgitation or vomiting of food, and a lingering death from inanition.

The first effect of these poisonous agents is best combated by the use of substances capable of neutralizing their chemical properties, combined with oleaginous and demulcent drinks to shield the mucous membrane. Opiates may be given to relieve pain; and as soon as deglutition can be performed, bland and nutritious diet may be allowed.

When secondary contraction of the œsophagus has taken place, the probability of relief is slight; fluid forms of nutriment administered in a concentrated form, and injections of a similar kind may prolong life. Dr. Cumin* succeeded in saving the life of a child by the use of elastic catheters, after potash had been taken; and Mr. Foster,† after the same poison, sought a like result, though less successfully, by the formation of a gastric fistula.

CASE V. Poisoning by Sulphuric Acid—Death on the 11th day.—In an interesting case of poisoning by sulphuric acid, in October, 1855, in which death did not take place until the eleventh day, the mouth and throat were of a whitish colour; at the posterior part of the mouth, there was considerable injection of the mucous membrane, and on each side of the posterior pillar of the fauces there were whitish loose patches of membrane. The edge of the epiglottis was found minutely eroded, and the mucous membrane of the œsophagus was pale and covered with yellow membranous flakes. The prostration and collapse immediately following the reception of the poison were accompanied by vomiting of grumous blood, but in less than twelve hours the patient was able to swallow some milk and arrow-root; and on the fourth day appeared to take her food without difficulty. Death took place from the sloughing condition of the mucous membrane of the stomach, com-

* Dr. Cumin. 'Transactions of the Medical and Chir. Soc.:' Edin.

† 'Guy's Hospital Reports.'

bined with inflammation of the duodenum, and of the whole tract of the intestine. The ability to swallow in this case was restored in a very short time, considering the fearful injuries which resulted to the whole of the mucous membrane. (See more full account of the state of the stomach in our remarks on that viscus.) Case XL.

Two instances admitted into Guy's Hospital in 1857, illustrate the primary effects of poisons on the pharynx and Œsophagus. They are detailed in the Reports for 1859, by Dr. Wilks :

CASE VI. "*Poisoning by Soap Lees.*—Charles T. C——, æt. a year and a-half, was admitted under Mr. Hilton, on September 4th, 1857, at six o'clock in the evening. About an hour before, the child had drunk from a cup about a mouthful of soap-lees; some oil and mucilaginous fluids were administered, and he was brought to the Hospital. The child was then very ill, and, in the course of an hour or two, some difficulty of breathing came on, but this did not appear sufficiently extreme to warrant tracheotomy. The most marked symptom after this was an intense heat of skin. The child died at five o'clock on the following morning, twelve hours after swallowing the fluid.

"*Post-mortem appearances.*—The mouth and tongue were slightly excoriated, and of a light brown colour. The fauces, tonsils, and mucous membrane of the pharynx had a slightly swollen appearance, and had a yellowish-brown hue. The whole of the Œsophagus presented a similar condition, the mucous membrane having a brownish colour, particularly the longitudinal rugæ. The membrane was changed in character by the alkali, but was nowhere destroyed. The greatest pernicious effect had been produced at the very extremity of the Œsophagus, where the interior was of a dark-brown colour; this terminated at a defined line, the mucous membrane of the stomach immediately below being quite unaffected. The stomach was contracted. It was found, on opening it, quite empty; the rugæ were well marked, and the whole mucous membrane had a slightly pink hue, being more than usually injected. These appearances were, however, so slight, that, unless especially looked for, they would probably have been disregarded. As before stated, the termination of the Œsophagus was of a dark brown colour, but this terminated abruptly at its margin. Near the pyloric end of the stomach, near its greater curvature, there were a few rugæ of a very dark colour, produced, no doubt, by the action of the alkali. The mucous membrane, thus altered, was not at all soft, nor could it be stripped off; but, on the contrary, was hard, and had a horny feel. The duodenum was healthy. The larynx, at its top, was almost closed by the greatly-swollen epiglottis, the enlargement being due to an effusion of serum within it; the glottis itself was only slightly swollen; and upon raising the epiglottis, and looking into the larynx, the passage was seen to be quite free; neither the vocal cords, nor any other part, having been touched. The lungs showed some lobules in the first stage of inflammation. The heart was healthy, and firmly contracted."

CASE VII. "*Poisoning by Sulphuric Acid.*—William V—, æt. 56. The patient's mind was not perfectly sound, and, therefore, the account he gave of himself was received with some doubt. When he was admitted, on the evening of October 28th, 1856, he walked up stairs to his bed, and did not appear very ill, although he was dejected, and did not speak much. He stated that he had been to a friend's house; and there, by mistake, drank about a dessert-spoonful of oil of vitriol. His mouth was of a brown colour, but not excoriated. Magnesia and milk were given him. On the following day, and also on the third, he appeared depressed; but he was not otherwise ill, and it was thought, from the mildness of the symptoms, that he would recover. On the fourth day, however, he died rather suddenly, or at least, unexpectedly.

"*Post-mortem examination.*—The body was that of a strong, muscular man. A yellow fluid, of acid reaction, ran from the mouth. The brain was not quite healthy. The mucous membrane of the mouth was of a yellow colour; but when this yellow epithelial layer was removed, the mucous membrane left was healthy. The front part of the tongue was also discoloured, but not the back. The Œsophagus throughout was of a yellow colour. The mucous membrane was only affected in the most prominent ridges, but the walls of the organ were swollen to three times their natural thickness. This was due to a sero-albuminous exudation into the submucous tissue. The top of the larynx was also slightly swollen in the same manner. The stomach appeared natural externally, and was of usual size. Upon opening it, it was found to contain about a pint of a bright yellow fluid. The mucous membrane was especially affected at the pyloric half of the stomach. The fundus, in which the fluid was found lying as usual, had only a yellow tint like the Œsophagus, and the mucous membrane was softened; but, towards the middle of the stomach, the whole of the pyloric half of the interior was of a black colour, and raised up in projecting masses or ridges, which were in a sloughing condition, and would soon have been cast off. This black matter consisted of carbonized and decomposed mucous membrane, with blood within it. The whole coats of the stomach were soft, and readily tore. The charring of the stomach ended at the pylorus, but about two inches of the duodenum were of a purplish colour, and the rugæ were blackened; below this, the intestines, both small and large, were unaffected. The small contained a yellow matter, similar to that in the stomach. The contents of the stomach were not acid, nor was any of the poison discoverable. The heart was healthy, and contained a firm, decolorized fibrinous clot on the right side, not acid in its reaction."

The following case illustrates the *secondary* effect of a corrosive poison in the thickening of the whole of the Œsophagus and obstructed pylorus, which led to a fatal termination, in a man who died three months after having taken an ounce of nitric acid.

CASE VIII. "*Poisoning by Nitric Acid.*—James T—, æt. 24, was ad-

mitted under Dr. Barlow's care, in March, 1852, in a state of extreme emaciation; more than two months previously he had taken about an ounce of nitric acid, and had completely swallowed it before he discovered the fatal mistake. The primary effects gradually subsided; but vomiting after food increased, and he steadily lost flesh and strength; he vomited, with some pain, all the food which he swallowed; the abdomen sometimes became extremely distended; the bowels had only been opened twice during the two months preceding his admission; the tongue was injected. He lived eighteen days after admission. On inspection, the epiglottis appeared healthy; the mucous membrane of the whole of the Œsophagus was thickened and readily separated; the submucous tissue and all the coats of the Œsophagus were also thickened; the stomach was enormously distended, reaching to the anterior superior spinous process of the pubes; the pylorus was obstructed, thickened, and contracted; the lungs and heart were healthy; the liver was small, deep in colour, and the gall-bladder contained about 3jss. of dark-coloured bile; no other viscus was diseased.

CASE IX. Poisoning by Nitric Acid. Recovery from the Primary Effects.
—A young man, æt. about 22, a hawker, whilst at his tea, on March 13th, took by mistake for vinegar, a mouthful of nitric acid, and swallowed it. A severe burning pain in the mouth was at once produced, which extended to the epigastrium. A druggist prescribed an emetic; vomiting then came on, and he brought up about half a cupful of blood. The vomiting continued through the night, and on the following day he was brought to Guy's; the countenance was anxious; the mouth and tongue were stained of a yellow colour, the tongue enlarged and injected; the throat was intensely injected, and presented irregular shreds of whitish membrane upon it. He was unable to swallow, and speaking produced cough and much distress in the throat. He stated, that he suffered pain in the throat and epigastrium when retching came on, but not when quiet. He was a muscular man, and in health at the time of the accident. Milk and eggs were given, and magnesia mixture with tincture of opium \mathfrak{m} v. every four hours. On the 17th, he was sitting up, taken food, and he stated that he felt much more comfortable; he had slight pain in the throat when he swallowed, but had no other discomfort. The throat was still very much injected, and sloughy mucous membrane was separating. In a few days he left the hospital, and considered himself well.

The immediate effects of the poison were in a few days relieved, and the dysphagia disappeared; but after such severe injury to the Œsophagus we should look with great caution to the result; thickening of the coats and constriction may, and perhaps will, follow; the acid probably reached the stomach, for we had pain produced at the scrobiculus cordis on vomiting; but there was no evidence that serious injury had been done to that viscus.

VIII. *Organic obstruction of the œsophagus* arises—1st, from annular constriction; 2nd, from ulceration, sometimes communicating with the trachea; 3rd, from cancer; and 4th, from the pressure of aneurismal or other tumours. The history of the symptoms will alone enable us to distinguish these cases from each other. In some of them it is to be hoped that remedial means may be used which have hitherto scarcely been attempted; in others, it is evident that nothing can be done for cure, but the pain and the severity of the symptoms may be mitigated.

Annular constriction of the œsophagus consists in the effusion of fibrinous material into the submucous cellular tissue; the new tissue contracts, and becomes exceedingly dense, forming a firm constricting band, whilst the tube above dilates, and the obstruction increasing, at last the passage of food becomes impossible. A beautiful specimen of this form of constriction is shown in the Preparation No. 1789, in the Museum of Guy's. Many of these cases are probably the result of corrosive poisons, or they arise from injury to the mucous membrane, or from inflammation set up by adjoining disease. No. 1789⁵⁰, in the Guy's Museum, shows the œsophagus of a lad, in which about two inches above the diaphragm the parietes of the tube are thickened, the mucous membrane is contracted, and is apparently cicatrized for the space of an inch and a half. There is an absorbent gland in the neighbourhood of the stricture, adherent to the walls of the canal. The boy had had difficulty of swallowing from infancy, and a bougie had been passed occasionally. Inflammation about the gland had, perhaps, led to this thickening of the canal.

An exceedingly interesting case, bearing some analogy to the preparation just mentioned, was recorded by Dr. Ogier Ward, in the Transactions of the Pathological Society for the year 1850:

“A boy, æt. 10, died rather suddenly of meningitis of the base of the brain, extending down the theca vertebralis. He had enjoyed good health till two months before his death, when he struck himself against a post, and after that time he was subject to sick headache. He was ill a week with pain in the chest and head immediately after the accident, and afterwards, for about a month, he complained of pain in the chest, and a difficulty of swallowing the first few mouthfuls at his meals, which, he said, seemed to stick in his throat; but this complaint ceased before his last illness. On post-mortem examination the œsophagus was found dilated and deprived of its epithelium opposite the bifurcation of the bronchi, where there was a

longitudinal ulcer, one and a-half inches in length, through the mucous membrane, with two small orifices at its centre communicating with each other, and by a canal three and a quarter inches long, with an enlarged bronchial gland filling up the interval between the bronchi. This gland was in a partial state of suppuration, but no pus flowed along the canal on pressure."

In recorded cases of annular stricture, the obstruction has gradually increased in severity, and unless we have a history of poison having been taken, or of the discharge of pus from an abscess, we know of no direct symptom by which this form of obstruction can be distinguished from that arising from cancerous disease. The passage of a bougie may reveal to us the presence and position of obstruction, without indicating its true character, but the mucous from the bougie should be examined, and may sometimes guide to a correct diagnosis.

Ulceration.—In the Museum at Guy's Hospital, there are several specimens showing ulceration of the Œsophagus, of a non-cancerous character, extending into the trachea, and there is some obscurity as to their correct pathology. Difficulty of deglutition was the most prominent symptom during life in these cases; in some, dysphagia had been gradually developed, in others, deglutition had suddenly become impossible. The patients had complained of pain at the sternum or between the shoulders; and on attempting to swallow urgent dyspnoea came on, and the forcible ejection of food through the nares took place. The patients became emaciated; and life was prolonged for a short time by the use of nutrient enemata. On inspection after death, the only disease found has been a perforation of the Œsophagus opening into the trachea; the openings, two or three in number, extended over one or two inches, their edges were smooth, and without any thickening; and in several cases the opening into the trachea was smaller than that into the Œsophagus. The examination of these cases does not give any evidence of cancerous disease; the early symptoms appear to arise from the Œsophagus, the difficulty in respiration following that of deglutition; nor do we find other evidence of disease, either in the larynx or in the lungs. These facts appear to show, that the disease has not commenced either in the mucous membrane of the trachea nor in disease of its cartilages; and we are led to

suppose, either, that an abscess has formed between the œsophagus and trachea, and led to fistulous openings into those canals, or that ulceration has taken place in the œsophagus, and gradually extended in depth through the adjoining structures. It sometimes, however, happens that ulceration extending into the œsophagus arises from disease of the tracheal cartilages, and the following remarkable specimen is of that kind:—

CASE X. *Diseased Cartilages of the Trachea. Ulceration of the Œsophagus.*
—A carrier, æt. 42, at Hampton, came under the care of Mr. Holleston and Mr. Jepson, in 1853, for crowing respiration, with abundant expectoration, but no very urgent dyspnoea, nor difficulty in swallowing. He gradually sank; but six months before his death, he expectorated a portion of ossified tracheal cartilage (Preparation 1711⁸⁷), and six weeks later a second portion. On inspection, at the commencement of the œsophagus, immediately beneath the cricoid cartilage, a vertical opening, half an inch in length, with smooth and rounded edges, was found to extend into the trachea; there were three other communications, resembling fissures, merely separated from each other by shreds of mucous membrane. (See Preparation 1711⁸⁷.) The cartilages of the trachea were ossified, and there was ulceration of the mucous membrane of the larynx at the cricoid cartilage. The inferior lobe of the right lung was consolidated, but no other part of the body was diseased; and there was no trace of cancerous nor of strumous disease.

Dysphagia was almost absent, as far as can be learned, in this case; and the symptoms were those indicative of disease commencing in the larynx; thus differing remarkably from the cases presently to be recorded, where dysphagia was the most prominent complaint of the patient. It is probable that their pathology is also different. No history of syphilis is given, but the expectoration of a portion of diseased cartilage, six months before death, indicated the character of the disease.

CASE XI. *Ulceration of the Œsophagus. Perforation of the Trachea.*
—A married woman, æt. 24, who had never enjoyed robust health, about a year previous to her admission under Dr. Barlow's care, had enlarged glands of the neck, which diminished under the use of iodine; and six months afterwards she began to experience difficulty in swallowing, with pain in the chest, uneasiness in the throat, and some shortness of breath. These symptoms increased in severity till admission, but a short time previously they had suddenly become very much aggravated. She was much emaciated; no swelling could be found about the neck, nor disease detected in the chest. She experienced the greatest difficulty in swallowing fluids, and food was at once forcibly ejected. Mr. Hilton passed an œsophageal tube; and found

that when the patient breathed, air passed from it, indicating a communication with the trachea. She was fed for six weeks before death entirely by injections. On inspection, the trachea and œsophagus were found extensively diseased from the level of the cricoid cartilage, nearly as far as the bifurcation of the trachea, and the two canals communicated by three openings. The anterior wall of the œsophagus near its commencement was destroyed, with the exception of two slips of muscle, which still remained; and at this part there was an oval ulcerated opening passing into the trachea; below this a small portion of the calibre of the œsophagus was considerably contracted; still lower the œsophagus was again destroyed, and two more openings passed into the trachea. At this latter part, the posterior wall of the œsophagus was also destroyed, and the body of the last cervical vertebra was exposed; so that an abscess had been formed bounded by the cellular tissue of the trachea, by the remains of the œsophagus, and by the muscles of the neck. The openings into the trachea were oval, transverse, perfectly smooth, and covered with mucus; and not the least thickening nor heterologous deposit could be detected by careful examination, aided by the microscope. In the ovary, and in an adhesion on the surface of the liver, there were slight strumous granular deposits; the other viscera were healthy, and there was no evidence of cancerous disease. There was difficulty in breathing, and Mr. Hilton performed tracheotomy, but without any permanent advantage to the patient. (See Preparation 1714¹⁰, and Drawing 246²⁴.) The stomach was small, contracted, and almost perpendicular; it contained a small quantity of bilious-looking alkaline mucus. The large intestine was dilated, and contained healthy fæces; the cæcum contained some acid mucus; the rectum presented several small ulcers, and was covered by a firmly adherent diphtheritic deposit.

The following case occurred in Guy's, in the year 1840. There is no history of the symptoms on record; but the patient was a man æt. 33, and he died four days after admission. The post-mortem inspection was as follows:—

The body was exceedingly emaciated. Near the middle of the œsophagus, the mucous membrane, for about two inches, was of a very red colour, and irregular from ulceration; the canal was much contracted, and would have scarcely admitted the end of the little finger. Below the stricture, the œsophagus was much dilated, and an abscess had formed behind it, containing four ounces of dark fluid of a sour odour; there was a small sinus leading to the abscess; the mucous membrane, both above and below the diseased part, was quite healthy; there was no evidence of cancer in the affected part; nor was any other organ diseased, except that the kidneys were found to be granular. (See Preparation 1789²⁷.) It was supposed, that a corrosive poison must have been taken, but of this there was no evidence.

An exceedingly interesting record will be found in the 'Patho-

logical Transactions' for 1852, by Mr. W. Trotter. A young woman, in St. Mary's Hospital, æt. 25, had ulceration of the œsophagus, which extended into the pericardium, and led to sudden syncope and death. For three months she had had nausea, dysphagia, occasional vomiting, and pain at the top of the sternum, and at the epigastrium. Solids were swallowed with much difficulty. There was found, after death, simple ulceration without contraction; the ulcer had extended from the bifurcation of the trachea nearly to the diaphragm, and had perforated the pericardium. No other organ was diseased.

The last two cases were instances of simple ulceration below the bifurcation of the trachea; in the others the disease was above this part; still, they appear very similar in character, and the modification in the symptoms arose from the difference of the adjoining structures which were implicated.

There are many instances of pain at the upper part of the sternum on swallowing, when no trace of pressure nor aneurism can be found; and I have seen this symptom disappear under the use of tonics, sometimes with iodide of potassium. The idea of cancerous growth has been precluded; and it has been therefore a question whether some abrasion of the mucous membrane, or slight ulceration, such as we sometimes find in the pharynx, had not led to this complaint.

It is exceedingly difficult, during life, to decide as to the character of these cases of simple ulceration: the emaciation, dysphagia, and distress, being the same as in cancerous disease. In all the cases which have come within my notice, the age of the patient has been very much less than in most of those of cancer. This alone, however, is not sufficient to enable us to decide with certainty as to the character of the disease.

The treatment is exceedingly unsatisfactory; the spasmodic contraction of the ulcerated part prevents the passage of œsophageal tubes; no food can be swallowed, and the administration of nutrient enemata prolongs life only for a few days or weeks. It is painful to find, after death, that simple ulceration of the œsophagus, or a fistulous communication with the trachea, is the only existing disease; and that if food could have been introduced beyond this point, life might have been prolonged. The operation of œsophagotomy is a very difficult one, and in

many of these cases, if performed, would be quite ineffective, because the disease is often situated at the root of the lung, or behind the first bone of the sternum; in either case, the operation could not be performed below the seat of stricture. It having been found that the peritoneum may be divided without fatal result, and without the terrible effect seen to follow from ruptured viscera, the propriety of forming a gastric fistula in some of these cases is worthy of very serious consideration. It appears certainly warrantable, as it would afford a chance of life to those who have only the prospect of certain death. In the human subject, several cases of gastric fistulæ accidentally produced have been recorded, and the experimenters on animals purposely make such openings, under the influence of chloroform, without the production of severe peritonitis.

Since the first edition of this work was published, a gastric fistula has been made in two cases; the first was suggested by myself to relieve the agonising distress of starvation in a man dying from cancerous occlusion of the œsophagus. (See cases of cancer.)

Syphilitic Ulceration of the Œsophagus.—In the 'Dublin Quarterly,' of Feb. 1860, Mr. J. West describes two cases of dysphagia occurring in young girls, in whom the dysphagia gradually increased, and at last led to a fatal result; four inches down the œsophagus there was constriction arising from fibrous deposition in the submucous tissue; disorganization of the lungs had also taken place. Mr. West believed the disease in each case to be of syphilitic origin; but I have never witnessed any case of obstruction of the œsophagus that could be positively ascribed to this cause.

In Cancerous disease of the Œsophagus and Pharynx the symptoms are very similar to those mentioned as occurring in ulceration of the œsophagus; the patients are generally beyond the middle period of life, and difficulty in swallowing, which gradually increases in severity is the first and most prominent symptom. In some instances, however, the dysphagia does not become extreme till the extension of the cancerous ulceration to the lungs, or to other structures, leads to symptoms, which almost mask the original disease; vomiting, or rather regurgitation of the food is always present in a greater or less degree. There is

also pain at the sternum, in the back, and sometimes in the upper part of throat, of a dull or burning character; but its intensity varies greatly. Dyspnœa comes on when the trachea or bronchi are involved, and is occasionally associated with loss of voice, especially in those cases in which the disease is situated at the base of the pharynx, and extends into the larynx. The dysphagia and emaciation increase, and after six or seven months the disease generally proves fatal.

In organic obstruction of the œsophagus, and especially in that of a cancerous kind, the dysphagia is very peculiar; the patient hesitates in making the attempt to swallow, he takes a considerable time, as it were, to prepare the muscles of the pharynx for the effort, and when the attempt is made, the food or fluid is at once rejected; and sometimes severe suffocative cough is produced.

Mr. Travers* thus describes cancerous disease of the pharynx and œsophagus, "Scirrhus strictures followed by ulceration and cancerous fungus are met with in the pharynx and the top of the œsophagus in elderly persons, chiefly *females*, in my experience. They are productive of constant nausea, dry burning sensation in the throat and stomach, difficult breathing, frequent spasms and alarms of suffocation, and excessively impeded deglutition; upon the gentlest introduction of the finger or bougie, hæmorrhage follows, which afterwards becomes spontaneous. The patient has a faded sallow countenance, a disturbed circulation, and is emaciated to a skeleton."

Hæmatemesis is sometimes a symptom of cancer of the œsophagus. Dr. Bristowe exhibited at the Pathological Society a specimen of ulceration of the œsophagus, extending into enlarged veins, and causing fatal hæmorrhage; and in another instance recorded by him, the superior intercostal artery was opened by cancerous ulceration. The hæmorrhage produced by the extension of cancerous ulceration is occasionally repeated several times in the progress of the disease, it indicates its progressive character, and is often the precursor of a fatal result. Dr. Balding, at the same Society in 1857, showed an ulcer of the œsophagus of doubtfully cancerous character, in

* 'Med.-Chir. Trans.,' xv, p. 252.

which a sloughing cavity connected with the œsophagus had also formed a communication with the right subclavian artery. The walls of the aorta are often partially injured, and in some cases perforated, thereby leading to sudden and fatal hæmorrhage, as in Case XXXII.

Sometimes the lymphatic glands are enlarged in the neighbourhood of the cancerous mischief, and it is well always to examine the neck and axilla for evidence of glandular disease. Pressure on the bronchi from similar enlargement leads to greater distinctness of respiration in one lung than in the other; this, however, is also a sign of aneurismal, or, in fact, of any kind of tumour exerting direct pressure on adjoining structures, and is only of corroborative value.

In cancerous obstruction of the œsophagus we do not generally find much distension of the canal above the seat of the disease; for the ulcerated surface leads to excessive irritability, and food is very quickly regurgitated; or a state of spasmodic contraction equally favours the instantaneous rejection of ingesta. Dr. Alderson, however, in writing on Carcinoma of the Œsophagus, mentions that "immediately after deglutition, there is a remarkable bulging out or protrusion of the œsophagus above the strictured point; it is, in fact, a bag or pouch, which is formed by the effort of the patient to swallow a larger quantity of food than the œsophagus, in its natural state, can contain." In annular obstruction of a non-cancerous nature, the canal becomes more enlarged and dilated; the disease is of a more chronic character, and there is less sensibility of the surface than in true cancer. *Monro** mentions an instance in which as much as a pint of fluid could be retained for ten minutes. This general dilatation of the œsophagus above an obstruction must be distinguished from pouches connected with the canal, and presently to be mentioned.

The part of the œsophagus which is most fixed by its connexion with other organs, is at the root of the lungs; and it is there that cancerous disease is most frequently found, and not at the termination of the canal, although, occasionally, the disease extends downwards, so as to reach the diaphragm.

* *Monro* 'On Morbid Anatomy of the Gullet, Stomach, and Intestine.'

This localization of the disease is, possibly, on account of the canal being less yielding opposite the root of the lung, and irritation being, consequently, more easily set up.

Epithelial cancer is the most frequent form of disease, but medullary and scirrhus cancer also occur. In numerous instances, the growth has presented modifications of epithelial scales; some cells had very large nuclei, other growths showed large nuclei thickly set together, or brood cells. In Case XVI, some papillæ were observed on the surface of the growth, covered by healthy squamous epithelium, and containing a capillary filled with blood. Another papilla presented a similar general appearance, but the capillary was filled with granular cells, somewhat resembling white corpuscles of blood. A third papilla closely resembled some of the brood cells; its central portion contained nuclei and nucleated cells, and it was surrounded by flattened scales or cells resembling epithelium. It appears probable, that in some cases degeneration of papillæ may lead to the formation of these clusters of cells, rather than endogenous growth, which is the method usually received. The disease generally extends by mere contiguity of structure, involving the adjoining bronchial glands, at the roots of the lungs, and thereby encroaching upon the bronchi; pneumonia is thus frequently set up, and if sloughing occur gangrene of the lung follows. I have several times found the pneumogastric nerves destroyed on one or both sides; and this destruction of nerve supply induces congestion of the lungs, which is followed by pneumonia, without actual extension of disease into the lung passages.

The glands and other viscera are rarely implicated in epithelial cancer; but Case XV, presents a marked instance of the disease existing in other structures besides the primary organ affected, cancerous elements being discovered in the liver, pancreas, stomach, supra-renal capsules, &c. In the liver, lungs, and pancreas, cells of an epithelial character, and precisely similar to those found in the ulcerated œsophagus, were observed. In this case, also, there was chronic pneumonia existing, not alone, but associated with deposit of small cancerous tubercles. In some cases, a doubt might be felt as to the cancerous character of these tubercles, but such doubt

did not occur in this instance. The vomiting and pain were exceedingly severe, more so than usual, but were explained by the condition of the pneumogastric nerve, which was exposed at the base of the ulcer; several of its branches were truncated, and others passed across the surface in a perfectly exposed state. Cancerous infiltration also extended around the right semilunar ganglion, and encroached upon its component tissues. All the gland-structures in the abdomen were more or less atrophied; but whether this atrophy was due to the diseased condition of the ganglion, or merely resulted from the exhausting disease, and the advanced life of the patient, is doubtful. Mr. H. Gray records a case of villous and epithelial cancer at the termination of the pharynx, in the 'Pathological Transactions' of 1855; and at the termination of the œsophagus colloid cancer has been observed.

In Case XIII, the disease was perfectly local, but its structure more closely resembled medullary than epithelial cancer.

Sometimes the cancerous ulceration extends through the diaphragm after destroying the œsophagus, as in Cases XXVIII, and XXIX. In the former the œsophageal opening into the stomach remained; but a large sloughing cavity was formed, bounded by the pancreas, spleen, and diaphragm, which communicated with the posterior mediastinum by an opening in the diaphragm. Immediately behind the pericardium was a large sloughing cavity, presenting above the truncated œsophagus and pneumogastric nerves, and terminating below as just described. It was surprising that the patient could have lived as long as he did, but only three days before death he had come from Broadstairs, and was not at all aware of his perilous condition. In the latter case there was remarkable absence of dysphagia.

Mondière in 'Arch. Gén. de Méd.,' tom. xxx, mentions from Keppelhont a case in which ulceration of the cardia and œsophagus communicated with an abscess of the liver; and also from Dr. Anesant, an instance of scirrhus ulceration of the inferior part of the œsophagus, rendered fatal by extension to the spinal cord.

The fatal issue arises from several causes; but these may

generally be divided into two. 1st. Inanition, the dysphagia having become complete; and 2ndly, from the extension of disease to the lung or the surrounding tissues. The character of the disease of the lung deserves our especial attention. In only 2 instances out of 21 did death result from simple inanition, and even in these the lungs were not altogether free from disease.

In 11 cases there was pneumonia, *one* was acute *and* chronic, the pneumogastrics being free; *two* were chronic pneumonia, with sloughing of the œsophagus, in one of which there was cancerous tubercles in the lungs, in the other communication between the trachea and œsophagus. In *seven* the pneumonia was acute, 3 from direct extension of disease to the lung, and 4 from injury to the pneumogastric nerves. *One* was strumous pneumonia with cancer of the pharynx.

In 4 cases there was gangrene of the lung.

In 1 there was acute bronchitis and laryngitis.

In 1 there was pleurisy and disease of the kidneys.

In 1 there was cancer of the lung with extreme congestion.

In 2 death took place from inanition; in one of these, however, there was chronic bronchitis, in the other lobular pneumonia.

In 1 fatal hæmorrhage took place from perforation of the aorta, but the lung was also involved.

As to the causes of the pneumonia,—1st. The pressure or destruction of the pneumogastric was followed by acute pneumonia on the same side, or by gangrene. 2nd. The pneumonia appeared to result from the extension of disease into the bronchi setting up, if not pneumonia, acute bronchitis or laryngitis. 3rd. The sloughing of the cancer was followed by septic changes in the blood, and consequent inflammation of the lungs. 4th. Cancerous growth or tubercles in the lung acted as the cause of congestion or inflammation. 5th. Strumous disease of the lung already existed.

The chronic affection of the lung is interesting in its relation to cancer; in Case XV, the tubercles were of a cancerous character, and set up chronic pneumonia; in Case XXVI, in which cancer of the throat existed, the lung was affected with ordinary strumous disease, and the most careful examination could not detect any trace of carcinomatous product in the lung; it is probable that the disease in the lung existed prior to the development of the cancer, and was scarcely concerned in the cause of death.

In Case XXVIII, the vomica at the apex of the lung was evidently of a chronic character; the dense iron-grey lung tissue around it, and calcareous degeneration, indicated that it had existed for a considerable period.

In Case XXIX, there was also a vomica at the apex, and the history indicated that cough had existed long prior to the dysphagia. There were evident signs of phthisis in the flattened apex, loud bronchial and amphoric respiration and bronchophony; had there not been present the cancer of the Œsophagus, it would have been considered as an ordinary instance of pneumonic phthisis. In the exhaustion which was consequent on the obstruction of the Œsophagus, the cough continued troublesome, and a few days before death acute disease of the lung was set up, arising, perhaps, at the time the cancerous growth began to disintegrate, or from atmospheric changes.

Of the 21 cases here recorded, 15 were men, and their average age was nearly 56½; 6 were women, and their average age 46½; in two of the latter, the pharynx rather than the Œsophagus was affected; the statistics from such a small number are, however, of comparatively little value.

Among these instances, the longest period which elapsed between the commencement of dysphagia and death was about two years, several were three to seven months, and in two still less, the interval being only 5 and 7 weeks.

The *diagnosis* is sometimes obscure; this has been mentioned in reference to annular stricture, and perforating ulcer into the trachea. Where we find chronic disease of the lung with dysphagia, the diagnosis is much increased in difficulty, because in ordinary phthisis, the dysphagia is sometimes exceedingly

severe. This remark applies especially to the bronchitic phthisis of advanced life, as in Case XXX, where the patient was sixty-nine years of age.

Dysphagia with chronic emaciation very often arises from cancerous disease; but, sometimes the dysphagia is very slight, and the sudden onset of acute secondary disease masks the primary mischief; and, again, very extensive sloughing of the œsophagus may render the rejection of food from the stomach almost impossible. It is sometimes very difficult also to distinguish between cancerous disease of the œsophagus and pressure upon the tube by aneurismal or other tumours; in the latter instances the dysphagia is less persistent; and often varies according to the position of the patient, the œsophagus falling from an aneurismal tumour of the aorta, as the patient leans forward; paroxysms of dyspnœa are also frequently present in cases of arterial disease. See Case XXXIV.

The *prognosis* is in all these cases very unfavorable; but in some, after the avoidance of irritating and solid food, or after the use of nutrient enemata for several days, the dysphagia becomes diminished in a marked degree, the patient is able to partake of solid food, and we may be led to take a more favorable view of the case than is warranted by the nature of the malady. Two cases admitted into Guy's Hospital with symptoms of cancerous disease of the œsophagus, men nearly sixty years of age, with nearly complete dysphagia were so much relieved as to leave the hospital; when, however, we find the disease extending into the respiratory passages, or into the large vessels, we may fear a speedy and fatal termination.

These remarks suggest to us the proper mode of *treatment*. The most bland and unirritating diet should be given, as milk, eggs, jellies, soups, &c. Solid articles of food should be abstained from, at least for a time, for the attempt to swallow solids produces distressing spasm of the œsophagus; and if the dysphagia be very severe, nutrient injections should be administered, so as to allow complete rest to the diseased structures. Stimulants and cod-liver oil afford partial relief, and check the progress of disease.

Solution of potash and iodide of potassium, with vegetable infusions, afford relief in the earlier stages of the complaint;

so also, nitric and hydrochloric acids, with morphia or opium. In advanced cases, where there is extensive cancerous ulceration and excessive irritability from exposure of the branches of the pneumogastric, internal remedies are of no avail, and nutrient injections are the only means of prolonging life.

Opium in one form or other is the best remedy for the secondary pneumonic or bronchitic complications; to give mercurials and antimony, &c., is to exhaust still more rapidly the already ebbing life of the patient.

It is a question of great importance, how far bougies may be used with advantage in the treatment of organic obstruction of the œsophagus. The frequency of communication between the œsophagus and the trachea or bronchi, the occasional entire destruction of the canal, and the injury which sometimes results to the walls of the aorta, are each of them serious objections to its use, see Case XXXII; and I have very frequently seen instances in which a bougie would certainly have passed into the bronchus, and led probably to instant death. In the earlier stages the bougie may usefully serve to indicate the precise character and seat of the disease; but should always be used with extreme caution. The cauterization of stricture of the œsophagus was resorted to about the close of the last century; and Sir Everard Home, in his 'Practical Observations on the Treatment of Strictures of the Urethra and Œsophagus,' records several instances, in which the use of caustic bougies was followed by relief of the severe dysphagia. Simple bougies have been more frequently used, often with benefit, though sometimes to the injury of the patient, as in a case mentioned by Mr. Fletcher, in his 'Medical and Surgical Observations,' in which perforation was produced at the termination of the pharynx, and suppuration among the muscles of the neck followed.

As an instance of the beneficial use of the bougie, I may refer to the following case:

A patient recently under my care in Guy's Hospital, suffering from severe dysphagia, complained of pain about the level of the sternum. He was forty-three years of age, by trade a letter carrier, and for twelve months the symptoms of obstruction in

the Œsophagus had troubled him ; but he had not found that the symptoms had generally much increased since the time when he first noticed them. The dysphagia sometimes became greater, but it was always with great difficulty that any portion of solid food could be swallowed. He was a spare man, no enlargement could be felt in the neck, no obstruction could be found in the pharynx, nor was there any evidence of pulmonary, cardiac, or arterial disease. Several attempts were made to pass bougies into the stomach, but without success ; they rested about the level of the sternum ; even an elastic catheter could not be introduced beyond that point. After a few days I requested the patient to abstain altogether from food, and fed him for one week entirely by nutrient enemata ; at the end of that time a large bougie could be easily passed into the stomach ; the rest to the canal had allowed the irritation and spasmodic constriction to subside, and, although organic stricture still existed, solids could be more easily swallowed. The bougie met with an obstruction, as before said, about the level of the sternum, and about three inches below the first, a second obstruction occurred, indicating probably the upper and lower limit of the diseased surface ; and when the bougie reached the constriction the coughing of the patient enabled the instrument to pass onwards. Cod-liver oil and a nourishing fluid diet were given, and the patient left the hospital relieved.

Cases of Cancer of the Œsophagus showing their Complications and Terminations.

- CASE XII.—James R—, æt. 45. Sloughing Pneumonia. Pneumogastric involved.
 „ XIII.—John R—, æt. 50. Communication with the Trachea. Pneumonia. Diseased Kidneys.
 „ XIV.—George E—, æt. 73. Gangrene of the lung. Cancer of the cervical gland and thyroid body.
 „ XV.—Jane B—, æt. 63. Cancer of the Stomach, Liver and Pancreas. Chronic pneumonia with cancerous tubercles. Destruction of the pneumogastric nerve. Granular kidneys. Diseased semi-lunar ganglion.

- CASE XVI.—Charlotte W—, æt. 32. Disease at the termination of the pharynx. Secondary laryngitis.
- „ XVII.—Catherine S—, æt. 38. Communication with the trachea. Cancer of the lung and of the kidney.
- „ XVIII.—Mrs. B—, æt. 54. Death from inanition.
- „ XIX.—George D—, æt. 45. Gangrene of the lung.
- „ XX.—John H—, æt. 66. Pleurisy and diseased kidneys.
- „ XXI.—James M—, æt. 49. Communication with the trachea. Gangrene of the lung.
- „ XXII.—James S—, æt. 57. Extension into the lung. Gangrene.
- „ XXIII.—John D—, æt. 64. Pleuro-pneumonia. Extension into the lung.
- „ XXIV.—John D—, æt. 71. Pneumonia. Pneumogastric nerve involved.
- „ XXV.—Richard E—, æt. 63. Pneumonia. Pneumogastric nerve involved.
- „ XXVI.—Martha M—, æt. 81. Cancer of the palate and pharynx. Strumous pneumonia.
- „ XXVII.—William E—, æt. 50. Pneumonia. Pneumogastric nerve involved.
- „ XXVIII.—George W—, æt. 53. Communication with the left bronchus. Pneumogastric involved. Old vomica in the lung. Extension of disease through the diaphragm.
- „ XXIX.—John H—, æt. 45. Pneumogastric nerves truncated. Sloughing extending into the lung and through the diaphragm.
- „ XXX.—William G—, æt. 69. Chronic and acute pneumonia. Vomica. Pneumogastric nerves free.
- „ XXXI.—William H—, æt. 47. Inanition. Slight lobular pneumonia. *Gastrotomy*.
- „ XXXII.—Margaret H—, æt. 60. Perforation of the aorta. Sudden fatal hæmorrhage.

CASE XII. *Cancer of the Œsophagus. Sloughing Pneumonia; the Pneumogastric Nerve involved.*—James R—, æt. 45, was admitted into Guy's Hospital, November 21st, 1854, under Dr. Gull's care; and died November 30th. He was a married man, a labourer, and intemperate in his

habits. For nine weeks prior to his admission he had been unable to swallow food with comfort, and he had suffered from severe pain at the lower part of the sternum. From that time he lost much flesh; and cough, with pain in his side, came on. He vomited occasionally, and had burning pain at the sternum; and there was a sense of nausea, when he began to eat. On admission, he had a cachectic, pale, and wretched appearance; he was troubled with cough, and the expectorated matters were exceedingly offensive. At the apex of the left lung the respiration was coarse; and at the base of the right lung there were signs of consolidation; the voice, both at the base and apices, was increased in resonance. Cinchona and morphia were administered, but the patient sank in a few days. The severe pulmonary symptoms in this case completely masked the original disease of the œsophagus; for a short time it was believed that the case was one of pneumonia with old disease of the lung, and that the burning pain at the sternum, and vomiting, were the consequence of his former intemperate habits. At the commencement of the œsophagus extensive ulceration was found on inspection; the ulcer was four or five inches in length, irregularly tubercular on its surface; and several tubercles were situated in the mucous membrane, both above and below the ulceration. The disease extended as low as the root of the lung, but the lungs themselves and the pleura were free from cancerous disease. The tissue external to the œsophagus was extensively infiltrated, especially on the right side, and some of the bronchial glands were affected; the right pneumogastric nerve extended through the diseased structures. The lower part of the pneumogastric appeared wasted, but it could not be traced satisfactorily throughout, having been divided in the inspection. The right lung, at its lower lobe, was of a greenish colour, and it had a faint gangrenous odour; it was infiltrated with dirty serum, and was imperfectly consolidated. The bronchi were intensely congested. The remaining parts of the lungs and the larynx were healthy. The heart, stomach, liver, and intestines, &c., were also healthy; and no cancerous disease could be detected in any other part.

As to the character of the growth, it had the general and microscopical appearance of epithelial cancer. There was no direct communication between any of the large bronchi and the ulceration of the œsophagus; and it appeared probable that the right pneumogastric, becoming involved in the disease, had predisposed to the pneumonic inflammation on the same side. The disease proved fatal at an earlier period than usual, for the patient died ten weeks from the recorded commencement of difficulty in swallowing; and the diagnosis was rendered obscure by the extreme severity of the pulmonary symptoms.

CASE XIII. Epithelial Cancer of the Œsophagus communicating with the Trachea. Pneumonia. Granular Kidneys.—John R—, æt. 50, was admitted

into Guy's Hospital, under Mr. Hilton's care, March, 1856, and died the following day. No history, except that he had felt ill for three months, could be obtained. On admission he was suffering from urgent dyspnœa; there was great congestion of the face, and he was apparently dying from apnœa. Tracheotomy was performed by Mr. Callaway, but the patient died in a few hours. The body was spare but muscular. The epiglottis was pale, and its mucous membrane slightly œdematous; near the arytenoid cartilage was a small circular ulcer. At the commencement of the trachea was the artificial opening, and two inches and a quarter below it, and an inch above the bifurcation, was a vertical opening about half an inch in length, extending into the œsophagus; the membrane around the perforation was of a dull grey colour; and the mucous membrane of the trachea and of the bronchi was much congested. Nearly three inches above the commencement of the œsophagus was an ulcer, three inches in length, with raised, irregular edges, and with a ragged surface; at its base was the opening into the trachea; the remaining part of the canal was healthy. (Preparation 1793.*^m) Neither bronchial nor cervical glands were affected. The lower lobe of the right lung was in a state of red hepatization, becoming grey; and in the left lung was a lobule broken down from acute disease. On the right pleura there was effusion of lymph. The kidneys were small, granular, contracted, and contained cysts. The other viscera were healthy.

As in the last case, there was no affection of the glands; the patient was one in whom the constitution was impaired by chronic disease of the kidneys. When admitted he was too ill to allow of proper investigation being made; but from the supposition that the dyspnœa arose from the larynx, tracheotomy was performed.

CASE XIV. *Cancer of the Œsophagus, of the Cervical Glands, and of the Thyroid Body. Gangrene of the Lung.*—George E—, æt. 73, was admitted into Guy's Hospital, November, 1853, in an extremely emaciated state, and died February, 1854. He was a table-cover maker, and in his early life had been intemperate. Eight months before his admission he received a severe fall, from which he never recovered; and two months later he began to suffer great pain in eating solids, and he had occasional attacks of vomiting. These attacks became more and more frequent, and latterly almost incessant. He could not take solid food, and complained of intense pain at the cardiac extremity of the stomach. Mr. Callaway passed an œsophageal bougie, but without meeting any obstruction in its passage. His vomiting diminished soon after admission. The bowels became constipated, and he continued to suffer severe pain. He became gradually weaker, and on February 5th, he vomited a considerable quantity of dark-coloured fluid; he died on the 13th. Inspection was made twenty hours after death. In the brain there was considerable sub-arachnoid effusion, and disease of the arteries. At the central part of the œsophagus, opposite the

root of the lung, there was a large, irregular ulcer, two inches in length, which involved the whole of the tube; at the upper part was a raised circular margin, and a partially detached ulcer of similar character, about half an inch in diameter. At the root of the right lung there was a mass of sloughing tissue, which was infiltrated with sanious fluid, and the adjoining lung was consolidated. At the base of the left lung there was a circumscribed mass of pulmonary apoplexy and lobular pneumonia, and a vomica containing thin purulent fluid. The cervical glands and the thyroid body were infiltrated with carcinomatous product, white, and of a medullary character. There were fatty and fibroid degeneration of the heart. In the peritoneum were old adhesions, and a granular condition of the surface of the liver. The kidneys were also granular and contracted.

Although the œsophagus was extensively ulcerated in this case, a bougie could easily be introduced; showing that spasmodic contraction was the principal cause of the obstruction, and of the rejection of the food. In this instance also, the bougie might easily have passed through the diseased walls of the œsophagus into the posterior mediastinum, or into the pleura; and, in a diagnostic point of view, it might have led to the supposition, that the disease was at the stomach, rather than in the œsophagus, on account of the absence of obstruction to its passage. As we have previously stated, great care must be exercised in the use of these instruments.

CASE XV. Epithelial cancer of the Œsophagus, Pancreas, Liver and Kidneys. The pneumogastric nerves involved. Granular kidneys. Chronic pleuro-pneumonia, with cancer. Fibrous tumour in uterus. Cancer of suprarenal capsules and semilunar ganglion.—Jane B—, æt. 63, was admitted Aug. 23rd, 1855, under Dr. Addison's care. She was married, of light complexion, was thin and anæmic, and had resided at Snow's Fields; she had had three children. She had suffered for nine months; and the first symptom was pain after swallowing; no tumour could then be felt, but cancerous disease was suspected. After admission, a firm mass, at the scrobiculus cordis, about the size of a hen's egg, could be felt; it was well defined, sensitive on pressure, and tolerably distinct pulsation could be perceived, the food returned at once, or rather was at once regurgitated. The tongue was clean, and the bowels constipated. She complained much of flatulence, and at night regurgitated water into the mouth. At first, vomiting several hours after food was the principal symptom. Soon after admission the food was at once returned; sometimes, however, it was retained for several days. She took creosote three times a-day, and opium at night, with considerable relief for a short time. On December 8th, I examined some of the water ejected from the mouth, but could not discover any cancer cells nor sarcina. She varied much, sometimes the stomach being excessively irritable, and rejecting everything, at other times she was able to take food. On Decem-

ber 19th, the tumour had not increased in size. She became more and more prostrate, and during the last month of her life suffered severely. She died March 26th. *On inspection*, the body was much emaciated. The brain was very much atrophied; and the convolutions separate. There was subarachnoid effusion, and an increase of fluid in the ventricles themselves. The septum in the ventricles was atrophied, and almost destroyed. *Chest*.—At the commencement of the œsophagus, the mucous membrane began to present an irregular granular appearance, with one or two whitish tubercles about the size of pins' heads; passing downwards, these tubercles became more numerous, till nearly opposite the root of the lung an ulcerated surface was found, with a raised margin and partially sloughing; still lower in the canal than this ulcer, the walls of the œsophagus were completely destroyed for about three inches, and the side of the right lung was in a sloughy condition; posteriorly, the pericardium bounded this sloughy mass, and there was an opening about the size of a sixpence extending through that membrane, opposite the left auricle, which was slightly affected with granular cancerous growth at that part. Nearer to the stomach, the walls of the œsophagus were again observed intact, but infiltrated with cancerous product, and nearly in a sloughy condition. At the floor of the cancerous ulcer were several branches of the pneumogastric exposed; the right one could be traced down to the ulcer, and several branches were completely truncated; another branch of the right nerve passed obliquely across the ulcer to the opposite side, to join the left nerve. On the left side, a branch was also observed to be truncated, and a large one ran for about two inches exposed in the sloughy tissue. The branches to the lungs were entire, and were situated above the cancerous growth. The ulcer in the œsophagus presented the elements of epithelial cancer. At the left lung were pleural adhesions; and the apex presented several white tubercles, at first supposed to be strumous, but found to consist of cancerous elements; the surrounding lung and the whole apex were of an iron-grey colour, from chronic pneumonia. The microscope showed large nucleated cells, resembling those of epithelial cancer, which were found in other parts, also smaller nucleated cells, a great number of granules, pigment, and some elongated fibre cells; other tubercles of a similar kind were observed in the lung, a few near the root, others at the periphery; the right lung was more free from disease. The larynx, trachea, and bronchi were healthy. The heart was small, destitute of fat, and atrophied. *Abdomen*.—In the stomach, near the œsophageal opening, was a raised tubercular growth about half an inch in diameter; it was ulcerated at its apex; its section showed that it principally involved the mucous membrane, but was extending into the muscular coat beneath. Some large nucleated cells were observed in the raised edges of the growth, and degenerated gastric follicles; some of the follicles were much enlarged, containing highly refracting particles, others contained nuclei. The rest of the mucous membrane and the pylorus were healthy. The head of the pancreas formed the hard mass which had been felt at the scrobiculus cordis; it consisted of hard, granular, whitish tissue, soft and breaking down in the centre, with some dense, firm, semi-transparent bands passing through it. On examination, it presented large epithelial cancer cells, elongated cells

forming fibres, and some cells undergoing degeneration. The adjoining lymphatic glands were infiltrated and adherent, the lesser curvature of the stomach was also adherent; the rest of the pancreas was normal. On the adjoining surface of the liver was an irregular tubercle, evidently produced by contact, and in the substance were several other small tubercles, but consisting of the same epithelial elements. The liver was small; its cell structure healthy. The gall-bladder was moderately distended, so also some of the bile ducts. On the right side, the cancerous infiltration extended to the right semi-lunar ganglion, which appeared to be infiltrated with cancerous product; cancerous cells being observed among the ganglionic cells. There were cancerous tubercles in both supra-renal capsules, but only involving a small portion of the organ. The duodenum, ileum, and colon were healthy, but the intestines were atrophied. The kidneys were granular, very small, and only four ounces in weight. The spleen was small, and there was a dense white patch on its surface. The cavity of the uterus was occupied by two soft polypi, and a large dense tumour, about three inches in diameter, was found in its walls; the tumour was dense and fibrous at its periphery, but did not present any trace of cancerous elements; its centre was tough, grey, and semi-calcareous; the ovaries were atrophied. (See Preparation 1799³.)

In this case the diagnosis was obscure, on account of the food being sometimes retained for several hours; which appeared to indicate disease of the stomach rather than of the Œsophagus. A tumour could also be felt at the scrobiculus cordis, the disease of the Œsophagus was, however, too extensive to produce obstruction, the walls of the lower part of the canal being entirely destroyed; the injury of the pneumogastric was very extensive, and the exposure of its branches was probably the cause of the severe pain from which the patient suffered.

CASE XVI. *Epithelial Cancerous Tumour in the Pharynx, closing the entrance into the Œsophagus. Effusion of False Membrane in the Larynx and Trachea. Acute Bronchitis.*—Charlotte W——, æt. 32, was admitted under Mr. Cock's care, February, 1856, and died March 6th. She was a married woman, short, anæmic, and somewhat emaciated. She had been out of health for a year, but for three months she had experienced very great difficulty in swallowing, and for several days it had become almost impossible to swallow anything except a small quantity of fluid; and the attempt now led to regurgitation through the nares. The effort of swallowing did not produce urgent dyspnœa. Respiration on admission was easy and normal, but there was slight hoarseness. On examining the chest, the respiration was found to be less free at the right apex. At the left side of the neck, below the angle of the jaw, was a prominent round tumour about one inch in diameter; it could be partially separated from the structures beneath. Mr. Cock attempted to pass a small bougie, but it was found to be quite impossible. The tumour in the throat could not be seen nor felt. A short time

before death very urgent dyspnœa came on, and she died from apnœa. At the lower part of the pharynx, attached to the cricoid and arytenoid cartilages, or rather the mucous membrane opposed to them, were four round tumours closely placed together, or rather one lobulated growth, extending as high as the upper margin of the epiglottis, and quite occluding the opening into the œsophagus. After removal, a probe could only be inserted by slowly passing it round the growth. The surface of the growth was grey, but was not ulcerated; its section was pale, and near the surface it presented red lines consisting of vessels full of blood. The soft palate also was considerably thickened. The inner surface of the epiglottis, of the larynx, and of the trachea, was covered by a layer of easily separable false membrane; the bronchi, especially the larger ones, were full of tenacious mucus. The tumour in the neck was soft, and of a pale yellow colour. The lungs did not collapse, but appeared quite healthy. The heart, liver, intestines, spleen, and kidneys, were healthy; so also the uterus and ovaries. The bronchial and abdominal glands were normal. On examining the growth from the pharynx, its base was found to consist of large cancer cells, containing a large granular nucleus, and the cells were closely arranged together. The growth in the neck had a similar structure. The surface was not ulcerated, but presented epithelium, which was normal in some parts. The appearance of the papillæ has been previously referred to; some were in a normal condition; in others, the central capillary was obstructed, and some were still more degenerated, closely resembling brood cells. (Prep. 1785⁷⁶.)

The obstruction at the commencement of the œsophagus in this case was mechanical, and the cause of death secondary laryngitis. The diagnosis of tumour was easy, but it was found to be quite impossible to pass any instrument beyond the growth. Many instances of non-cancerous polypi are recorded; and some may be removed by operation; in this case the growth was too low to be reached; and its character was less suited for operative interference.

CASE XVII. *Carcinoma of the Œsophagus, Communicating with the Trachea. Cancer of the Lung and of the Kidney.*—Catherine S—, æt. 38, admitted under Dr. Barlow's care, April 9th, 1856, and died April 17th. She had been a servant in a family for twenty years, and began to suffer from her present illness about six months before her death. On admission, she was in a state of great emaciation, and the dysphagia was extreme. The attempt to swallow food was at once followed by the regurgitation of it through the nose and mouth. The circulation was exceedingly feeble, and Dr. Barlow feared lest gangrene of the extremities might come on. She appeared to die from exhaustion. The body was much emaciated. In the neck, on the left side, was an enlarged cervical gland, about one inch in diameter, firmly adherent to the œsophagus and to the trachea; a smaller gland was situated on the right side; the former tumour could be felt before

the division of the skin. The lungs did not collapse freely. On dividing the trachea an opening into the œsophagus, somewhat oval in form, slightly pointed above and below, and about one inch and a half long, was found immediately above the division into the bronchi; the edges of this opening were thickened and slightly irregular. The corresponding part of the œsophagus in its whole circumference presented a nodular surface for three inches in length. The edges were raised and irregular, and the surface ulcerating; and there was slight vascular turgescence of the mucous membrane. Several cervical glands which were adherent to the œsophagus, were infiltrated with cancerous deposit; they were of a firm consistency, and were white in colour; but in the centre yellow. Other glands at the root of the lung were not at all infiltrated. The bronchi were intensely congested, and contained much dirty grumous fluid. The lower lobes of the lungs were much congested, and the right contained beneath the pleura a small mass, about half an inch long and a quarter of an inch broad, composed of yellowish-white cancerous substance. The left renal vein was filled with adherent clot, and its walls were considerably thickened. In this kidney were several cysts, and a minute tubercle composed of elements resembling the other cancerous structures. On examination of the œsophageal ulcer, a small quantity of juice from the section presented numerous nuclei; and in the section, some epithelial plates, cells with large nuclei, and caudate cells. It also presented some elongated nuclei and fibres, some of which had a curved arrangement inclosing nuclei and brood cells. The raised edges of the ulcer were composed of masses of these nuclei and cells, with some intervening elongated nuclei and fibres; and on the addition of acetic acid, some elastic coiled fibres were observed. The growth in the lung presented similar aggregation of nuclei. The cervical glands were of a much firmer texture, and much fibrous tissue was observed in them, forming irregular interspaces, in which nuclei were found. The central portions were yellow, and contained highly refracting granules (degenerating cancer). The great number of large nuclei resembled those found in medullary cancer, and this case appeared to be almost intermediate between medullary and epithelial disease.

CASE XVIII. *Cancer of the Œsophagus. Death from Exhaustion.*—Mrs. B—, æt. 54, a stout married woman, who had had ten children, and had ceased to menstruate for four years, experienced seven months before her death pain at the middle of the sternum, at the scapula, and in the loins; the pain was increased on taking food; she had slight palpitation of the heart, vertigo, and flatulence; her sleep was disturbed by pain; the passing of a probang down the œsophagus much increased the pain, which became very severe, and was aggravated by a chronic winter cough; deglutition also became very difficult; nutrient enemata were used, but she gradually sank, seven months after the commencement of the dysphagia. The central portion of the œsophagus was converted into a softened discoloured brain-like substance, from two and a half to three inches in extent. No gland nor other structure was affected; and the remaining part of the tube was healthy.

CASE XIX. *Cancer of the Œsophagus. Gangrene of the Lung.*—George D—, æt. 45, a very intemperate man of irregular habits, was admitted into

Guy's, suffering from symptoms of stricture of the œsophagus, with supposed phthisis. The breath was exceedingly offensive. At the apex of the left lung was a large gangrenous cavity. There was extensive ulceration of the œsophagus, which had ulcerated through to the vertebra, and communicated with the left bronchus. The sternal and mesenteric glands were enlarged.

CASE XX. *Cancer of the Œsophagus, with Pleurisy and Contracted Kidneys.*—John H—, æt. 66, seven months before his death first experienced difficulty in swallowing solids. The dysphagia increased, and swallowing at last became impossible. Till two days before his death he had no pain in his throat, but he had severe pain in the left side; this arose from pleurisy, the right pleura being found to contain more than a pint of purulent serum after death. There was constriction of the œsophagus, with some ulceration and considerable thickening around it. The kidneys were granular.

CASE XXI. *Cancer of the Œsophagus. Gangrene of the Lung. Communication with the Trachea.*—James M—, æt. 49, was admitted, under Mr. Hilton's care, into Guy's Hospital, on October 20, and died November 25, 1858. There was great dysphagia, and for a short time he had coughed up blood. *Inspection*—Two inches above the bifurcation of the trachea was a cancerous ulcer, two inches in length, in the œsophagus; its walls were defined and thickened; the surrounding parts were involved, and anteriorly was an opening into the trachea. The lymphatic glands were infiltrated. The left lung was universally adherent, the whole of the lung was in a state of early inflammation, and the root of the lung was gangrenous. In the right lung were patches of recent pneumonia. There was no disease in other parts.

This instance is simply mentioned as an exemplification of gangrene of the lung from direct extension of disease; so also the following case.

CASE XXII. *Cancer of the Œsophagus. Extension into the Lung. Gangrene.*—James S—, æt. 57, was under my care in Guy's Hospital; admitted May 12th, 1858, and died June 4th. For many years he had been a coachman, and temperate in his habits. In 1832 he had been ill for six weeks with pain across the chest, and for many years he had had cough. Eight weeks before admission, on attempting to eat, he felt that he was unable to swallow, and from that time he could take no solid food; the food seemed to pass as far as the scrobiculus cordis, and was then rejected. He was able to take fluids, but he had slight pain afterwards, and he was troubled with flatulence. He was a pale and emaciated man; the chest was normally resonant, the respiration murmur was very distinct, especially posteriorly and at the left apex. The heart sounds were normal; the abdomen was supple, and no tumour could be detected in the abdomen, but pulsation was very

distinct at the scrobiculus cordis. The superficial epigastric veins were slightly enlarged; the bowels were constipated; the pulse was quiet and compressible; he complained of weakness and vertigo. On the 21st he had gaseous eructations, which increased the dysphagia. On the 25th food was instantly rejected, and the vomiting produced a "cutting" pain in the epigastrium; at other times the pain was of a dull character. The vomiting and exhaustion became very trying to the patient. When unmixed with food, the rejected matters consisted of tenacious mucus; no sarcina were detected by microscopic examination. He died June 4th. On inspection the termination of the Œsophagus was diseased. For the space of an inch there was cancerous deposit infiltrated into the mucous and submucous tissues, and also into the muscular layer. The canal was much contracted, so that a small probe only could be passed. On the right side the growth was adherent to the lung; the cancerous tissue was at that part ulcerated, and communicated with the lung tissue, which was in a sloughing state. The upper lobe of the right lung contained pneumonic deposit, and several small cavities. The other viscera, and the lymphatic glands were in a healthy state. The examination of the growth showed much firm fibrous tissue, and some cells of epithelial cancer.

CASE XXIII. *Carcinoma of the Œsophagus. Pleuro-pneumonia from direct extension of disease.*—John D—, æt. 61, was admitted into Guy's Hospital August 19th, and died September 16th, 1857. He was an organ-builder, who resided at Cripplegate, and was a man of temperate habits. Two months before admission he was seized with cough, and pain in the left side, and he began to waste rapidly. On admission he was almost too weak to speak. The left side of the chest was found to be dull, and lobular respiration was observed. Stimulants were given, and he lingered for a day or two. The difficulty in swallowing was not particularly noticed, and was ascribed to debility. The body was extremely wasted. Opposite the bifurcation of the trachea the Œsophagus was entirely destroyed anteriorly, by a corroding epithelial cancer. It occupied a surface of about two inches, and although the whole circumference of the tube was affected internally, the disease had not penetrated the coats *behind*. The Œsophagus was seen to be quite entire at this part, but on its sides soft carcinomatous matter was seen to be protruding, and the trachea and bronchi were implicated in the disease. A large hole, through which the finger could be passed, formed a communication between the trachea and the Œsophagus, at the bifurcation of the former, and it extended into the left bronchus; soft sprouting cancer existed at the opening. The pneumogastric nerves were not implicated by the disease, the left was especially examined and found to be quite healthy and unaffected. The left pleura contained three or four pints of serum. The lower lobe of the left lung was hepatized, the upper presented small masses of deposit, as seen in phthisis; the right lung was healthy. No other organ was diseased.

CASE XXIV. *Cancer of the Œsophagus. Pneumonia. The Pneumogastric Nerve Involved.*—John D—, æt. 71, was admitted into Guy's

Hospital, on January 26th, 1859. For two years he had been under observation, and he had also previously been in the hospital. The difficulty in swallowing, and the emaciation, had increased to an extreme degree; but he was able to get some food down, until two or three days before his death, when the symptoms of pneumonia came on. On inspection, eleven hours after death, the body was found to be much wasted. The œsophagus was contracted one inch above the bifurcation of the trachea, the walls were thickened, and there was adventitious deposit effused in a circumscribed manner in the submucous cellular tissue. Some of the neighbouring glands were slightly infiltrated. There was pleuro-pneumonia of the left lung; the whole of the left lung was grey and solid, and a dark-coloured grey fluid exuded from it; but there was no sloughing. The left pneumogastric was involved in the diseased structure of the œsophagus. The heart and liver were healthy; so also the gall bladder and the ducts. The kidneys were degenerated, small, five ounces in weight, and they contained a few cysts. The diseased œsophageal substance consisted of fibrous and elastic tissue, squamous epithelium, large nuclei and cells in considerable quantity, and aggregated in clusters; and some of these clusters of cells were limited by membrane, as if forming part of a glandular structure.

CASE XXV. Cancer of the Œsophagus. Pneumogastric Nerves Involved. Pneumonia.—Richard E——, æt. 63, was admitted under Dr. Gull's care, September 14th, and died October 14th, 1859. There had been dysphagia for six months, with gradual emaciation; one month before admission he brought up blood; he was seized with pneumonia ten days before death; but was able to swallow fluids till the last. There was a large cancerous ulcer about the centre of the œsophagus; with a well defined and raised margin; it extended a short distance round the aorta, but did not implicate the vessel; it did not involve the bronchi, but penetrated the pericardium, which, at its posterior part, was softened, and of a greenish black colour. The right lung was universally adherent, and consolidated by recent and old pneumonia; several portions of very fetid lung existed in the upper lobe; and in the lower lobe were several hepatized lobules, with red intervening tissue. The left lung presented numerous patches of hepatized tissue. The pneumogastric nerves were involved. The stomach and other viscera were sound.

A brief notice of the last eight cases is given to illustrate further the causes of death. Case XVIII from exhaustion, complicated with chronic bronchitis. Cases XIX, XXI, and XXII from gangrene of the lung. Case XX was complicated with renal disease and acute pleurisy. In cases XXIII, XXIV and XXV, pneumonia was produced either by extension of disease to the lungs, or by injury of the pneumogastric nerves.

CASE XXVI. Carcinoma of the Throat. Strumous Pneumonia.—Martha M——, æt. 31, admitted December 5th, 1855, and died on the 20th, at two

p.m. She was a short woman, married, and had been confined fourteen months previously, but since that time had not been well, having suffered from a slight cough. For three weeks she had had difficulty in swallowing, and this had increased to such an extent that she was, on admission, unable to swallow food, except with extreme difficulty. She could, with much distressing pain, swallow solids, but fluids were at once regurgitated through the nose. She suffered from hunger, but still more from thirst, and she was extremely emaciated. The glands at the angle of the jaw on the right side were much enlarged, giving her emaciated countenance a miserable appearance. Her voice was nasal, and she was extremely exhausted. She was too ill to allow the chest to be examined, and died on the 20th. Her relatives, brother, &c., died of phthisis. On removing the larynx and tongue the soft palate was found to be about twice its natural thickness, irregularly tubercular, and brawny; the posterior pillars of the fauces were affected in a similar manner (Preparation 1785⁷⁰). On the right side there was a communication from the pharynx into an irregular cavity, situated opposite or rather behind the angle of the jaw, about two inches and a half in length, and half an inch in breadth, and containing almost black sloughy substance. The glands were infiltrated with firm, cancerous product. The tissue of which the soft palate was composed consisted of an immense number of nuclei. In the lungs, there were firm adhesions at the apex of the right lung, the pleura being semi-cartilaginous. In the remaining part of the lung numerous minute tubercles were observed beneath the pleura, and at the lower lobe there were also moderately firm adhesions. The left pleura was free. At the apex of the right lung was an irregular vomica, capable of holding about two drachms of fluid, with a smooth lining, and surrounded by iron-grey lung, with several opaque tubercles. At the lower lobe a considerable portion of the lung was red and consolidated; and several lobules were infiltrated with pale yellow, low organized deposit, which was breaking down in several parts, and precisely resembled the lung observed in cases of acute pulmonary phthisis. There was considerable congestion of the bronchi, and they contained tenacious mucus. The left lung was congested, but was otherwise healthy. The bronchial glands were black from pigment, and those quite at the base of the neck were firm, white, and dense, consisting of nuclei resembling those in the palate. In the lung, the tubercles presented no nuclei like those in the palate, nor did they consist of cancerous growths, but were composed of imperfectly developed nuclei, dark pigmental granules, and some nucleated cells. The tissue of the thickened pleura consisted of fibrous tissue. The heart was exceedingly small, and destitute of fat; its cavities contained moderately firm clot, and the valves were healthy. The liver, and also the spleen, were healthy. The stomach and the intestines were contracted and healthy, except the rectum and sigmoid flexure, the mucous membranes of which were congested in longitudinal stripes; and numerous minute superficial ulcers were scattered along these patches.

This case is one of great interest in the connexion of cancerous disease of the pharynx with strumous pneumonia, diseases rarely

conjoined, occurring at different periods of life, and considered antagonistic the one to the other. In advanced life, when death has occurred from cancer, we sometimes find the remains of strumous disease of an earlier period of life, in the form of calcareous deposit at the apex of the lung, or in wasted glands; such cases are, however, rare and exceptional.

CASE XXVII. Cancer of the Œsophagus. The left Pneumogastric involved. Pneumonia.—William E—, æt. 50, was admitted July 30th, 1856, under Mr. Callaway's care, and died September 18th. He had suffered from dysphagia for six months, and could not swallow solids. He had cough, and expectorated tenacious mucus. His cough and expectoration became worse, and the lung tissue involved. During the last week of his life he swallowed with more ease. Inspection was made eighteen hours after death; the body was much emaciated. The head was not examined; the abdominal viscera were quite healthy. On removing the sternum the disease was at once exposed; and on trying to remove the lungs, the ulceration of the Œsophagus broke through into the left pleura. The ulceration in the Œsophagus extended from the cricoid cartilage to the bifurcation of the trachea; the edge was well defined, raised, and yellowish; the central part was ulcerated, and the whole circumference of the Œsophagus involved; in front, the cartilages of the trachea were exposed, and immediately above the bifurcation was an opening about the size of a sixpenny piece, with irregular serrate margins. The ulceration extended downwards and outwards, and was closely connected with the external surface of the left bronchus; it had involved the pneumogastric nerve on that side, one of the larger branches of which was truncated. Posteriorly, the vertebræ formed the boundary of the ulceration. The greater part of the lower lobe of the left lung was in a state of grey hepatization, and towards the apex there was some iron-grey hepatization, with whitish tubercles. These tubercles appeared to be of a cancerous character. In the right lung was another small mass of condensed lung. There was slight infiltration of the adjoining bronchial glands. On microscopical examination the ulcer and bronchial gland were found much decomposed; but there was no doubt of its being of the character usually called epithelial cancer.

The more easy deglutition towards the last week of life is possibly explained by the extension of the ulcer having destroyed the whole of the circumference of the Œsophagus, and thus prevented any spasmodic obstruction.

The pneumonia of the left lung, was no doubt accelerated by the injury to the nerve on that side; but it must be borne in mind, that the cancerous growth extended to the left bronchus.

CASE XXVIII. Cancer of the Œsophagus. Communication with the left bronchus. The Pneumogastric involved. Old vomica in the Lung. Extension

of disease through the Diaphragm.—George W—, æt. 53, was admitted, under my care, September 3rd, 1856; he was an old man, emaciated and grey; he had been a blacksmith at Chatham, and, until six weeks before, he had enjoyed good health; at that time he experienced pain in swallowing food, especially solids, which were almost at once rejected. He had pain across the sternum. The day of admission he came from the North Foreland; and was exceedingly prostrate, emaciated, and cachectic; the pulse was feeble; the heart was irregular, with a slight bruit; no tumour, nor enlarged glands could be felt; but the abdominal aorta pulsated very distinctly. The abdomen was collapsed; in the chest were general bronchial râles; the breath was not offensive as in sloughing lung; and he had no distressing cough, but hiccough. On the 6th, he was more prostrate; the hiccough was distressing; the motions were black; and he was bringing up brownish-coloured blood; he gradually sank, and died on the 8th, at 2 a.m. Inspection was made twelve hours after death. The body was much emaciated. *Œsophagus.*—Two or three inches from the commencement of this canal were several small ulcerated surfaces of a pale, yellowish colour, with central depression; an inch further the whole of the walls of the Œsophagus were destroyed, and the margin defined; beyond this part was an irregular flocculent grey tissue, floating out when placed in water; it was found upon a dense fibro-cartilaginous base, firmly adherent to the trachea, aorta, and other tissues; an inch from the left bronchus was a circular opening, about three-quarters of an inch in diameter, forming a communication between the Œsophagus and the bronchus; the latter tube contained a flocculent grey mass, which almost obstructed the bronchus. Some of the bronchial glands were partially infiltrated. The pneumogastric nerve extended into the dense tissue at the base of the ulceration, and some of its branches were exposed at the floor of the ulcer. The destruction of the Œsophagus extended to the diaphragm, and the ulceration passed through it, so as to form an irregular sloughing cavity below that muscle, bounded by the stomach, by the cellular tissue, by the large vessels, and partly by the left lobe of the liver. The cardiac opening into the stomach remained in its normal condition; but a second opening had been formed near it, from the abscess just mentioned. The ulceration also extended into the liver. The branches of the sympathetic were partially destroyed, but could not be satisfactorily dissected; some of them were very hard; on microscopical examination, nuclear nerve fibre, apparently undegenerated, could be detected. The coronary artery of the stomach was obstructed by clot; some of the glands at the lesser curvature of the stomach were infiltrated. The ulceration almost extended into the thoracic aorta; that vessel was exceedingly diseased, from atheromatous and ossific deposit, and in two parts had a greenish appearance; there appeared to be minute communication beneath a bony plate and the ulcer in the Œsophagus, but no probe could be passed. On examining the upper margin of the Œsophageal ulcer, large cancer cells were detected and some nuclei; the surface of the flocculent growth consisted of pointed processes filled with granules, sometimes several proceeding from one trunk. The stomach was exceedingly contracted, and of hour-glass form; the mucous membrane was healthy. The left lobe of the liver,

which was somewhat enlarged, almost obscured the stomach; the liver itself appeared healthy, its weight was 3 lbs. The pancreas and spleen were healthy; the gall-bladder contained some bilious mucus. The bronchi were congested and full of frothy mucus. At the left apex was an old vomica, surrounded by iron-grey lung and calcareous deposit; its lining was smooth, and it was capable of containing about 3ss of fluid: its contents were dirty mucus. The pleura, on the left side, was universally adherent; on the right side, it was partially so at the apex. The right lung also contained a small vomica, but there were no tubercles in it; a small, white, dense tubercle, of questionable character, was situated beneath the right pleura. The remaining part of the lungs were œdematous. The pericardium contained an excess of fluid; the heart and its valves were healthy; the aorta was exceedingly diseased throughout, being rough, scabrous, and bony; weight of the heart was 9½ ounces. The kidneys were atrophied, and contained several cysts.

The pain at the sternum, the difficulty in swallowing solids, the emaciation, the cachexia, the age, all indicated organic disease of the œsophagus. The general bronchial râles pointed to some communication having been set up; and such was believed to be probable. There was no pain at the scrobiculus cordis, nor was there any apparent indication of the abscess which existed. The prostrate condition of the patient, probably, prevented the development of more manifest peritonitis.

The disease had, probably, existed for a longer period than six weeks, if we judge by the destruction of nearly the whole œsophagus, and the firm character of the tissue which bounded it. It was evidently cancerous; although no other part except those in immediate contact were affected. But the villous flocculent character of the growth in the œsophagus, with evident cancer cells at the margin of the ulceration, appeared to indicate that it somewhat differed from ordinary so-called epithelial cancer. The small vomica at the left apex was not diagnosed; it appeared to have been in a passive condition, but its association with cancerous disease is an exceedingly interesting circumstance. It is doubtful whether any blood oozed from the aorta, or whether that effused, was from the coronary of the stomach.

Stimulants, beef tea, milk, julep of ammonia, and ether, with lead and opium, were prescribed; it is, however, probable that the stimulants passing into the false cavities which had been formed, tended rather to irritate than to produce effectual

benefit. Nothing more, however, could have been done, except perhaps, by the use of nutrient injections; but the patient could swallow fluids and retain them, so that these means appeared scarcely called for.

CASE XXIX. *Cancer of Œsophagus. Pneumogastric nerves truncated. Sloughing extending through the lung, and through the diaphragm.*—John H., æt. 45, was admitted into Guy's Hospital February 17th, and died March 2nd, 1858. He was a tall, emaciated man, who had been ill for several years. He had no dysphagia, but the food was generally rejected at once; sometimes, however, it was retained. He had no pain between the shoulders, nor on pressure at the region of the stomach. He gradually sank. *Inspection*—In the right lung the lobules were consolidated very generally, and were infiltrated with offensive serum. At the root of the lung, below the vessels, was a circumscribed slough communicating with the diseased œsophagus. The left lung was affected in a similar manner, but in a less degree. *Œsophagus.*—At the root of the lung the tube was irregularly truncated, and a large sloughing cavity formed, which was bounded by the lungs, one of which was encroached upon anteriorly by the posterior surface of the pericardium. At the lower part the diaphragm had sloughed; and the sloughing cavity was limited below by the pancreas, by the anterior surface of the stomach, and a small portion of the liver. The Pneumogastrics were both truncated. In the stomach, at the cardiac orifice, there was an irregular infiltration of the mucous membrane by cancerous product, and two openings extended into the sloughing cavity before mentioned; these openings were bevelled on their inner aspects. The rest of the stomach was healthy. There was no glandular enlargement, nor disease of the spleen, liver, intestines, &c. The pericardium was adherent, the heart small, the valves healthy.

The physician who had the care of this case regarded it as one of pyloric disease on account of the remarkable absence of pain and difficulty in swallowing after the patient came under observation in the hospital. This immunity probably arose from the manner in which the œsophagus and its nerves were truncated.

CASE XXX. *Cancer of the Œsophagus. Chronic Pneumonia. Vomica. Acute Pneumonia.*—William G., æt. 69, applied to me Dec. 4th, 1856, suffering from dysphagia. In early life he had been an attorney, but had evidently been reduced in circumstances; for twelve months he had had cough and shortness of breath, sometimes palpitation of the heart, but no hæmoptysis; for twelve months also he had pain across the chest, but no expectoration; his health continued tolerable till two months before I saw him, when he first experienced difficulty in swallowing; this gradually increased in severity, so that in December he was only able to swallow

liquids, and that with considerable pain. The pain was situated about the level of the third rib, at the sternum; the sensation being as if a foreign body was retained at that part; the ability to swallow was occasionally relieved, but never completely so, nor did he ever become free from pain. On examining the chest it was found flattened at the right apex, and there was tubular breathing at that part, with mucous rattle on the whole of the right side; the voice was slightly increased in resonance; at the left apex the respiration was coarse; and sibilant râles existed at the base of that lung. The heart was feeble, but nothing abnormal was detected in it, nor in the abdomen; there was slight pain in the back, between the shoulders; he had an aged appearance, and was moderately emaciated; the tongue was clean; the bowels were regularly open; the urine was healthy. The arteries at the wrist and in the neck were rigid; and no enlarged glands could be felt in the neck. He was recommended not to attempt solid food, but to take milk, eggs, &c., and a cough mixture was prescribed. December 17th, he was worse, and for three days had been scarcely able to swallow any food; it appeared to lodge in the throat; nothing, however, could be seen, and there was no evidence of aneurism. He had more pain between the shoulders; there was occasionally, for half an hour, slight relief to the dysphagia, but he was more distressed; with the bronchial breathing at the left apex, there was gurgling, and his cough was troublesome; he said that he expectorated the milk which he had attempted to swallow; he was exceedingly feeble, and was not able to take any sleep. There was evidence of old disease at the apex of the right lung; and acute bronchitis with it; with these were associated tolerably clear evidence of organic disease of the œsophagus, probably cancerous; such was my opinion at that time, and it appeared probable that the disease had extended into the bronchi, as is frequently the case. In this condition, he was admitted into Guy's, under my care; he was requested not to attempt to swallow, for this effort produced spasmodic contraction of the œsophagus, and several nutrient enemata were given. The following day he swallowed with greater facility, and could take beef tea, eggs, and milk, with a little brandy; his cough, however, was more troublesome; the sputum was purulent, nummulated, and, on microscopical examination, presented no evidence of cancer cells, but some curved elastic fibre, resembling lung structure, and large inflammatory granule cells. After admission no food or milk was vomited. He continued in the same state, sometimes for a few days being much more comfortable. A small quantity of cod-liver oil was given and morphia at night. During the month of January, he continued to emaciate, and became more anæmic; he took his food with more relish; but the attempt, at my request, to swallow a portion of softened bread produced great discomfort. Morphia, hydrochloric acid, and calumba afforded slight relief. On February 23rd, his mind was wandering; the stools were discharged involuntarily; the following morning he died. *On inspection*, the lung was found to be very firmly adherent at the right apex, and a thick dense layer of fibrous tissue was with great difficulty separated; the whole of the right pleura was destroyed; on making a section of the right lung, a small vomica was found at the apex surrounded with iron-grey lung, the surface was smooth; the lower lobe was

in a state of hepatization, and some lobules were red and consolidated; there were also some smaller granular masses. The left lung was in a similar condition, except the disease at the apex; the pleura over the lower lobe was covered with a thin layer of lymph. Some of the bronchial glands were slightly infiltrated with cancer, but there was no evidence of cancer in the lungs.

In the *œsophagus* was an ovoid mass, about six inches in length, and one in thickness, attached at the root of the lung, and reaching nearly to the cricoid cartilage; the canal was dilated; the mass was of a pale yellowish colour, and was softened in the centre; it was attached only to one side of the tube, and no smaller tubercles were observed on the mucous membrane; no communication with the trachea, nor bronchi, existed; the tumour consisted of nuclei and nucleated cells resembling medullary cancer; none of the blood cells usually found in epithelial cancer were observed. The pneumo-gastric nerves were free, and the disease appeared to have commenced in the mucous membrane. The heart was healthy; so also all the abdominal viscera; the intestinal canal was much contracted, but contained solid *fæces*. The liver was slightly congested, and the gall bladder was much distended.

The existence of a disease so closely resembling pneumonic phthisis as that found in this case was very interesting when we consider it in connexion with the cancerous disease of the *œsophagus*, and with the age of the patient. It was my opinion, during life, that the disease in the *œsophagus* had extended into the bronchi, but this was not found on inspection. The only other disease which appeared to be probable as a cause of the dysphagia was aneurism; but, the persistence of the dysphagia in every position, and the absence of other signs of aneurism, led me to believe that the obstruction was of a cancerous character. If the patient had been much younger it might easily have been supposed that the case was one of ordinary phthisis, with severe ulceration about the larynx and epiglottis: we had evidence of chronic disease of the lung, with acute disease; and in phthisis the dysphagia is sometimes exceedingly severe and distressing; but the patient did not lose his voice, the food was never regurgitated through the nose, nor did it produce spasmodic cough; the obstruction was evidently below the epiglottis.

No attempt was made to explore the *œsophagus* with any bougie or tube; the danger and discomfort which would have arisen from it did not warrant such an attempt being made. The use of nutrient enemata, even for a single day, removed the

very urgent dysphagia which existed on his admission. He had previously tried to swallow, till he found himself exhausted.

CASE XXXI.—*Cancer of the Œsophagus. Artificial opening made into the Stomach.*—Walter H—, æt. 47, was admitted into Guy's Hospital October 8th, 1857, under my care. He had resided at Tunbridge Wells as a stableman, was of ordinary stature, light complexion, and moderately nourished. He stated that for sixteen years he had had winter cough, but that he had never had dropsy. On admission there was considerable dyspnœa; the lips purplish; the pulse compressible, but regular; the chest was resonant on percussion, and the respiratory murmur indistinct; distant prolonged expiratory murmur was everywhere audible, with some sibilant râle; the voice was also indistinct, and tactile vibration diminished. The heart-sounds were regular and normal; the expectoration frothy and moderately abundant. The abdomen was moderately full and rounded, and there was a small hard gland felt about the anterior margin and upper part of the sterno-mastoid muscle. After he had been in the hospital for a short time, he began to complain of severe pain in the throat during coughing; but, on carefully examining the part, nothing could be perceived. In a few weeks pain was also produced in swallowing, especially when solids were taken, and cough continued unrelieved. On December 14th, he continued to suffer severely, and became more anæmic; the countenance was expressive of great distress, and the mind irritable; deglutition had become very difficult, so that he could only take fluid forms of food, and some stimulant. The cough also was very troublesome, producing very severe pain in the throat; it was violent, and small drops of blood were spirted out in the act of coughing; the expectoration was thin and watery. The chest continued resonant; respiration very feeble, on the left side indistinct, and the expiratory murmur was prolonged; the larynx was free in its movements. Nothing could be seen in the throat, except slight œdema and redness towards the right side. The gland at the angle of the jaw remained of the same size; the pulse was compressible: the tongue clean; the bowels confined. Various means were tried, as conium and carbonate of soda, with hydrocyanic acid, steel, &c. The bowels were acted upon by colocynth and henbane, by magnesia mixture, or by injections. Counter-irritation was applied to the throat—hot-water fomentation, or cataplasms, blisters, &c. The inhalation of steam afforded some relief, but still more the smoking of stramonium leaves. Tincture of aconite, applied externally, was also of some benefit. Mr. Cooper Forster examined the throat for me, but could not detect any cause of obstruction. The patient continued during January and February without any improvement, the emaciation increased, and both respiration and deglutition became more difficult, especially the latter. Morphia occasionally given, and the stramonium inhaled, afforded partial relief. On again examining the throat, Mr. Forster felt below the epiglottis, towards the right side, a rounded tumour, which was evidently obstructing the commencement of the œsophagus, and he believed that its surface was ulcerated.

The respiration, although noisy and accompanied with a loud inspiratory sound, was not hurried, and sufficient air appeared to enter the larynx. The propriety of performing tracheotomy was discussed, but it was decided that no benefit would be likely to accrue from it. The examination of the growth in the throat was followed by temporary relief; and the patient was able, for three or four days, to swallow solid food. The stramonium and other means of relief, were continued, and nourishment was given in any form that could be taken. On March 1st the emaciation had very much increased. During inspiration a loud noise was produced in the throat; and this sound had for some weeks been increasing in intensity, so that he had been unable to sleep for several nights, on account of the "roaring," as he termed it. His voice had become more feeble, but the cough had almost ceased; he could only swallow fluids, and those very slowly; his nourishment latterly had consisted of milk and rum, with eggs. Deglutition was much relieved for two or three days by two small blisters applied on either side of the larynx; but it again became so difficult, that nutrient injections were resorted to. These injections, by allowing the throat to rest, had enabled him to swallow with more comfort. The pulse had become very compressible and small; the bowels occasionally constipated. March 2nd. The respiration became more difficult, and Mr. Stocker had tracheotomy performed in the night. The incision was made as low as possible, but the trachea appeared flattened from behind, and the patient could not bear the tracheal tube inserted; when it was attempted, he appeared to be quite incapable of breathing. The operation did not afford relief, and a deep-toned rhonchus could be heard in the lungs. There was no congestion of the face; the pulse was very compressible; the cough slight; he was able to get down his rum and milk, and some blanc-mange, &c. On the 9th he was breathing more comfortably; the opening in the throat was patent, and thin pus covered the red margins of the wound; there was also less noise on inspiration. 24th. The emaciation and prostration of strength increased; his bones appeared barely covered with thin skin, and the face expressive of starvation. He said "he was famished." He endeavoured to relieve his distressing thirst by moistening the mouth; but for four days he had not been able to swallow a drop of fluid. The attempt to swallow at my request was preceded by much hesitation and preparation, and was followed by a paroxysm of severe coughing. The expectoration had changed in character, and had become muco-purulent. On examining the chest sibilant râles were everywhere faintly audible. There was no dulness on percussion, but preternatural resonance. The voice was very feeble, and scarcely audible; the pulse slow and very compressible; the tongue clean; the larynx was moveable; the gland at the angle of the jaw as before; the opening in the skin made during tracheotomy remained open, and the skin was undermined, there being evidently no power to repair the wound. The abdomen was exceedingly contracted, the pulsation of the aorta being visible, and the arteries most distinctly traceable. There was no evidence of enlargement of the liver, nor of disease of the abdominal viscera. He complained of pain towards the right side, and tied a handkerchief firmly around him to relieve the sense of hunger. The skin was dry. He passed about a pint and a half of urine during the day.

The sleep was tolerable; the mind clear and active. Nutrient injections of beef tea, eggs, and rum, thickened, if possible, with flour, had been given, at first four times and then six times a day. Milk also was ordered, and \mathfrak{M} v of tincture of opium to be added to each injection. On the 25th he appeared to be sinking, and the rectum ejected the enemata almost at once. His hands were cold; but he complained of a sense of heat.

It now became a question, whether life was to be allowed gradually to die out, or an attempt to be made by any other means for the introduction of food; the patient appeared to have chronic bronchitis, with epithelial cancer at the commencement of the Œsophagus, possibly extending into the trachea, and death threatened from inanition. Three modes of relief suggested themselves—1st, the forcible introduction of an Œsophageal tube; 2nd, opening the Œsophagus in the neck; and 3rd, opening the stomach. In reference to the first, there was evidence of a growth at the commencement of the Œsophagus; and the trachea appeared partially compressed, as shown in the operation of tracheotomy. The disease in the throat was probably of the form of epithelial cancer, and the passage of a bougie must have been constantly repeated. The great irritation and coughing produced by attempting to swallow, showed that the epiglottis was extensively ulcerated; or, that there was a communication between the Œsophagus and trachea, which would render the passage of a bougie very dangerous. In some cases of cancer of the Œsophagus, a bougie has been passed into the pleura, and led to a speedy death; and probably the passage of a bougie could not have been effected; this decided against the first proceeding. As to the second, namely opening the Œsophagus, since the most frequent seat of cancer in that tube is opposite to the root of the lung, about the third dorsal vertebra, and consequently beneath the position at which the canal could be opened, the operation would have been very formidable, dangerous, and even useless. The third proposition, that of opening the stomach, appeared to be the only operation which could possibly relieve the patient.

Wounds of the stomach, as that of Alexis St. Martin, the cases recorded by Mr. South, those by Dr. Murchison, &c. showed that life could be continued after fistulous communication had been thus made. The operations on the lower animals

proved that it could be performed with some probability of success; such an operation would give a chance of prolonged life, where death was certain; and where the peritoneum was healthy, there was less danger than in abnormal conditions of that membrane. If life were prolonged only for a short time, and food introduced, there would be relief to the distressing thirst and the fearful sense of starvation: and, lastly, it was evident that the patient was dying from inanition rather than from the disease, nutrient enemata being refused. On the other hand, however, I felt that the disease was probably of a cancerous character, and would sooner or later terminate life; that the operation was a hazardous and uncertain one; and that life might possibly be continued for a few days by a small portion of the injection being retained. After carefully weighing these facts, I asked the assistance of my colleague, Mr. Cooper Forster, and if he considered the operation of opening the stomach through the anterior abdominal parietes, for the purpose of introducing food, a feasible and warrantable one, I decided that it should be attempted. The operation was accordingly performed, and the stomach was opened by an incision commencing at the extremity of the eighth rib, and the edges were stitched to the wound. March 26th. The operation took place about half-past 2 p.m., and was borne without a movement on the part of the patient. The pulse, which before the operation was 62, and exceedingly compressible, rose to 116. Six drachms of milk with part of an egg were introduced through an elastic tube into the stomach. At twenty minutes past 3, about two ounces more milk and egg were introduced; the patient then complained of feeling a sense of heat, but appeared comfortable. He was now removed to bed. At 4 p.m. the pulse was 120 and still very feeble; it was decided to introduce every half hour, if the patient were awake, two ounces of milk and egg, and every second time two drachms of rum with it. At 9 p.m. he was comfortable; there had been slight pain in the left side; the pulse was fuller, 124; the skin less parched; and he had slept occasionally for a short time. Messrs. Greenwood, Gayleard, Owen, and Tuck, kindly volunteered to remain with him in rotation. During the night he had four hours sleep; he passed urine, and there were three

slight watery evacuations from the bowels. 27th. About 10 a.m. he coughed violently, and the contents of the stomach were forcibly ejected through the wound. His pulse continued 120. At 1 p.m. he was cheerful, his eyes more bright, his voice stronger, the skin less parched, his tongue moist, thirst and the sense of starvation relieved; he had pain in the left side; the pulse 120, and very compressible; his hands were cold, feet and legs warm; the coldness of the hands had been very marked for several days. The operation had evidently mitigated his suffering. At 1:30 p.m. half an ounce of rum, with sugar, and an ounce and a half of water, and fifteen minims of lemon-juice were given. The stomach received it well, contracting upon the tube. He said that it produced a comfortable sense of warmth throughout the abdomen. At 3:30 the pulse was firmer and fuller than at 1 o'clock, and the hands warmer. Since the operation, during the twenty-four hours, he had six eggs, beaten up in twelve ounces of milk, given in small divided doses, with four ounces of rum. Milk and egg, or beef tea thickened with flour, were ordered every half hour, and occasionally half an ounce of rum, as just mentioned. At 8:30 p.m. faintness came on, the face became cold and perspiring; pulse 136, and scarcely to be felt. The stomach appeared to have lost its power of contracting on the food introduced.

Stimulants were ordered to be given repeatedly and freely, with nourishment as before; and two or three times, as a stimulant, $\mathfrak{m}\mathfrak{x}\mathfrak{x}$ of tincture of sesquichloride of iron. During the night he was evidently sinking, the pulse sometimes became scarcely perceptible, but rallied after stimulants were introduced. On the 28th he slept for a short time about 10 a.m., and expressed himself as comfortable; but gradually became unconscious, and died at 10:45, rather more than forty-four hours after the operation.

Inspection was made twenty-eight hours after death. The body was extremely emaciated. The head was not examined. At the lower part of the neck, immediately above the sternum, was the wound made in tracheotomy, gaping and undermined, and on the trachea a few drops of pus. At the left hypochondrium was the opening made by the operation of gastro-

tomy, also enlarged by the plug which had been introduced a few hours before death. The mouth and soft palate were healthy, also the epiglottis. At the posterior surface of the cricoid cartilage there was a growth connected with the mucous membrane, about a quarter of an inch in elevation, and extending from side to side, soft and slightly injected; passing downwards, there was irregular ulceration, and towards the trachea destruction of all the coats of the œsophagus; on either side and below, the ulcer was bounded by a sharp undermined edge. The cellular tissue of the trachea and its muscular fibres were destroyed for about half an inch; the mucous membrane was bare, and perforated by a small opening about one sixteenth of an inch in diameter, so that fluid could pass from the œsophagus into the trachea; below the ulcerated surface in the œsophagus the canal was much contracted by infiltration into the surface of the mucous membrane; the passage was so much diminished at this part that a probe could only be passed after death, and it was probably quite impervious to fluids during life. The constriction *was situated at the level of the first bone of the sternum.* The rest of the œsophagus was healthy. One or two glands in the neck were infiltrated and diseased, but none of the mediastinal nor other glands. The rima glottidis was free; the vocal cords and aryteno-epiglottidean folds were quite healthy; so also was the trachea. The bronchi contained thick tenacious mucus. The pleura on the left side was healthy; on the right, there were general, but not firm adhesions. The lungs were both much distended with air; they were pale, emphysematous, and covered the heart. At the right apex the lung-tissue was puckered; there were numerous lobules of iron-gray consolidation, with intervening crepitant lung, but no disorganization. The lower lobe of the right lung afforded a beautiful specimen of emphysema, but there were numerous gray tubercles studded in small clusters; they were non-cancerous. The lower lobe of the left lung was much congested; and one or two lobules were softened and breaking down, from acute changes, which took place, probably, a very short time before death. There was no enlargement of the bronchial glands. In front of the surface of the heart was a small collection of pus, only a few drops, apparently from the inflammation

of a small gland. The pericardium and heart were healthy; the heart was contracted and firm. On opening the abdomen, the intestines were found contracted; *the peritoneum was healthy; no inflammation, as shown by effusion of lymph, serum, or diminution of the normal smoothness, could be detected.* The stomach was partially distended; it was situated lower than usual, and its anterior surface was looped up to the opening in the anterior abdominal parietes made by Mr. Forster at the linea semilunaris. The mucous membrane of the stomach was pale, slightly injected at the opening. On gently drawing aside the stomach at the opening, the opposed serous surfaces were found slightly adhering. The small intestine was healthy throughout, but atrophied; the food introduced had only passed four feet down the intestine; below that point the intestine was exceedingly small. The lower part of the ileum was healthy. In the colon there were several patches of congested mucous membrane. The gall-bladder was distended, the liver healthy, so also the kidneys; the spleen was very small. There was no evidence of any cancerous disease affecting any part except the Œsophagus and one or two adjoining glands.

It is probable that chronic disease of the right apex had existed for a long time, so also the emphysema of the lungs. Other miliary tubercles were perhaps of recent deposition. The lobular consolidation and congestion of some parts of the left lower lobe of the lung were evidently only of very brief duration. As to the cause of the cancerous disease we have no evidence; it was probably of about six months' duration; its existence with chronic disease of the lung is a fact of some pathological interest. The microscopical examination of the growth in the neck presented the form of cell-growth common in epithelial cancer of that part.

In reference to the diagnosis in this case, it was evident at the time of admission that the patient had chronic disease; he had had cough for fifteen years; there was, however, no evidence of serious obstruction to the heart or to the portal circulation; the dysphagia was then a new symptom. The examination of the chest did not give any indication of phthisis, the respiration was exceedingly feeble at the apices, but there was no dulness

on percussion, and the voice was diminished rather than increased in resonance ; at the bases the expiratory murmur was prolonged ; it appeared probable that there was no marked consolidation of the lung, but rather that the feebleness of the respiratory act arose from emphysema. The question arose, whether there was any pressure on the right bronchus ; but of this there was no proof ; the respiration at the right base was as strong as at the left, and there was no increase of resonance at the right apex, though the respiration was less distinct. As to the cause of the dysphagia, it was naturally suggested, whether it was a case of phthisis, in which the principal disease manifested itself in the throat, by ulceration of the epiglottis. I have several times observed in severe ulceration of the epiglottis very severe dysphagia ; and the condition may be mistaken for obstructed Œsophagus. Few, however, would have had that idea in this case, for there was wanting at the early stage the raucedo of phthisis ; and the difficulty in swallowing was evidently at a later stage of the process than that produced by ulceration of the epiglottis : there the attempt to swallow is scarcely commenced before the food is ejected, often through the nares ; here, it passed beyond, there was no such ejection of food, but rather severe pain, and that extending to the ears. The blood also which was spirted out during coughing was not as we observe it in ulcerated larynx, but was evidently from the pharynx. Subsequently a tumour could be felt at the commencement of the Œsophagus. The emaciation was not that of simple phthisis : there we generally have a rounded abdomen, and often more or less diarrhœa ; here there was uniform constipation and collapsed abdomen. There was no evidence of syphilis to account for the affection of the throat, nor of aneurism. It was presumed that the affection of the pharynx and Œsophagus was of the form of epithelial cancer ; the enlarged gland in the neck confirmed this idea, and there was nothing to indicate any such disease of other viscera. As to the prognosis, that was from the first unfavorable, but it was not for several weeks anticipated that such serious disease of the throat would present itself ; all idea was then given up of ultimate recovery. In the *treatment*, the object was to relieve the irritation of the bronchi and the engorgement of the portal system ; and to strengthen the patient,

meat diet was given. The excreting organs and glands were acted on by squill and blue pill, by purgatives, &c. The spasmodic contraction of the bronchi were relieved by conium, alkalies, hydrocyanic acid, and stramonium. The pain produced by swallowing and coughing was most distressing, and sometimes kept the patient awake for several nights. Inhalation of steam afforded relief, so also the smoking of the leaves of stramony. The application of nitrate of silver aggravated the distress; but small blisters on either side of the throat were productive of the greatest benefit; tincture of aconite also, applied externally was more palliative than chloroform; morphia administered internally produced transient composure; however, all these means, could not check the progress of the disease. When it became necessary to resort to nutrient injections, life was fast ebbing away. It may be a question as to what is most effective in this case; beef tea thickened with flour, an egg, and a small quantity of rum were used. Dr. Gull suggested the propriety of using pepsine mixed with the fluid, and since the rectum is incapable of rendering the aliment in a condition ready for absorption, it may be well deserving of trial. The longest period in which I have known a patient nourished by injections alone, was in a case mentioned by Dr. Barlow, in which for seventy days food was administered only in this manner. In this patient the rectum much more quickly refused injections. All other means of affording relief being taken away, as stated in describing the case, I was brought to the consideration of gastrotomy. I had previously ventured to suggest this operation in impending starvation, in cases of perforation or communication between the Œsophagus and trachea, where deglutition is sometimes impossible. Death in these cases sometimes takes place simply from inanition. In this instance, it was well known that there was incurable disease, and that any operation which might be performed would only be palliative. It was submitted to the patient, that such an operation might be quickly fatal, or prolong his life for a few weeks; even with such a slight hope he most readily assented, so terrible was the sense of starvation and of thirst. These symptoms were relieved, and the horrors of such a death partially mitigated. In the treatment of the case after the stomach had been opened, it might have been

well to have repeated nutrient injections by the rectum, and to have given food less frequently, although the quantity introduced at each time was only two ounces. There was fear lest the operation might suddenly terminate the flickering flame of life, and lest no rallying should take place; and afterwards, lest the sutures should give way, and thus the contents of the stomach be freely extravasated into the peritoneum. After twenty-four hours, faintness came on; the patient was evidently sinking, and any treatment would have been alike ineffective. Stimulants were given very freely, and at each time were followed by a slight revival in the action of the heart. There appeared to be nothing to call for the use of opium; but, at last, a small quantity of tincture of iron was administered, as being one of the most powerful stimulants. The consideration of the complete particulars of this case lead to the conviction, that if the operation had been performed earlier more permanent benefit might have accrued. It was done with a comparatively trifling addition to the sufferings of the patient; it was effected with ease, without collapse or peritonitis; and the thirst and sense of starvation were relieved in a degree which were scarcely anticipated. In cases of equally advanced starvation, death has ensued as quickly; and it is probable, that had the operation not been performed, life would have terminated even more speedily. The patient would certainly have been deprived of the relief which for twenty-four hours he experienced. Under these circumstances we should strongly urge a repetition of this operation, if a favorable case were presented, but at an earlier period of the disease.

Mr. Forster has a second time performed this operation on a child for occlusion of the Œsophagus from the action of caustic alkali; but so feeble had the child become that the stitches between the coats of the stomach and the skin gave way, and extravasation took place into the abdomen.

CASE XXXII. Cancer of the Œsophagus. Sloughing. Perforation of the Aorta. Sudden and Fatal Hæmorrhage.—Margaret H—, æt. 60, was admitted, April 17th, 1861, under Dr. Wilks's care, in a prostrate condition; she made no especial complaint, but had worked hard, and appeared worn out, rather than to be suffering from any positive disease. About a fortnight after admission, she spat up a little blood; but, on careful examination of the chest,

no disease could be detected. Two days before death, she again spat up a little blood, and appeared very prostrate. She rapidly sank. On inspection, a circumscribed sloughing cancer was found in the centre of the œsophagus; the disease extended into the mediastinum, and involved the lung on the left side. The sloughing had extended into the aorta, the walls of the vessel were perforated, and the stomach was distended with blood. It could not be ascertained that she had ever suffered from dysphagia.

The diagnosis of this case was very obscure, and even when the first oozing of blood came on it was supposed to have been poured out from the lungs; the hæmorrhage from the aorta led to fatal syncope; the blood, however, passed downwards into the stomach, and was not rejected by the mouth. The absence of the dysphagia increased the obscurity of the symptom.

Polypus at the commencement of the Œsophagus. In the 'Medico-Chirurgical Transactions' (vol. xxx), a case of this kind is recorded by Dr. Arrowsmith. A lobulated growth, freely moveable and about the size of a walnut, was attached to the mucous membrane of the œsophagus, and produced fatal dysphagia. The tumour was vascular and homogeneous in structure.

Obstruction of the Œsophagus from pressure of aneurismal or other tumours. Tumours of a cancerous, strumous, or aneurismal character sometimes exert pressure upon the œsophagus, and cause dysphagia. Cancerous and strumous diseases of the glands occur more frequently in the anterior than in the posterior mediastinum, and in the former situation they do not exert pressure upon the œsophagus; but where those at the root of the lung, or at the lower part of the neck are enlarged, there is closer proximity with the œsophagus; and pressure upon, or sup-puration communicating with the canal may be the result.

The œsophagus is also in close contact with the aorta, and we frequently find that dysphagia is one of the symptoms of aneurismal disease of that vessel; the dysphagia is, however, less severe and constant than in direct constriction of the œsophagus. It is no uncommon thing to find death suddenly taking place from rupture of the aneurism into the œsophagus; but a considerable time may elapse, as in an instance recorded by Mr. S. Cooper, in which eight weeks elapsed after the first rupture before the fatal hæmorrhage occurred; in other instances death

follows from the condition of the heart, or from pressure on the respiratory and laryngeal nerves, producing fatal syncope or apnoea, even although the pressure on the œsophagus may have been sufficient to produce sloughing. As to other symptoms, they are, pain at the sternum, at the side, between the shoulders, and very frequently down the arm; paroxysms of dyspnœa often cause intense distress; and raucedo from pressure on the laryngeal nerves is by no means unfrequent. The sounds of the heart are frequently modified, and if any pressure be exerted on the bronchus, it is manifested by the less free admission of air into the lung on the side of the obstructed bronchus. The pain, dyspnœa and dysphagia, in some of these cases, are much relieved when the patient bends the body forward, so as to remove the pressure from the structures beneath. In an instance mentioned by Mr. Armiger, in the 'Medico-Chirurgical Transactions' (vol. ii), the patient was most easy on his knees and elbows.

The *diagnosis* is often very obscure; the emaciation is not generally so great as we find in cancerous disease, but the paroxysms of dyspnœa and pain are more marked and are exceedingly severe, although the patient is at times able to swallow with ease. The *treatment* is simply palliative; yet much may be done for the comfort of the patient and the prolongation of life, by the regulation of the diet, the avoidance of mental and physical excitement, and the occasional use of sedatives and anti-spasmodics, as of morphia, chloric ether, stramonium, &c., to relieve the paroxysms of dyspnœa. A word of caution must be given in reference to the use of bougies, for I have seen several instances in which the attempt to pass a bougie would have at once broken down the communication with the aneurismal sac, and have led to sudden fatal hæmorrhage.

CASE XXXIII. *Aneurism of the Aorta and Sloughing Œsophagus.*—James F—, æt. 34, was admitted, November, 1855, and died in January, 1856; he was a temperate man, married, and a labourer at Dartford. Six months before his admission, after having been engaged a short time previously in carrying very heavy weights, he experienced pain in the left breast; this pain became much more severe, and also extended between his shoulders, but there was no tenderness in the back. December 4th, the pain at the left nipple became more fixed, and there was a slight systolic bruit. January 1st, it was noticed that the radial pulse was weaker on the right side, and he was found to have difficulty in swallowing solids. This dysphagia increased

in severity, and his dyspnœa became more distressing. January 20th, he was unable to swallow food; his face was livid, dyspnœa urgent, and his pain severe. He died on the 25th. On examining the chest, the lungs were emphysematous, pale, but moderately collapsed. There was acute inflammation of the pericardium, and considerable injection of the pleura on both sides. On turning aside the lungs, an aneurismal tumour, about the size of a large orange, was found at the termination of the arch of the aorta; its walls were thin; the posterior part of the vessel was entirely destroyed, and communicated with a cavity in front of the vertebræ, one of which was absorbed. There was scarcely any fibrin in the sac. The aneurismal tumour had pressed upon the œsophagus, and quite obliterated its canal; the whole of its walls were of a greenish colour, very offensive, and in a sloughing condition. Still no perforation had taken place. Both bronchi were compressed. Two other aneurismal tumours were found connected with the ascending and transverse portions of the arch of the aorta. Other viscera were healthy.

CASE XXXIV. *Aneurism of the Ascending Aorta Rupturing into the Pericardium. Communication of the Œsophagus with the Left Bronchus.*—Frederick K—, æt. 23, was admitted under Dr. Gull's care, January 23rd, and died April 26th, 1856. He was a hawker, and had been living in the Old Kent Road; he had enjoyed good health till five months previously, when he struck his chest against a box hanging from a crane; a fortnight afterwards he experienced pain at the part; this gradually increased till three weeks before admission, when he was obliged to give up work. On admission, he complained of pain in the chest; a distinct pulsation could be felt between the second and third ribs on the right side, and a jar with the second sound of the heart. There was pain at the seat of pulsation, and along the border of the pectoralis major, and down the inner side of the arm. The pain continued severe, and a systolic bruit became audible at the seat of the tumour. He could obtain no rest at night. On April 19th, he had difficulty in swallowing, which increased in severity. On the 28th, after talking with his friends, he died very suddenly. On removing the sternum, an aneurism of the ascending aorta was opened; it had extended to the sternum on the right side. On opening the pericardium, it was discovered to be full of blood, and a small irregular communication was found at its upper part with the aorta. The heart was of normal size; the left ventricle was not hypertrophied; the valves were healthy. The ascending aorta formed an aneurismal sac, about two inches and a half in diameter, principally on the right side. The lung was adherent, and it was nearly perforated. The aneurism extended as far as the origin of the left carotid; below the left subclavian was another small dilatation. At the centre of the œsophagus was a slough, and an opening had been formed into the left bronchus; there was no communication, however, with the aorta. The remaining viscera were healthy.

Although it appeared that the greater pressure from the aneurism, was on the right rather than on the left side, we can find

no other explanation for this sloughing condition of the œsophagus, and its communication with the bronchus, beyond the pressure which all these parts suffered from the distended aneurismal sac.

CASE XXXV. Aneurism. Pressure on the Œsophagus and on the Left Bronchus. Difficulty in Deglutition. Sudden Death.—James F—, æt. about 40, was admitted, under Dr. Addison's care, August 24th, 1859, and died September 24th, at 6 p.m. He was a spare man, pale, but strongly built; for nine months he had suffered from pain at the sternum and between the shoulders; he had lost flesh; and for several months had suffered from difficulty in deglutition, so that on admission he was unable to take solid food; but fluids were easily swallowed. A few days after he had been in the hospital, an attack of very urgent dyspnœa came on, but gradually subsided, leaving the respiration easy and of ordinary frequency. In that state he remained till the day of his death. The physical signs were obscure; the resonance on the left side was very slightly diminished; the respiratory sounds on that side were much less distinct than on the right; and were accompanied at the base with slight mucous râles; the voice was not increased in resonance. No abnormal pulsation nor sound in the interscapular or sternal regions could be heard. The heart sounds were normal as to rhythm; but the second sound was "ringing;" the pulse was compressible, more feeble on the left side. He had no vomiting, neither was there any evidence of enlarged glands in the neck, &c., nor had he any pain in the arm. On the 24th, after taking his tea, he washed up the things for the nurse, and immediately afterwards he was seen sitting on the edge of his bed, unable to speak, and scarcely able to breathe. He never spoke afterwards, but died in about a quarter of an hour, the heart continuing to beat after he had ceased to breathe. *On inspection.*—About one inch below the left subclavian artery was an oval opening into the aorta, with rounded edges, extending into a large cavity, about four inches in diameter; this cavity was bounded posteriorly by the vertebræ, which were corroded; the tumour pressed upon the left bronchus, and upon the œsophagus, and probably upon the thoracic duct. The lungs were healthy. The bronchi on the left side were rather more congested than usual. The ascending and the arch of the aorta were atheromatous; the valves and the muscular fibre of the heart were healthy; the pericardium was normal, and no disease could be found in any other part.

The sudden fatal apnœa was due, probably, either to injury of the laryngeal nerves and spasmodic action of the larynx; or to sudden distension and increased pressure on the bronchus. The diagnosis was very obscure; the symptoms were slight, and the interference with the entrance of air into the left lung might have been regarded as arising from a tumour in the mediastinum.

Before concluding the subject of dysphagia, I must refer to its occasional presence in pericarditis and pleuro-pneumonia, as mentioned by Dr. Stokes, in his valuable work on 'Diseases of the Heart.' Although very numerous instances of these diseases come under my observation, I do not remember ever having witnessed difficulty in swallowing associated with the other symptoms of pericarditis; and even, when the pericardium was found distended with 36 oz. of pus, and nearly filling the left side of the chest, this symptom had not been observed: in a patient recently under my care, extensive pleuritic effusion was found with dysphagia; but close watching of the case led me to believe that the latter symptom had its origin in local disease. Transient enlargement of the bronchial glands might, however, induce the dysphagia in these cases, and to quote the words of Dr. Stokes, it may probably "be less the result of a mechanical condition, such as pressure on the œsophagus, than of some excited irritability either of that tube, or of parts immediately in contact with it."

Dilatation of the Œsophagus. Pouch.—Three forms of dilatation of the coats of the œsophagus and pharynx are met with; in the 1st the canal is uniformly dilated; in the 2nd a pouch or bag is formed, consisting of all the coats; and, in the 3rd, there is a hernial protrusion of the mucous membrane penetrating through the muscular coat.

The *first* is found occasionally present in organic occlusion of the œsophagus, especially where it is of a non-malignant character, and of slow formation. Mayo, in his 'Outlines of Pathology,' narrates a remarkable instance of œsophageal dilatation. "Mary B—, æt. 33, ill for ten years, was in a state of extreme debility and emaciation; the food was returned in three or four minutes mixed with mucus, and death took place from inanition. The œsophagus from its junction with the pharynx, which was rather less capacious than usual, enlarged to an extraordinary degree of dilatation. The greatest width that it attained exceeded two and a half inches when distended, and occurred about four inches above the cardia. The tube then narrowed more abruptly, so as to render the cardiac termination like the pharyngeal of nearly the usual dimension. The structure of the cardiac end for about an inch, and that of the pharyngeal

end for about an inch and a half, were healthy. Intermediately the lining tunic was thickened and opaque; the mucous membrane had the appearance of having yielded or opened into flat shallow depressions, which above followed a longitudinal direction, and below formed irregular pits. At the depressed surfaces, the membrane had the natural colour; between them, it was opaque and whitish. The muscular fibres were of the natural colour and thickness; they had grown with the expansion of the canal."

In vol. x of the 'Pathological Transactions,' Dr. Barker has recorded an instance of dilatation of the œsophagus commencing two inches below the rima, and forming an ovoid swelling the size of a swan's egg in the posterior mediastinum. There was strumous deposit in the bronchial glands forming masses as large as the fist, and firmly adhering to the front of the œsophagus and of the trachea, and connected with fibrous tissue surrounding the origin of the great vessels. There was also a perforating ulcer in the duodenum, communicating with a small abscess in the peritoneum.

As to the *second* form, consisting of a muscular pouch, Mr. W. C. Worthington, in vol. xxx of the 'Trans. of the Royal Med. and Chir. Society,' narrates an instance under his own care. A gentleman, æt. sixty, three years before his death, suffered from slight dysphagia, which gradually became extreme with progressive emaciation; and for three weeks he was sustained by injections. The œsophagus was constricted at the commencement, so that an urethral bougie was alone capable of being passed. Opposite the cricoid cartilage, between the trachea, the œsophagus and the cervical vertebræ, was a pouch three and a half inches in length, like the finger of a glove, two thirds of it were covered by the muscular fibres of the constrictors; the opening into the pouch was free. The mucous membrane of the œsophagus at the constricted part was healthy. The same author quotes from Sir Charles Bell an instance of Mr. Ludlow's of muscular pouch, between the œsophagus and vertebræ, in a man æt. 60, who had dysphagia for five years; and also mentions from Sir Charles Bell one of the *third* form of pouch, consisting only of mucous membrane protruded through the muscular layer. This last form may be free from symptom. A case of

this kind recently occurred at Guy's Hospital; J. C—, æt. 28, was brought in a dying state with peritonitis from perforation of the ileum during typhoid fever. At the commencement of the Œsophagus was a pouch, half an inch in length, consisting of the mucous membrane, and very slightly covered at the neck of the pouch by a few muscular bands; it was full of mucus. (Prep. 1784⁷².)

Minute warty growths are sometimes observed in the Œsophagus; but, I am not aware that any symptom is indicative of their presence; they are, probably, of similar pathological origin to some of the non-cancerous growths occasionally found attached to this part of the mucous membrane. (Prep. 1784⁷⁷.)

Gastric Solution.—In studying the diseases of the Œsophagus, *gastric solution* of its lower extremity must be borne in mind. This subject has been very clearly brought forward in the communications to the 'Guy's Reports,' by Mr. Wilkinson King, in the years 1842 and 1843. It is exceedingly frequent to find the mucous membrane of the Œsophagus abruptly terminating at the cardiac extremity of the stomach, from the solvent action of the gastric juice having extended to that line; but on opening the canal of the Œsophagus itself for several inches near its lower extremity, the upper margins of the rugæ are often found deprived of mucous membrane; and long shreds are observed on stretching out the tube, these portions having escaped digestion. This solution extends into the mediastinum, as found in cases mentioned in the communication just referred to, or into the pleura itself, the contents of the stomach thus escaping into the left pleural cavity, which is in closer relation with the Œsophagus than the right pleura.

Only two cases of this perforation of the Œsophagus have occurred at Guy's during the last few years, one in a case of fever, another of hydrocephalus, so that it is a circumstance of unfrequent occurrence. Mr. E. Canton, in the 'Lancet' of 1859, gives an instance of an infant, æt. 2 months, who died comatose, insensibility having come on two hours after the ingestion of breast milk and soaked bread. An oval opening three-fourths of an inch in length was found on the left side of the Œsophagus, the rent commencing a quarter of an inch above the diaphragm. Its edges were thin, flocculent, and irregularly

fringed; a second aperture also was found separated from the other by a small strip of undissolved texture. The causes of gastric solution are now more clearly understood than formerly. The position of the body, the development of gases in the intestines pressing upon the contents of the stomach, the non-contracted state of the œsophagus itself, are causes which produce the passage of the gastric juice into the œsophagus. Sometimes, indeed, this pressure forces the contents into the pharynx, and we find them gravitating into the trachea and bronchi.

Ecchymosis.—Hæmorrhage from the œsophagus generally arises from the rupture of aneurismal tumours, or from cancerous disease; but in cases of fatal purpura, we sometimes find the whole mucous membrane covered by points of effused blood, and blood is also effused into the surrounding cellular tissue. This part, however, is affected only in common with the whole mucous surface of the alimentary canal, as well as with other membranes and gland-structures.

CASE XXXVI.—*Rupture of the œsophagus.*—The following case warrants the belief that rupture of the coats of the œsophagus sometimes takes place during life; the specimen is in the Museum of Guy's (No. 1799⁴⁰).

M. C—, æt. 24, a cabinet-maker, of intemperate habits, attended a public supper in September, 1842: during supper he felt sick, and left the table; he vomited slightly, and returned home with assistance. He then took a dose of castor oil; at two in the morning he complained of severe pain across the epigastrium, and great difficulty in breathing; the abdominal muscles were rigid, the respiration laborious; the patient was found sitting up in bed, leaning forwards on his hands; his countenance was anxious, the pulse soft, the bowels had not acted; an emetic of antimony and ipecacuanha was administered, but without effect; at 7.30 a.m. there was less pain, but increased dyspnœa, and there was emphysema of the face and throat. The stomach-pump was used, but without effect, and he died at noon. On inspection a large rent was found in the œsophagus at its lower part, filled with ingesta, which were also extravasated into the left pleura; the pleura also contained castor oil. The stomach and intestines were exceedingly distended with flatus; and the stomach partially dissolved by gastric juice. The rent in the œsophagus appears in the preparation to extend into the stomach, but was perhaps increased after death.

It is probable that the œsophagus was much dilated with

food, and that its coats were softened either by previous disease, or by digestion from gastric juice regurgitated into it from the stomach, and there remaining sufficiently long to corrode its walls. There is no evidence that the stomach-pump increased the rent, for the castor oil which was found in the pleura was taken several hours before the stomach-pump was used; still, if it had been known that such a rent had existed, this remedy would not have been applied; the severity of the symptoms rendered it probable that some poisonous substance might have been taken with the food, and the emetic had failed to act; under the circumstances which existed the use of the stomach-pump probably tended to relieve rather than aggravate the symptoms.

Mayo quotes a case from Boerhaave of rupture of the Œsophagus after an emetic had been taken by a robust, but gouty man; a rent one and a half inches in length was found communicating with the left pleura; and the fatal result took place twelve hours after the emetic had been administered.

CHAPTER III.

ORGANIC DISEASE OF THE STOMACH.

ALTHOUGH it is probable that every change of function is marked by physical difference, still very many of the local alterations of structure are of a character so transient, and so completely beyond recognition by the senses, that we are compelled to separate them from others in which the structure of the part is more evidently modified. This uncertain division is especially found in the study of stomach diseases, for their greater number are of the kind in which no structural lesion can be traced; but the chemical, anatomical, and physiological researches of late years have diminished the class of simple functional disease, and have greatly increased the knowledge of organic lesion.

The following conditions of the stomach are met with :

Atrophy and Hypertrophy of the Mucous and Muscular Coats.

Solution by the action of the Gastric Juice.

Inflammation.

Catarrh.

Diphtheritic Inflammation.

Suppuration in the Walls of the Stomach.

Ulceration, Superficial, Follicular, Chronic, and Perforating.

Fibroid Disease of the Pylorus.

Cancerous Disease.

Atrophy of the Mucous Membrane.—Wasting of the mucous membrane of the stomach takes place in common with that of other organs and glands, and the subject has been elucidated by Dr. Handfield Jones, whose microscopical investigations have directed particular attention to the subject.

The mucous membrane of the stomach consists principally of

small glands or follicles, which open into minute pits on the surface, and secrete the gastric juice, the most important solvent of food. The follicles rest on a stratum of cellular tissue of varying thickness, which contains a very large quantity of elastic fibre. The delicate capillary branches of the blood-vessels, derived from the coronary, hepatic, and splenic arteries, extend between these gastric follicles nearly in a straight course, and they form a beautiful plexus of vessels around the minute crypts, and also beneath the follicles themselves. These structures are easily observed under a low magnifying power of the microscope, and in a portion of congested membrane present a beautiful appearance. The sympathetic nerve filaments are also seen at the base of the mucous membrane, sometimes upon the capillary vessels, and at other times apparently leaving them; but their ultimate division I have not been able to trace. There is no doubt that this nerve forms an important element in the structure of the membrane.

The surface presents columnar epithelium and mucus, and the follicles contain spheroidal epithelium and nuclei. As in every other gland, these minute and simple ones appear to have varying degrees of functional activity, and undergo degenerative changes. Thus in many cases of fatal disease, with gradually increasing exhaustion, only a small quantity of food is taken for many days before death, whilst, in other instances, the appetite is maintained to the last; we consequently often observe, in the one case, the follicles full of secreting cells and nuclei; in the other, they are comparatively empty.

Microscopical investigation has done much to increase the knowledge of pathology; but, with increase of microscopical power we must add equal caution to remove all the causes liable to mislead us. The mode which I have adopted in preparing sections, and which will generally be found a successful one, is to stretch the membrane over or between the fingers, and then, by means of Valentin's knife, make a section of the required depth and thickness. This is afterwards removed by scissors, and spread out in water by needle points. I have examined with great care a considerable number of stomachs from the post-mortem table of Guy's Hospital; but, it is not necessary to mention the cases in which the membrane appeared in a healthy

condition. In many of these examinations I have observed appearances precisely corresponding to the descriptions and drawings of Dr. Handfield Jones; but, I think we must carefully consider the manner in which many of these appearances may be produced, by the mode of making the preparation, or by changes after death. I refer to wasting of the follicles, nuclear deposit around them, and the development of cysts. The gastric follicles change very rapidly after death; and in a short space of time nothing can be observed but the termination of the follicle itself upon the submucous areolar tissue, and above this, an irregular aggregation of granules and nuclei. The basement membrane also rapidly becomes dissolved; and this condition will be found, on microscopical examination, before the ordinary appearances of gastric solution are observable in the stomach. The greater curvature of the stomach is in this way generally too much changed to allow us to place much dependence upon its microscopical examination; and for this reason, it is evident, that we have to avail ourselves of portions of membrane above the line of solution. That, from the lesser curvature, however, and from the pyloric region, is less generally dissolved by the gastric juice, and is also the part most subject to morbid changes; but, the cardiac portion should be also examined when possible. Not only does the membrane become dissolved, but, in some cases, by decomposition, it becomes emphysematous, and presents minute vesicles and blebs, which occupy the substance of the tissue. The appearance of the very minute emphysematous vesicles in the mucous tissue precisely resembled the appearance of well-defined cysts, surrounded by nuclei, in Case XXXVIII, recorded in this chapter. This is one source of fallacy; and another will be found in the fact that nuclei are readily separated from the follicles in the preparation of the section, and become diffused between the structures. The contents of the follicles are easily detached; and, by the action of acetic acid, of cold water, or by mere pressure, a perfect cast of the follicles will be often extruded, and project from the surface of the membrane. A third fallacy, which may considerably mislead us, is the appearance of the mucous coat, altered by the state of contraction of the submucous and muscular coats beneath. The mucous coat will expand to the largest amount of distension that the

muscular coat allows. When the muscular coat is contracted, the usual appearance of rugæ is presented; but, a further contraction produces a mammillated appearance of the membrane. This may be sometimes observed, after removing a portion of healthy mucous membrane, a short time after death; when, by immersing it in cold water, for a few hours, this state of mammillation is produced. A thickened, chronically-inflamed membrane will, I believe, present *true* mammillation of the stomach; but in that artificially produced, the manner in which the fissures extend nearly to the submucous cellular tissue, might lead us to attribute this appearance to a morbid contraction of the membrane itself. Dr. Handfield Jones gives, in his observations on the stomach, an original and interesting account of the production of mammillation; and he attributes these depressions to wasting of the membrane, the breaking up of nuclear masses, and to the contraction of the tissue beneath. This opinion requires confirmation, for, as far as my observations have gone, it would appear that mammillation is more common than the existence or evidence of solitary glands or separate nuclear deposits in the membrane; and that this appearance of simple mammillation may be easily produced artificially in a healthy mucous membrane. A fourth fallacy may arise from the direction of the section. The surface of the stomach being not that of a plane membrane, and its follicles opening into crypts, an oblique section may readily give the appearance of fibrous tissue abnormally developed, where such does not really exist.

The whole of the coats of the stomach are sometimes exceedingly wasted, but in fatty degeneration or atrophy of the mucous membrane this is not generally the case. There are several degrees of this wasting or fatty change. Thus, sometimes the cells of the follicles, instead of presenting a simple nucleus, contain a great number of minute highly refracting particles, and almost resemble an inflammatory granule cell, while the appearance of the stomach itself is otherwise in a perfectly healthy condition; although these cells are also found in other states, as extreme congestion with superficial ulceration, &c., they appear to indicate a diminution of vital activity, rather than an excess of it. At other times, the stomach is found to be pale, and here and there studded with white points, somewhat

resembling solitary glands, but not at all elevated above the surface. A horizontal section, in such a case, shows around the crypts at this whitened portion, minute highly refracting granules and fatty particles; and a vertical section presents a dark border on the surface, consisting of the same elements; these are also sometimes observed, more or less distinctly, beneath the follicles.

A more advanced condition of atrophy shows the follicles to be entirely destitute of secreting cells, and only containing granules of fat, or perhaps wholly destroyed, with irregular patches of pale mucous membrane. This state was well marked in the first briefly-recorded case.

Beside these forms and degrees of atrophy, which may be called secondary, there are others which arise from chronic inflammation of the membrane, in which the structure appears thickened, dense, and the mere rudiments of gastric follicles remain. These may arise from fibroid degeneration or cancerous disease slowly encroaching upon the membrane adjoining it, and thus leading to atrophy and degeneration.

The symptoms observed in some of the cases in which this fatty change in the mucous membrane of the stomach existed, were a sense of great prostration and exhaustion, with complete loss of appetite. The tongue was clean; there was no pain, neither was there thirst, nor vomiting, but an inability to take food; vomiting has sometimes taken place, but possibly from other causes. This form of atrophy has been observed in phthisis, in struma, in exhausting suppuration, and is often associated with a fatty condition of the liver.*

CASE XXXVII.—*Strumous Disease of the Third Rib. Phthisis. Ulceration of the Intestine. Strumous Disease of the Kidney, Ureter, Bladder, Prostate Gland, Vesiculae Seminales, Spermatic Cord, and Testis. Fatty Degeneration of the Stomach and Liver, &c.*—John S—, æt. 50, was admitted under Dr. Barlow's care, September 27, 1854, in an almost dying condition, but he was apparently suffering from phthisis; no history could be obtained, and he died October 2nd.

The case is a good illustration of a strumous subject dying from exhausting disease, almost every organ and mucous membrane being affected with

* Handfield Jones.

struma; the stomach, however, presented only atrophy and fatty degeneration, probably dependent on the exhaustion and diminished nervous power.

Inspection was made forty hours after death. The body was much wasted. A fistulous opening passed from the third costal cartilage into a pleural abscess. The lungs were disorganized and tubercular.

The *stomach* was flaccid: its mucous membrane was covered with a thick layer of mucus; and it presented, especially towards the pyloric extremity, several opaque white patches, about a quarter of an inch in circumference. These parts were found to consist of degenerated mucous follicles. The follicles had their usual outline, but were filled with minute fat-particles, and were destitute of secreting cells.

The small and large *intestines* had been ulcerated, and long cicatrices extended for several inches. The mucous membrane was congested and much puckered, and the muscular coat hypertrophied; the rectum was similarly affected, but in a less degree. Some of the mucous follicles in the cæcum were examined, and were found to contain abundant secreting cells: in this respect, they were very different from those of the stomach.

Although the stomach in this case presented marked evidence of diminished functional activity in common with the general wasting; still, in those parts of the intestine in which there was obstruction to the free passage of the contents, on account of the contraction of cicatrices, the muscular coat was hypertrophied.

CASE XXXVIII.—*Thinning of Mucous Membrane, with the appearance of minute Cysts (probably post-mortem) from the Stomach of a man who died five weeks after taking a solution of Chloride of Zinc.*—He was an Irish labourer, whom a fellow-labourer induced, as a joke, to swallow some of the solution of zinc. Its imbibition was followed by vomiting, promoted by an emetic. After his admission into Guy's Hospital, emaciation rapidly came on, and he died in five weeks.

Examination thirty hours after death.—The stomach was distended and injected; at the cardiac extremity and greater curvature several gray lines of discoloration existed; internally, it was generally covered by a layer of mucus. The rugæ were not observable. The lining membrane was *emphysematous* at the greater curvature. Throughout the stomach the mucous coat was thinned, while the muscular one was hypertrophied. At the pylorus, the mucous coat regained its usual thickness, and presented a frilled margin, as at the circumference of an ulcer.

On the examination of the mucous membrane, above the emphysematous line, the ends of gastric follicles were observed, but they were not covered with the usual thickness of membrane. Near the pylorus, the membrane appeared equally thin; but also presented numerous very minute transparent vesicles, which projected upon the surface of the membrane, and appeared to consist of a cyst wall, containing fluid and nuclei. The cyst, when ruptured, presented a halo of fluid and granules around it. Other more minute cysts were found in the substance of the membrane, particularly towards the greater curvature. They were about 1-30th to 1-10th of an inch in diameter.

At first, it appeared that the development of cysts in this case had taken place before death; but the greater curvature presented large blebs of air, and the smaller vessels were, no doubt, of a similar character. The separation of the elements of the membrane by the development of gas had given rise to this deceptive cystic appearance. I have observed a similar emphysematous condition of the mucous membrane in some other cases; and it probably arises from rapid decomposition taking place in connection with partial gastric solution, modified, perhaps, by a diseased condition of the membrane.

It occasionally happens that we find structures resembling solitary glands of the intestine in the mucous membrane of the stomach. Thus, a short time ago, in examining the stomach of a child who had died from chorea, I found the whole membrane presenting numerous whitish specks, which consisted of these structures imbedded in the substance of the membrane. There was, however, no evidence that such glands existed in the stomach of the man poisoned by zinc.

Hypertrophy.—An apparent thinning of the mucous membrane of the stomach is consequent on great dilatation of the viscus; and conversely the lining membrane seems to be preternaturally thickened when the stomach is contracted, and large rugæ are formed by the inversion of the mucous membrane; and, as we have before remarked, simple contraction of the submucous tissues causes a mammillated appearance, resembling that produced by chronic change. Still there are instances arising from chronic irritation and inflammation, and from prolonged con-

gestion, which we shall have to refer to when speaking of catarrh, in which the mucous membrane is both thickened and hypertrophied. In other cases, indeed, polypoid masses are formed by the mucous membrane, and I have observed large folds of this kind surrounding the pyloric orifice, without evincing symptoms of disease during the life of the patient.

The more marked hypertrophic changes which take place are those connected with the muscular coat, when there is obstruction at the pylorus; the muscular tissue then becomes stronger and thicker, at first in the proximity of the obstruction; the hypertrophy gradually extends its area, reaching two or three inches, and even over the whole stomach. This change is a preservative one, and it tends, though with less and less efficiency, to force the dissolved aliment through the diseased part; but at length the wasting, consequent on the system not receiving its proper supply of nutriment, becomes extreme, and the patient succumbs.

Post-mortem solution.—The secretion poured into the stomach from its numerous follicular glands has a very important function to perform in dissolving the nitrogenous portions of food. This secretion, the gastric juice, is a clear, somewhat viscid fluid; it has an acid reaction, from the presence of hydrochloric or lactic acid, and it contains an organic substance, pepsin, in small proportionate quantity. By the mutual reaction of these agents on the organic animal principles of ordinary food, assisted by the temperature of the body and the churning movements of the stomach, the solution of the protein compounds and of the gelatin and chondrin takes place, and a fluid is formed containing peptone compounds, as they are called, and which is discharged into the intestine through the pyloric valve. The solvent power of the gastric juice is of a simple chemical kind, and it can be exerted external to the living organism when the necessary conditions are carried out; it would act also upon the structures of the viscus itself, even during life, but is prevented by the protective covering of mucus and epithelium, constantly renewed by the living power of the part. After death, however, the chemical action is unchecked, and the walls of the stomach are dissolved; sometimes, indeed, with great rapidity, and in every instance to some extent, so that pathological researches are

interfered with, and the appearance of the stomach is necessarily modified.

John Hunter drew attention to post-mortem solution in connection with diseases and injuries of the head, and T. Wilkinson King, of Guy's, added definite facts in reference to the degrees and position of the solution; and Dr. Budd, in his treatise on Diseases of the Stomach, has still further and very fully elucidated the subject. The gelatinous softening which has been described by Andral, Cruveilhier, &c., as occurring during life, is now generally believed to be a form of this solution, and in this opinion I concur, although a different one is held by some talented pathologists.

It must be always borne in mind, that after death blood gravitates into the most depending vessels, that exosmosis takes place, and chemical action exerts its influence, unchecked and unmodified by vital action.

The amount of gastric solution depends in part on the quantity of gastric juice actually in the stomach at the time of death. Gastric solution is especially manifest when sudden death occurs during the process of digestion; and it is shown still more as a sequence of cerebral diseases, especially those of an inflammatory character in young subjects, the follicles having been stimulated to pour out secretion at irregular times, and in excessive quantity. Time is required for the solution, and the action proceeds more rapidly in summer than during the coldness of winter. The stomach is sometimes found completely perforated, although food has not been taken for several hours before death.

The simplest condition of this change is thinning and softening of the mucous membrane, so that it is with great readiness detached; if the blood-vessels be empty, the colour is pale, and it has a semi-gelatinous appearance; generally, however, the vessels contain blood, which gravitates into the most depending vessels; the hæmatine exudes into the substance of the stomach itself, and greenish-brown or almost black lines are formed in the course of the vessels, at the whole of the dissolved part, by the action of the gastric juice on the colouring matter of the blood. If the transudation have taken place into the cavity of the stomach, a greenish-brown fluid is produced by a similar action. This solvent action may be so slight that it is only

detected when we examine a section of the membrane with the microscope, or the mucous membrane is exceedingly thinned, or entirely destroyed; the submucous and the muscular coats are then dissolved, and at last the peritoneum is reached. The serous membrane occasionally gives way, so that a ragged perforation is formed, and the contents of the stomach transude into the peritoneal cavity. The adjoining viscera then become acted upon, unless adhesions exist which have obliterated the cavity, as we find in strumous peritonitis.

The extent of the dissolved part is generally marked by a defined line, showing the level to which the solvent fluid has attained. This is generally along the greater curvature; but sometimes, from the position of the body, we find that the solution is greatest in the region of the lesser curvature, or even that the duodenum is especially acted upon; and this part of the intestine may be perforated while the stomach is intact. Or from the evolution of gases, position of the body, &c., the fluid is pressed into the œsophagus; the mucous membrane of that canal dissolved, and sometimes all its coats perforated, so that the contents of the stomach are found in the pleural cavity.

John Hunter attributed these effects to the fact that chemical action is unchecked by the vital state of the parts; but Dr. Bernard and Dr. Pavy* have demonstrated that the gastric juice will act upon living tissues, as shown by introducing a rabbit's ear and the leg of a frog into a gastric fistula, thus proving the protective influence of the gastric epithelium and mucus. These experiments, however, are not conclusive, for the circulation could not be carried on in the usual free manner.

When the anterior part is acted upon, Dr. Budd† explains the fact by the small quantity of gastric juice, which was in the greater curvature, being neutralized either by ammonia evolved, or by the exudation of alkaline serum from the blood, or from dropsical effusion; whilst the small quantity on the anterior part has not been thus neutralized. The action of the gastric juice, Dr. Budd states, may be checked by alcoholic liquors, or by medicines administered before death. We are not acquainted fully with the causes of the greater gastric solution in some

* Guy's Reports. Vol. ii. Third Series.

† Budd on the Stomach.

cases than in others, for, whilst agreeing with the author just cited, that it is occasionally very manifest in cases of phthisis, renal disease, typhoid fever, and cancer of the uterus, or disease of organs in which the stomach is functionally disturbed, we shall find an almost equal per-centage of cases of solution when such causes do not exist. It is certainly more manifest in children, and in inflammatory disease of the brain, and is generally more marked in acute than in chronic disease.

Inflammation of the Stomach.—The instances of *acute inflammation* of the stomach, which have come under my own observation, have arisen from poisons, as alcohol, arsenic, oxalic acid, chloride of zinc, sulphuric and nitric acids; in these there are two symptoms which demand particular attention—the absence of pain at the stomach, in most instances, unless perforation have taken place, and the marked prostration of strength, with depression of the pulse.

After irritant and corrosive poisons have been taken, burning pain in the mouth and throat, charring of the mucous membrane, vomiting, irritability of the stomach, purging of blood or of loose fecal evacuations, are produced; and according to the strength of the fluid and its action on the pharynx, œsophagus, and epiglottis, there is dysphagia, or dyspnœa. The vomiting is generally excessive from the primary irritation of the poison, and the vomited matters vary according to the character of the agent and the extent of its chemical action. Subsequently, if the mucous membrane of the stomach or œsophagus have been destroyed, and fibro-plastic exudation have taken place, contraction is the result, and narrowing of the canal follows. The period of the retention of food is then according to the situation of the occlusion, whether in the œsophagus or at the pylorus; in the former case, the food is at once regurgitated; in the latter, it may be retained for several hours. If the absorption of nutriment be thus prevented, emaciation takes place; the whole frame becomes wasted to an extreme degree, and a fatal issue follows in several weeks or months after the poison has been swallowed. I must refer my readers to Dr. Taylor's valuable work on 'Poisons,' for an account of the distinctive symptoms produced by the various corrosive and irritant poisons.

In the following case of poisoning by alcohol, the appearance of the stomach resulted from the irritant action of the poison, but the morbid change was of such a character, that unless especial attention had been drawn to the part, it might very easily have been overlooked. The stomach was minutely injected with arborescent vessels, and the congestion was apparently the remains of an acute erythematous inflammation.

CASE XXXIX.—*Poisoning by Brandy.*—Timothy B—, æt. 27, a vagrant, on January 28th, 1837, was following a man who carried a gallon bottle of brandy. The bottle was accidentally broken, and the spirit spilled in the road. The patient drank the spirit from the ground, by putting his mouth to the earth. A short time afterwards he was observed to lean against a lamp-post, and gradually became insensible: he was taken to the station-house, and, since it was thought to be ordinary intoxication, he was left for several hours; he was then brought to Guy's Hospital about two in the morning; the stomach-pump was used, and some dirty fluid brought up, which appeared to be muddy water. Some coffee was afterwards given to him. He spoke once, asking for water, but died about twelve hours after admission.

The mucous membrane of the stomach presented considerable injection, giving a patchy appearance of rose-coloured spots to the surface; and in the cavity of the stomach was some muddy fluid: the duodenum and upper part of the jejunum contained similar fluid.

The œsophagus was healthy; the liver and kidneys congested. The heart contained a moderate amount of blood in both auricles and ventricles; but it was not distended on the right side as in death from apnœa; the lungs were congested in patches. The membranes and substance of the brain were much congested; but there was no smell of alcohol.

Absence of pain was shown in a marked degree in the case of poisoning by chloride of zinc, from Burnett's disinfecting fluid. **CASE XLI.**

In another case, in which a woman had taken some oxalic acid, the quantity of which, however, was not known, vomiting and prostration were the only symptoms, and the patient gradually recovered.

The following case of poisoning by sulphuric acid is exceedingly interesting in the same respect; but here the absence of complaint of pain can only be partially depended upon, on account of the mental condition of the patient.

CASE XL.—*Poisoning by Sulphuric Acid. Death on the eleventh day. Destruction of the Mucous Membrane of the Lesser Curvature and Pyloric extremity*

of the Stomach. Acute Inflammation of the Colon and small Intestines.—Charlotte D—, æt. 55, was admitted into Guy's Hospital Oct. 5th, and died Oct. 16th, at 11—30; she had been in Bethlehem Lunatic Asylum three times; and several members of the family had been affected with insanity; about seven o'clock on the morning of the 5th, she drank a wine-glassful of sulphuric acid before breakfast, when the stomach was empty. She was brought to Guy's two hours afterwards in a state of collapse, almost pulseless, and her extremities cold; there was no stain in the mouth, and she could not swallow magnesia mixture, which was administered to her; in two hours she became warm, and vomited some grumous, dark-coloured matter. She continued to vomit blood, and to pass blood by stool; and the vomited matter contained sulphuric acid. In the evening she was able to swallow a little milk and arrow-root, and was decidedly better. For the next three days she was very low, but there were no very urgent symptoms, although she continued to pass blood, and occasionally vomited grumous matter. On October 9th, four days after taking the acid, she swallowed without difficulty. She continued in a low condition, able to take a small quantity of food, but suffered from a kind of dysenteric diarrhoea; no blood was passed, but rather pale loose motions, with shreds of mucus. On the day of her death she spoke as usual, got out of bed, but in the evening died unexpectedly, without any one being present at the time.

Inspection seventeen hours after death.—The body was wasted, the eyes were sunken. The calvarium was exceedingly light and thin; the depressions for some of the Pacchionian bodies were so deep that only the external layer of bone appeared to remain. The Pacchionian bodies were large; the brain was atrophied. The mouth and throat were of a whitish colour; and at the posterior part of the throat there was considerable injection, and the mucous membrane of the anterior part of the tongue was whitened. On the posterior pillars of the fauces were white, loose, and irregular patches, which consisted of elongated cells and epithelium. Similar tissue was found adhering to the membrane, at the lower part of the œsophagus. The posterior part of the tongue was covered with ordinary squamous epithelium. The edge of the epiglottis was irregular, minutely serrate, as if it had been injured by the acid and ulcerated. The membrane of the œsophagus was pale, and filled with yellow membranous flakes. The cardiac extremity of the *stomach* was moderately distended; three inches from the pylorus the muscular coat was contracted. The peritoneal surface was not covered by false membrane, nor particularly injected. On opening the stomach it was found to contain about $\frac{3}{4}$ ij of yellow grumous fluid, and a large, thin, yellow membranous mass, which was attached by one extremity to the walls of the stomach; this mass was 4 to 5 inches in length, and as many in breadth; other smaller patches of similar membrane were observed to be partially detached; the surface of the stomach beneath this membrane presented whitish gray tissue, in some parts slightly flocculent. The mucous membrane along the lesser curvature, and for several superficial inches near the pylorus, was destroyed, and the surface was irregularly flocculent. On examining the yellow membranous membrane, it was found to consist of minute particles which highly refracted the light (fat from degenerating

tissue); some indistinct markings resembling gastric follicles were observed, and small blood-vessels were seen filled with altered blood. At the cardiac extremity the mucous membrane was thin and granular; but at the centre of the greater curvature the lining membrane presented numerous nodules, about $\frac{1}{4}$ of an inch in height, the intervening depression being of a dull gray colour. In some of these depressions were adherent yellow tissue, consisting of semi-detached mucous membrane. The raised isolated nodules consisted of mucous membrane, which had not been acted upon by the acid; and showed very distinctly the distended gastric follicles; the follicles were also evident at the cardiac extremity of the greater sac. Along the lesser curvature, and at the pyloric extremity, the section presented involuntary muscular fibre; this was covered over by fibrous tissue, in some parts thin and irregularly flocculent, but near the pylorus forming a thicker layer; this tissue dipped down between the bundles of muscular fibre. At the pylorus were several nodules of undestroyed membrane; but for more than an inch and a half in the duodenum the membrane was destroyed, as at the pyloric extremity of the stomach. The intestine then became gray, and was covered over by a delicate adherent tissue in irregular patches, which were contiguous the one to the other, and generally placed transversely. The jejunum was very much congested, the ileum still more so, and at the lower part of the latter, the mucous membrane was covered by adherent diphtheritic membrane. The whole of the colon was acutely inflamed from the cæcum to the rectum; the mucous membrane was covered by adherent diphtheritic membrane; the submucous cellular tissue was white and thickened (œdematous), and the muscular coat distinct; the diphtheritic layer consisted of imperfect cells, mucus, granules, and highly refracting particles. The rectum was less affected. There was a small ecchymosed spot beneath the endocardium below the left semilunar valve of the aorta. The heart was flaccid, but its cavities contained fluid blood. The liver was healthy, but small. The spleen also was small. The pancreas, supra-renal capsules, and kidneys were healthy.

The symptoms usually following the administration of a large poisonous dose of sulphuric acid, are discoloration and destruction of the mucous membrane of the lips and mouth, intense pain, difficulty in respiration and deglutition, vomiting of bloody grumous fluid, and collapse; and death generally follows in a few hours. The appearances after death vary, according to the strength of the acid; if concentrated, the mucous membrane of the mouth, œsophagus, and stomach, are charred, sometimes even every layer destroyed, and the adjoining structures acted upon.

The case detailed presents many points of interest, in a general as well as in a medico-legal point of view. The acid was taken not by accident, but by a melancholic woman, and the exact

strength of it cannot be ascertained ; but from inquiry, it seemed probable, that the commercial acid was diluted with three or four parts of water, and $\bar{z}ij$ to $\bar{z}iij$ of this diluted mixture taken. The acid had been used for cleaning brass ; and the bottle, previously nearly full, was found emptied of its contents. The symptoms were believed to come on at once, and two hours afterwards, when admitted into Guy's, she was in a state of collapse ; vomiting of bloody, grumous fluid took place, and loose evacuations containing blood were passed. On admission, the mouth appeared neither charred nor discoloured ; and this led some to doubt whether sulphuric acid had been taken.

Death from poisoning by mineral acids takes place either in a few hours, or the patient rallies from the immediate effect, and dies from the severe organic change and inflammation ; or, in a third class, the primary effect is recovered from, but the patient dies from exhaustion, consequent on stricture of the œsophagus or pylorus, or from loss of functional power of the stomach ; the case detailed belonged to the second class ; death did not take place till the eleventh day, from gradually increasing tendency to syncope. In poisoning by sulphuric acid, the bowels are generally constipated ; here the purging of blood was followed by dysenteric diarrhœa.

As to the parts affected by the acid, the mouth, pharynx, and œsophagus were less injured than the stomach, and more quickly and completely recovered themselves ; in none of these parts could the mucous membrane be found destroyed, except at the tip of the epiglottis.

In the stomach, the mucous membrane was destroyed along the lesser curvature, and at the pyloric extremity ; in these parts the muscular coat was covered by irregular fibrous or cellular tissue of greater or less thickness, and was of a slate-gray colour ; and the mucous membrane was separated in the form of a slough. The greater curvature was less affected, but its middle portion presented islets of raised, uninjured membrane between grooves, in which the mucous membrane was destroyed, and in some parts adherent in semi-detached sloughs. It might have been supposed that the part most affected would have been the greater curvature and cardiac extremity ; the stomach was, probably, nearly empty, and the acid passed at once along the lesser

curvature to the pylorus and into the duodenum. The condition of the duodenum was peculiar; it was partially injured, especially near the pylorus, by the chemical action; the rest was acutely inflamed. It may be a matter of doubt how far the acute inflammation of the small, but especially of the large, intestine was the result of the poison; we sometimes find in other cases that the large intestine and stomach are the parts most affected, partly, perhaps, on account of the contents being longer retained in the large than in the small intestine.

CASE XLI.—Poisoning by Chloride of Zinc (Burnett's Disinfecting Fluid). Death in Fourteen Weeks.—Sarah R—, æt. 37, was admitted December 24th, 1856, under Dr. Wilks' care. She was a single woman, engaged in domestic service; her previous health had been good, but her habits were intemperate; twelve weeks before admission she went to visit a friend in the evening, who gave her, in mistake for gin, three-quarters of a wine-glassful of Burnett's disinfecting fluid. The precise strength was not known. She swallowed the draught, and at once felt a painful burning in the mouth, and in about a quarter of an hour vomiting came on, and in half an hour purging; neither the vomited matters nor evacuations contained any blood; the latter were passed involuntarily. Milk and water were administered.

Neither pain nor abdominal tenderness was produced at the time, nor at any period prior to admission. She was conveyed home; her hands appeared swollen, and she staggered slightly. She walked up-stairs and undressed herself; the following day she remained in bed, but was up on the third, and moved about the house. Vomiting was the only symptom; the ejected matters consisted at first of thick, stringy mucus, and afterwards of bilious fluid; in a few days she washed a floor, and in about a fortnight tried a change of air in order to recover her health; but she remained in the same state till admission.

There had been dysphagia for a short time after taking the poison, but this symptom did not recur. It appeared to her, that the food lodged at the scrobiculus cordis.

On admission, she was rather emaciated, not particularly pale, and appeared to possess tolerable strength; the tongue was large and slightly furred; the pulse was compressible; the stomach could only retain food (though of the simplest kind) for a short time; there was no tenderness in the abdomen, which was collapsed; the muscles were rather rigid; the lungs and heart were normal. She had no cough, and appeared comfortable. Carbonate of magnesia, with dilute hydrocyanic acid, were ordered three times a day.

Dec. 31st.—She complained of pain at the scrobiculus cordis; a mustard poultice was applied, and on the following day a blister; the irritability of the stomach lessened; the evacuations were healthy.

Jan. 4th.—She became much worse; the countenance haggard, and she was

apparently prostrate; the gums were slightly ulcerated, as if from mercury. Chlorate of potash and bismuth were then prescribed.

Jan. 8th.—She had continued for the last three days in a dying condition, resembling a person with cholera, the eyes sunken, the countenance sallow, the voice scarcely audible, the hands out of bed, and almost without power of motion; the pulse was exceedingly small; the abdomen was collapsed, and free from pain and tenderness, the respiration was easy, but accompanied with a slight catch, and pain appeared to be felt at the base of the left lung, so as to give the idea that she might have some pleuro-pneumonia; there were sordes on the teeth, and the tongue could scarcely be moved. She was sensible, but appeared exceedingly prostrate and dying. She died the following morning.

Inspection.—Eighteen hours after death.—The head was not examined. The mouth and pharynx were healthy; but the lower half of the œsophagus presented irregular ulceration, in longitudinal lines, or rather a series of small, circular ulcers; there was slight congestion of the membrane.—*Abdomen.* The intestines were collapsed, and the peritoneum had partially lost its transparency. The stomach was very much contracted, and the pyloric portion only observable. It was found, that the omentum was adherent to the pylorus, so also the colon. On removing the whole of the stomach, with the spleen and the diaphragm, the small size of the stomach was still more manifest; it was only five and a half inches in length, from the œsophageal opening to the pylorus, and an inch and a half in breadth. At the cardiac extremity, it was found to be firmly adherent to the spleen and to the diaphragm; and on opening the stomach close to the œsophageal orifice there was a pouch, about an inch in length, resembling the finger of a glove; the extremity of this pouch was perforated, and passed into a cavity situated between the spleen and the diaphragm, containing dirty mucous fluid. The sides of the pouch were smooth, and adherent to the spleen. The mucous membrane of the stomach was destroyed near the œsophageal opening; and at the pylorus also, for one inch, the mucous membrane was completely destroyed and injected, the ulcer being bounded by a defined line. At the centre of this pyloric ulcer all the outer coats of the stomach were destroyed, and extravasation only prevented by the adhesion of the omentum. In the intermediate portion, near the lesser curvature, in several places the mucous membrane appeared slightly raised, and in others it was smooth and firm, as if cicatrized; on carefully examining a section of this smooth portion with $\frac{1}{4}$ -object glass, immediately beneath the surface a considerable quantity of fibrous tissue was observed, and still lower were the remains of distended gastric follicles. It appeared probable that the mucous membrane had at this part been superficially injured, and that the fibrous tissue was the result. The greater curvature was slightly discoloured by gastric juice, but appeared otherwise uninjured. The muscular coat of the stomach was considerably hypertrophied, especially near the pylorus, where it was a quarter of an inch in thickness. The duodenum and intestines were healthy throughout; the colon contained some scybala. The spleen was small; the liver and kidneys, uterus and ovaries, were healthy; the bladder was dis-

tended; the left pleura was firmly adherent; the lower lobe of the lung was in a state of red hepatization; it was soft, heavier than water. The right pleura and lung were healthy. The cavities of the heart contained fibrin; the heart itself was healthy and tolerably firm; its weight was 8 oz.

The absence of pain in this case, although there was extensive injury to the stomach, was very remarkable. The symptoms were exceedingly slight, vomiting shortly after food being the only prominent effect of the poison; the patient was, however, depressed, and the pulse was very feeble; she then became perfectly prostrate, and her death appeared to result from asthenia. The pneumonia which took place on the left side was, perhaps, set up by the abscess on the opposite side of the diaphragm, or it was the result of the absorption of decomposing material into the blood.

The effect produced on the stomach a considerable time after nitric acid had been taken, is well shown in case No. IX, related with diseases of the œsophagus, in which a young person lingered for three months, and suffered at last from the ordinary symptoms of obstructed pylorus; that part had become thickened and contracted, so as to lead to fatal obstruction, and the stomach was enormously distended.

CASE XLII.—Poisoning by Corrosive Sublimate. Death on the sixth day.—William P—, æt. 25, was brought to Guy's Hospital on October 16th, 1861, about 7 a.m., having swallowed ʒij of corrosive sublimate, dissolved in vinegar, about one hour previously. Vomiting came on in half an hour, and became very severe, the stomach at once rejecting everything. The stomach-pump was used, and eggs were given freely. At 11 a.m., he had an anxious expression; there was no whiteness of the gums nor of the mouth; the tongue was clean; saliva was abundant, and the vomiting was frequent. He had been purged, and the evacuations were of a dysenteric character, but no urine had been passed; there was *no* pain at the stomach, nor in the abdomen, but he complained of pain at the back of the mouth and at the upper part of the gullet, and the pain was greatly increased by pressure on the cervical spine. The pulse was small, soft, feeble, and very compressible, 80; the vomited matters consisted of mucus in large quantities.

On the 17th he had passed a restless night, sleeping for a few minutes at a time; he had passed bloody mucus, and suffered from tenesmus; there was a total suppression of urine. He complained of a sense of coldness, the pulse was 70, and he had tenderness about the umbilicus, and in the evening had a good deal of pain in the abdomen, especially in the right side, over the descending colon. There was no vomiting, and the mouth was parched.

On the 18th the stools consisted of blood and mucus, but the urine was still

suppressed; the pulse was 72; he had had a restless night. *On the 19th* he seemed better; he had slept at intervals, and had passed urine. There was less tenderness of the abdomen, and the pulse as well as the countenance were more natural. *On the 20th* he had had a little sleep. Three or four motions were evacuated, but no urine; there was soreness of the throat and inability to swallow; the face and the neck were somewhat swollen, and there was salivation; there was also tenderness over the umbilicus.—*21st*. He had had a bad night; the bowels were moved several times, the stools were of a dysenteric character. He complained of the throat, which was swollen, and he expectorated a small quantity of blood. His strength was evidently failing, and he died the following day at 2 a.m.

Inspection.—The body was well developed; the muscles were very prominent from post-mortem rigidity. Bloody mucus flowed from the mouth; the brain was healthy. The *œsophagus* was affected throughout; a fine layer of lymph covered the mucous membrane, and at the lower part the longitudinal ridges were covered with adherent granular lymph. The *bronchial tubes* were filled with bloody and frothy mucus. The *lungs* were extremely congested, and were in a state of early inflammation; the posterior parts of the lower lobes were especially affected; they were red, solidified, and devoid of air, but some fluid could be squeezed from them; other portions were in an earlier stage of pneumonia, and in some parts blood was extravasated. The *pericardium* was healthy; the right side of the heart was distended with dark blood and soft coagulum, but the ordinary tough, white fibrin was not present. The left ventricle contained a small quantity of semifluid blood. *Abdomen*.—The peritoneum was healthy. The stomach was of a pinkish colour, but the colon and the rectum were the portions of the intestine which were especially affected. The duodenum was healthy, it was filled with bilious matter. In the jejunum here and there, were patches of congestion, especially affecting the *valvulæ conniventes*. In the ileum there were some dark, congested parts; and at the lower part the mucous membrane was swollen; here and there were patches of granular lymph, and in the mucous membrane there were numerous dark spots of extravasated blood, the distribution and size of which corresponded to those of the solitary glands. The *cæcum* itself had escaped, but where the ileum joined the colon at the upper limit of the *cæcum* the congestion was extreme. The *ascending colon* was the part most severely affected; the mucous membrane there was swollen and of a dark-purple colour, and patches of lymph were adherent, especially on the longitudinal bands. Granular lymph was closely adherent, and the mucous membrane was superficially ulcerated, but no extensive ulceration was observed. The transverse and descending colon were congested in parts; but the *rectum* was in a similar condition to the ascending colon, being of a dark-purple colour, and covered with lymph. The liver was apparently healthy, but full of blood; so also was the spleen. The kidneys were swollen, injected, rather soft and coarse; the uriniferous tubes were filled with secreting cells and with cells filled with granules; and some fat-globules were observed. The mucous membrane of the bladder was very vascular.

tended; the left pleura was firmly adherent; the lower lobe of the lung was in a state of red hepatization; it was soft, heavier than water. The right pleura and lung were healthy. The cavities of the heart contained fibrin; the heart itself was healthy and tolerably firm; its weight was 8 oz.

The absence of pain in this case, although there was extensive injury to the stomach, was very remarkable. The symptoms were exceedingly slight, vomiting shortly after food being the only prominent effect of the poison; the patient was, however, depressed, and the pulse was very feeble; she then became perfectly prostrate, and her death appeared to result from asthenia. The pneumonia which took place on the left side was, perhaps, set up by the abscess on the opposite side of the diaphragm, or it was the result of the absorption of decomposing material into the blood.

The effect produced on the stomach a considerable time after nitric acid had been taken, is well shown in case No. IX, related with diseases of the œsophagus, in which a young person lingered for three months, and suffered at last from the ordinary symptoms of obstructed pylorus; that part had become thickened and contracted, so as to lead to fatal obstruction, and the stomach was enormously distended.

CASE XLII.—Poisoning by Corrosive Sublimate. Death on the sixth day.—William P—, æt. 25, was brought to Guy's Hospital on October 16th, 1861, about 7 a.m., having swallowed ʒij of corrosive sublimate, dissolved in vinegar, about one hour previously. Vomiting came on in half an hour, and became very severe, the stomach at once rejecting everything. The stomach-pump was used, and eggs were given freely. At 11 a.m., he had an anxious expression; there was no whiteness of the gums nor of the mouth; the tongue was clean; saliva was abundant, and the vomiting was frequent. He had been purged, and the evacuations were of a dysenteric character, but no urine had been passed; there was no pain at the stomach, nor in the abdomen, but he complained of pain at the back of the mouth and at the upper part of the gullet, and the pain was greatly increased by pressure on the cervical spine. The pulse was small, soft, feeble, and very compressible, 80; the vomited matters consisted of mucus in large quantities.

On the 17th he had passed a restless night, sleeping for a few minutes at a time; he had passed bloody mucus, and suffered from tenesmus; there was a total suppression of urine. He complained of a sense of coldness, the pulse was 70, and he had tenderness about the umbilicus, and in the evening had a good deal of pain in the abdomen, especially in the right side, over the descending colon. There was no vomiting, and the mouth was parched.

On the 18th the stools consisted of blood and mucus, but the urine was still

which the constriction of the œsophagus was relieved by bougies. Great injury may be done by the too-active interference of the attendant; depletion and mercurials, as well as powerful stimulants, generally aggravate the mischief.

Subacute Inflammation.—Catarrh of the stomach takes place, probably, in an acute form, and is the cause of some of the varieties of dyspepsia; but we are not cognizant of the conditions observed after death indicative of this state. It very rarely happens that any can have the opportunities possessed by Dr. Beaumont, of observing the appearance of the gastric mucous membrane; he found sometimes an erythematous condition, with deficient gastric secretion, arising from irritating food or stimulants. These cases of dyspepsia may not be more functional in their character than coryza or slight conjunctivitis. In ordinary catarrh, after exposure to cold, we find there is generally partial, often complete, loss of appetite, and occasionally diarrhœa, and the mucous membrane of the stomach joins in the general condition.

In the dyspepsia just mentioned, as arising from irritating food or stimulants, the *symptoms* are nausea or vomiting increased by food, injection of the tongue, and enlargement of the papillæ, with sometimes an abundant yellowish-white fur, tenderness and uneasiness at the scrobiculus cordis, pain, extending to the back between the shoulder-blades, loss of appetite, languor, headache, and an incapacity or unwillingness for mental or physical exertion, an anxious, care-worn countenance, with a sunken condition of the eyes. Oftentimes there is slight febrile disturbance, with a burning sensation at the palms of the hands and soles of the feet; the bowels become irritable or constipated; the urine is high-coloured, and, with nitric acid, assumes a deep-red colour, or it deposits lithates abundantly. When vomiting is severe, bright-green fluid and mucus are often rejected. This state, in not a few instances, passes into a more chronic form of disease, which is often called *chronic gastritis*, a condition which is indicated by soreness at the scrobiculus cordis; by pain, which extends to the spine; by sallowness of the complexion, and an anxious, distressed countenance; by injection of the tongue, which is sometimes like raw beef, or has red patches upon it, as if deprived of epithelium; by a com-

pressible pulse, and by emaciation. The pain at the stomach is increased by food in every form. This disease is often very obstinate, and persists month after month; and although we have no evidence of actual ulceration and destruction of the surface of the mucous membrane, still there is probably chronic inflammatory change. The gums sometimes become spongy, and the mouth and pharynx aphthous and painful; vomiting also is occasionally a troublesome symptom, and diarrhoea may be present.

There is a predisposition to this form of disease in strumous subjects; but we must distinguish this affection from a sympathetic irritation of the stomach, produced in the early stage of disease of the lungs and of the brain, and to which we shall have subsequently to refer. And, especially, towards the close of phthisis, of cirrhosis, of strumous peritonitis, &c., a state arises which, although associated with a general condition of exhaustion, closely simulates what we have described as chronic gastritis; in these instances, however, the small and large intestines are often implicated, and we find a flush on one cheek, profuse perspirations at night, preceded by dry, burning heat; the bowels are at one time confined, at another relaxed, and the motions are very offensive in character. These symptoms are often associated with great irritability of temper, fretfulness, and sometimes with delirium. (See Gastro-enteritis.)

In the disease to which my late colleague, Dr. Addison, gave especial attention towards the close of his life, "melasma supra renale," irritability of the stomach and great prostration of strength are the two most prominent symptoms; and so closely do the symptoms resemble those produced by poisons, and so frequent are the traces of gastric irritation found after death, that it has often been questioned how far these gastric symptoms, with bronzed discoloration of the skin, are due to sympathetic disturbance, and how far to disease of an inflammatory character. An instance of this kind is recorded, with cases of superficial ulceration of the stomach; and the following, more recently under my care, is exceedingly interesting in this relationship.

CASE XLIII.—*Bronzed Discoloration of the Skin. Violent Vomiting. Excessive Prostration.*—George A—, æt. 23, was admitted, under my care, into

the clinical ward of Guy's Hospital, on April 10th, 1861. He had been a lighterman, and till twelve months previously had enjoyed good health; at that time he was working very hard, slept in imperfectly ventilated cabins, several times he got his clothes wet from falling into the river, and soon afterwards began to feel weak, and unable to do his work with his usual energy; still he did not suffer from any cough or pain. Three years previously his skin had been sallow, but this colour disappeared; for 5 months, however, it had again assumed the appearance presented on admission. Some time before he had had syphilis, and had also been out on the coast of Africa. Three weeks before admission he went down to Sydenham, with the hope of regaining strength; but after taking a dose of castor oil violent vomiting came on, and continued with rapidly increasing prostration. He was a spare young man, with a peculiar sallow, semi-jaundiced appearance of the face, and numerous dusky-red blotches were observed on the cheeks and forehead; the discoloration was very much less on other parts of the body. On admission he was almost pulseless, cold and faint; he had had violent vomiting; the eyes were sunken, the breath sweet, the abdomen was contracted; and there was no evidence of disease of the chest; the tongue was clean; the motions were dark, the urine high-coloured. He was ordered some brandy at once, and effervescing mixture. On the 17th he was rather more comfortable, but the vomiting continued; food was *at once* rejected, and also a considerable quantity of green fluid and mucus; the mind was clear, the bowels open on the previous day. Ordered to take soda-water with brandy or with milk; to continue effervescing mixture, to which gr. ij of iodide of potassium were added, and at night, in the form of pill, calomel gr. j with extract of henbane and compound rhubarb pill, of each two grains. On the 20th the vomiting continued, and he was more prostrate; the tongue was injected and dry; the pulse slow, very compressible, and irregular; he seemed sleepy; was entirely free from pain; the abdomen was contracted, the urine was pale. He requested to have sherry instead of brandy; effervescing mixture with ammonia was given, but he died on the following morning. *On inspection* one or two lobules at the apex of the right lung were found to be white and consolidated, they were partially gray; the other thoracic viscera were healthy. The *stomach* had a considerable quantity of tenacious mucus covering its mucous membrane; there were numerous patches of arborescent injection of the capillaries, and some points of ecchymosis; this injection was most marked at the edges of the rugæ and towards the greater curvature; there was no ulceration. The first part of the duodenum was injected, and Brunner's glands were distinct. The other portions of the intestines were healthy, so also the peritoneum and kidneys. The liver was slightly congested. The spleen presented throughout its structure minute white points (Malpighian corpuscles). The supra-renal capsules were much enlarged on both sides, and contained white deposit.

As to *remedial* measures, stimulants aggravate the malady; but leeches may sometimes be applied in severe instances to the *scrobiculus cordis*; whilst, generally, cool drinks, soda-water,

or ice mitigate the symptoms, and may be used with solution of potash and demulcents, or with magnesia and opium. If the bowels are confined, saline aperients should be given, as the carbonate of magnesia and soda with lemon juice or citric acid. Mercurial purgatives are sometimes used, and have the advantage of thoroughly unloading the bowels and of diminishing capillary and portal engorgement, without increasing the irritability of the stomach; thus, calomel may be given in doses of three to five grains in combination with colocynth, aloes, rhubarb, henbane, &c.; or blue pill or gray powder may be administered in similar combination; or, again, the calomel is sometimes decomposed by alkaline carbonates, as in the preparation of soda with mercury in the Guy's Pharmacopœia. It is well also to follow these mercurial purgatives with a warm aperient draught, so as to produce a full and copious evacuation. To give mercurial preparations, so as to affect the system generally, we think to be positively injurious, and it is better when they can be altogether dispensed with; saline aperients, followed by vegetable tonics, and a bland, unstimulating diet, are all that are necessary. Subsequently, if the appetite fail, and the mucous membrane be relaxed, benefit accrues from the internal administration of the dilute nitric or nitro-hydrochloric acids, with vegetable infusions, as calumba, cascarilla, cusparia, or gentian.

In the more chronic forms the means best calculated to afford relief are nutriment of a character which is easily digestible, given in small quantity, and well masticated. Alkalies and salines relieve the irritability and congested state of the mucous membrane; for instance, solution of potash, the bicarbonate of potash or soda, calcined or carbonate of magnesia, administered with almond emulsion, with camphor mixture, or with any demulcent. If there be neuralgic pain, it is well to add a few drops of dilute hydrocyanic acid, tincture of henbane or conium, the solution of morphia or a preparation of opium; or to give the trisnitrate of bismuth, with chloric ether.

In some cases I have seen very great benefit from the administration of lemon juice; the pain has subsided, and the toleration of food and ability to digest it have considerably increased. It must be borne in mind, in these cases, that whilst vegetable food appears to be less easy of digestion, and often has to be

prohibited, if months are allowed to pass without its use the health fails on that account alone, and increased cachexia is produced; the administration of fruit, oranges, grapes, &c., is advisable; the juice of a lemon may be taken daily with relief to pain and distress at the *scrobiculus cordis*.

The application of leeches or blisters is often of service, and in some cases I have known benefit derived from the introduction of a seton.

The action of the skin should be promoted; and although sudden and violent exercise is injurious and could not be borne, still, exercise in the open air, and even horse exercise, is often very beneficial. A change of air should, if possible, be made, especially from a damp and relaxing situation to one of a more dry and bracing character.

Chronic Catarrh of the stomach and of the intestines, although arising from inflammation affecting them in common with the pulmonary mucous membrane, is more frequently observed as a consequence of congestive disease of the portal system. The mucous membrane becomes congested, often intensely so, or even ecchymosed; it has a swollen, oedematous or granular appearance, and is covered with a thick and tenacious layer of mucus. This is sometimes found to be alkaline in its reaction, is with difficulty washed off by water, and consists of mucous corpuscles, nuclei and epithelium.

Thickening of the mucous membrane, and gray discoloration from the deposition of pigmental granules from the long-continued congestion of the capillaries, are the result of chronic catarrh. The follicles of the stomach are found very distinct, and filled with nuclei and cells.

The cause of this condition appears to be long-continued congestion. In chronic bronchitis and emphysema, in valvular or obstructive disease of the heart, in cirrhosis of the liver, and in other conditions, the *vena portæ*, with its tributary branches, becomes over-filled with blood, and consequently the capillary vessels from the mucous membranes of the viscera also become surcharged; altered secretion, and the condition we have just described, is the consequence; thus, the congestion is not limited to the stomach, but extends through the whole of

the tract of the alimentary canal, in both the small and large intestines.

These conditions, then, are not in themselves primary, but are the result of cardiac, pulmonary, or hepatic disease. Sooner or later, in most cases, the signs indicative of gastric catarrh come on, pain at the scrobiculus cordis increased by food, pain between the shoulders, occasional vomiting, flatulence, oppression at the stomach, malaise, constipation; the flatulent distension after food becomes exceedingly distressing, so that scarcely any can be taken with comfort, and solid food is almost discarded; an attack of hæmatemesis, or of bleeding from hæmorrhoids, &c., may remove the congestion, and afford comfort to the patient, but the symptoms are very quickly reproduced. The dyspnœa and palpitation of heart disease, the cough and gasping for breath of chronic bronchitis, engage the attention of the patient, and obscure the less urgent symptoms of disease of the alimentary canal; it is when the former have been relieved that attention is directed to the abdomen. This state of catarrh is often relieved by the same means which mitigate the original disease. Emptying the portal system not only diminishes the distension of the right side of the heart and of the pulmonary vessels, but also the congestion, which is the direct cause of the catarrh of the intestines. Purgatives, saline, hydragogue, or mercurial, are generally used; and sometimes the more direct means of relieving the vessels, by the application of leeches to the anus. The administration of mineral acids, with demulcents, expectorants or tonics, according to the condition of the patient, affords great relief; so also the administration of steel.

The following are some of the cases in which we have found marked Catarrh of the Stomach :

Elizabeth E—, æt. 46. Catarrh of stomach, with bronchitis.

George P—, æt. —. Catarrh of stomach, with pneumonia.

Mary H—, æt. 25. Catarrh of stomach, with contracted mitral.

James M—, æt. 22. Catarrh of stomach, cæcum and colon, with diseased aortic valves and degenerated kidneys.

George C—, æt. 55. Catarrh of stomach, with diseased aortic valves.

Joseph S—, æt. 41. Catarrh of stomach, with superficial ulceration; diseased heart and kidneys.

Ann A—, æt. 23. Catarrh of stomach, with superficial ulceration; cystic disease of the ovary.

Diphtheritic Inflammation of the Stomach.—Acute inflammation of mucous membranes manifests itself by alteration in the secretion and condition of all the parts composing them. The capillaries and the blood within them, the formation of epithelium or mucus or of other secretions, are modified, and the whole vital condition of the part deviates from the healthy state. "The more the conditions of nutrition deviate from what is normal, the more will the material effused from the vessels deviate from the normal type."* This is exemplified in ordinary catarrh and bronchitis, as compared with the effusion of false membrane in croup and laryngitis. In these diseases the membrane becomes intensely red from congestion of its capillaries, swollen from effusion of serum into its tissue, hot and more highly sensitive, and its secretion changed. If the disease be slight, the mucus is altered in quantity rather than in quality, or its cells are found to be exceedingly abundant, and imperfect in their formation, or mere nuclei are produced. In croupous inflammation, the secretion consists of a blastema, with greater or less tendency to fibrillate, containing granules, nuclei, or variously formed cells. It is more or less adherent to the membrane beneath, though not incorporated with it. The larynx and trachea are most frequently the subject of this disease, or perhaps still more, the mucous membrane of the mouth, pharynx, and nasal passages.

The term diphtheritic inflammation was applied by Bretonneau to a form of acute inflammation of the mouth and pharynx, accompanied with the effusion of a grayish false membrane in small lenticular or diffused patches, and followed by superficial or deeper ulceration, and the disease extended to the nasal mucous membrane; the same term is applied to similar disease affecting other organs.

The stomach is less prone to acute inflammatory disease than either the small or large intestine, and we rarely have an opportunity of observing acute gastritis except as the result of irritant poisons. Croupous or diphtheritic inflammation is still more rare, and the following case, although in many respects imperfect, is of considerable interest; the symptoms of disease of the stomach were not clearly marked, but the patient was

* Paget, 'Surgical Path.' vol. i.

exhausted, and suffering from advanced syphilitic necrosis of the bones of the nose, and was also the subject of disease of the kidneys.

CASE XLIV.—*Syphilis. Diphtheritic Inflammation of the Stomach. Diseased Kidneys. Necrosis of the Bones of the Nose.*—Ann O—, æt. 47, was admitted, under Mr. Poland's care, Nov. 22nd, 1854, and died March 30th. She had had syphilis many years previously, for which she had taken mercury; and was admitted in a state of general cachexia, with necrosis of the bones of the nose. In this condition she continued till a short time before death, when she appeared more exhausted, and puffiness of the hands and face came on. She appeared to die from exhaustion.

Inspection fourteen hours after death.—The whole of the soft parts and bones of the nose as well as the palate were destroyed. In the brain there was serous effusion. The lungs and heart were healthy. The liver was fatty and nodulated, and it contained small lardaceous masses. The spleen was firm and waxy, and it also contained lardaceous masses; its weight was six ounces. The kidneys were much degenerated, presenting white deposit in the secreting structure, and the tubes contained highly refracting granules (fat).

The stomach presented a very remarkable appearance; it was of normal size. The mucous membrane was intensely congested; in numerous parts were small patches of thin, yellowish, lymph-like substance, which were very adherent, and were composed of mucous cells, granules, granule-cells, and some secreting cells. Other parts of the mucous membrane were covered with tenacious mucus. There was intense congestion of the capillaries of the mucous membrane, the follicles of which were distended with granules and with secreting cells. A dissolute life, and the impairment of general health by syphilis and mercury, were the predisposing causes of this disease of the stomach.

Suppuration in the Coats of the Stomach.—Local suppuration in the walls of the stomach is of exceedingly rare occurrence. The history of the following case is imperfect in its details, but is sufficient to show the general character and symptoms of such disease. It is probable that the case was one of pyæmia.

CASE XLV.—Elizabeth T—, æt. 40, was admitted, May 2nd, 1847, into Guy's Hospital. She was a married woman, and a nurse. For a fortnight she had suffered from pain in the limbs and back, and for a few days in the

stomach and chest. The abdominal tenderness subsequently increased. She had anorexia, and constant vomiting of a dark-coloured, bitter fluid, with intense thirst. Her death was preceded by restlessness and stupor.

Inspection twenty-four hours after death.—The body was tolerably nourished. The peritoneal cavity contained yellow, opaque, puriform secretion of uniform consistence, but of very offensive odour. At the pyloric third of the greater curvature of the stomach was a firm mass, measuring four and a half by three and a half inches. On opening the stomach, a small quantity of greenish fluid escaped. The mucous membrane was dotted over its surface with points of ecchymosis, and an irregular, dark-brown patch, about the size of a shilling, was found near the pylorus, at the centre of the thickened mass. When the peritoneal and muscular coats were divided, there was found to be a collection of pus in the submucous cellular tissue. The pus was semifluid. The intestines were distended with gas; but no disease could be found in the mucous membrane, except a small polypus in the rectum. The liver was dark, congested, and lacerable. The spleen and kidneys were congested. The uterus was full of menstrual blood. (Preparation 1802nd.)

Superficial Ulceration of the Stomach.—There are several forms of simple ulceration observed in the stomach:—1st. Superficial ulceration, affecting only the mucous membrane, which, although confined to the surface, is associated with marked gastric symptoms. 2ndly. Follicular ulceration. And 3rdly. Perforating and chronic ulceration.

Long-continued congestion of the mucous membrane of the stomach not only produces the state which we have described as chronic catarrh, but is also followed by superficial ulceration; but, this destruction of the mucous membrane is also the result of subacute inflammation. The membrane is generally found to be congested, especially at the rugæ; and the ulcers are situated at the lesser curvature, or in the neighbourhood of the pylorus. The ordinary size of these ulcers is about a quarter of an inch in diameter; they have irregular and sometimes rounded edges; as to colour, they are minutely injected or pale; and in depth they often reach to the submucous cellular tissue; a single point of ulceration may alone be observed, or several parts of the membrane be thus superficially destroyed. The intervening tissue may either have a normal appearance, or present arborescent and even general congestion, whilst the submucous and muscular coats of the stomach remain free from hypertrophic change, unless chronic irritation has existed. On further

and microscopical examination, the ulcerative process is found to have irregularly destroyed the gastric follicles laterally; and the surface presents mucus-cells, nuclei, and epithelium.

The *symptoms* of superficial ulceration sometimes have an acute character; vomiting, pain at the scrobiculus cordis, and between the shoulders, tenderness at the epigastrium, pyrosis, injection of the tongue at the tip and edges, loss of strength, and even great prostration, are the symptoms which have been observed in this disease; but in those instances which we shall presently mention, other coincident diseases existed; in one, disease of the supra-renal capsule, with long continued indigestion; in a second, the patient was affected with phthisis, after great intemperance, and he rapidly sank, apparently from exhaustion; in a third case, there was very severe chorea. The great prostration of strength was a marked symptom, and a most interesting one, when it is viewed in connexion with the intimate union of the stomach with the large plexuses and ganglia of the sympathetic nerve. The association of some of these cases of superficial ulceration with pyæmia, appears to show that a general diseased condition of the blood predisposes or excites this change.

In the class of cases associated with portal congestion, vomiting of coffee-ground substance sometimes takes place before death, and fluid of a similar kind is found in the stomach after post-mortem examination; chronic congestion followed by ulceration leads to effusion of blood, which gives rise to this red-coloured vomited fluid. The cases which I have very briefly referred to, were connected with renal anasarca and diseased heart, or with cirrhosis; but obstructive disease of the heart and lungs, and any condition which interferes with the free circulation of blood through the liver, predisposes to this form of disease. This state of stomach is often associated with chronic catarrh, and the symptoms before mentioned are generally present; an attack of hæmatemesis may afford great relief, but the symptoms may be so masked by the primary disease that this gastric ulceration is only recognised on the post-mortem table.

In the *treatment* of superficial ulceration the application of leeches, or of a small blister to the scrobiculus cordis affords

considerable relief; nitrate of bismuth with conium, or with morphia, and hydrocyanic acid, soothes the irritated membrane, and diminishes pain. Solution of potash, the bicarbonate of potash, or of soda, with anodynes and demulcents, render the mucus less irritating, and they diminish the congestion of the mucous membrane. Nitrate or oxide of silver in small doses relieves the pain, and renders the stomach more tolerant of food. Chloric ether also, with carbonate of soda, and mucilage mixture or almond emulsion is often of great service.

Ice and cold water are exceedingly grateful to the patient, but stimulants are not well borne, although on account of the prostration we are often tempted to give them; if they be absolutely called for, they should be diluted and mixed with food, as wine with arrow-root or jelly, or a small quantity of brandy with soda water. A farinaceous diet is more suited to these cases than animal food, which taxes in a greater degree the energies of the diseased membrane; it is well to allow three to four hours to intervene between each meal, unless the stomach be very irritable, or the patient prostrate, when small and often repeated quantities are to be preferred.

In some cases of anæmia, and chlorosis with leucorrhœa, steel may be taken with advantage; it should be administered in the milder forms, as the ammonio-citrate and tartrate, or in pills, as the compound steel pill with henbane and rhubarb, and always after a meal.

In the congestive forms of ulceration, purgatives relieve portal congestion, and thus remove much distress; but their action is followed by prostration, so that at last we are obliged to suspend them altogether. Diuretics and diaphoretics also tend to a similar result; small depletions afford temporary relief, but are not called for unless the respiration and the impeded action of the heart absolutely require them.

CASE XLVI.—*Superficial Ulceration of the Stomach. Diseased Supra-renal Capsules.*—John J.—, æt. 22, was admitted March 20th, and died on the following day. He was a stonemason by trade, residing at Lambeth, and during the winter had had pain in his stomach and vomiting. He slightly improved, but the day after Christmas was confined to his bed from great pain at the stomach and vomiting. The vomited matters consisted of watery fluid. At that time he had *tic-douloureux*. On admission the extremities were cold, he was almost pulseless, his hands were blue, he had not any

diarrhoea, but he had slight pain in the hypogastric region; he was sensible; the pupils were much dilated. He rallied a little after admission, but vomiting of bilious matter came on, and he appeared to die from syncope.

The *inspection* was made seventeen hours after death. The body was tolerably nourished, but the face was of a dingy hue, "*Melasma Addisonii*." The brain and its membranes were normal, but there was slight subarachnoid effusion. The trachea and bronchi were granular; at the apices of the lungs were lobules of iron-gray consolidated lung, with some calcareous deposit. The right side of the heart was moderately distended; the left was firmly contracted. On carefully examining the stomach, the cardiac extremity presented post-mortem solution, but towards the lesser curvature the mucous membrane was granular, and in several parts was destroyed by small patches of ulceration. These were quite superficial and irregular. In other parts above the line of solution there was arborescent injection. On microscopical examination, mucus and granule cells were observed. In the small intestine, Brunner's glands in the duodenum, and Peyer's and the solitary glands in the ileum, were very distinct. The liver and spleen were healthy; the kidneys coarse. The supra-renal capsules were atrophied, being only 49 grains in weight, and each was adherent to the surrounding parts by dense fibrous tissue; the left appeared irregular from contraction. The section was pale, red, and homogeneous, and presented fibrous tissue, fat, and cells about the size of the white corpuscles of blood.

There were evident symptoms of disease of the stomach in the pyrosis, pain, and vomiting from which this man suffered. His skin was discoloured, and the prostration of strength, which was very remarkable, is a condition which Dr. Addison draws attention to in his cases of disease of the supra-renal capsule. The case closely resembled one which we have previously detailed, No. XLIII; but in Case XLVI the symptoms of gastric disturbance were of longer duration.

The connexion of all these symptoms may be accounted for by the fact that the pneumogastric nerve not only supplies the stomach, and joins the large sympathetic ganglia of the solar plexus, but sends a large branch to join the sympathetic nerve of the kidney and supra-renal capsule, and this nerve is of considerable size. The exhaustion, collapse, fluttering pulse, present in many diseases of the abdomen, and sometimes produced by blows on the epigastrium, as well as the neuralgic pain in the side, with gastric irritation or ulceration, arise, no doubt, from this cause, namely, the connexion of the sympathetic with the pneumogastric and spinal nerves.

CASE XLVII.—*Superficial Ulceration of the Stomach and Duodenum. Phthisis. Ulceration of the Ileum, Cæcum, Colon, and Rectum. Great intemperance.*—James M—, æt. 64, was admitted into Guy's Hospital, April, 1854, in an anæmic and exhausted state. For a long period he had been exceedingly intemperate in his habits, and had become dissolute and helpless. He suffered from a slight cough, from palpitation of the heart, and, before admission, he had diarrhoea. He gradually sank. The skin was of a dingy colour. There were several vomices at the apices of the lungs; the pleura was semi-cartilaginous, and there was ulceration of the larynx, of the ileum, cæcum, colon, and rectum. Near the pylorus there was a small ulcer in the stomach, round, and about the size of a fourpenny piece, excavated, but without any external thickening. In the duodenum, near the pylorus, a similar ulcer was observed, but rather larger. The liver was pale and fatty; the kidneys were atrophied, and their tunic adherent; the heart was flabby; but, besides atheromatous deposit on the aortic and mitral valves, there was no disease. Preparation 1802⁷⁹. The state of the supra-renal capsules was not noticed.

The ulceration of the stomach and duodenum, in this case, was set up probably by the dissolute habits of the patient, and the symptoms of phthisis were obscured. The prostration, which was of a remarkable kind, was different from that of ordinary phthisis.

CASE XLVIII.—*Chorea. Vegetations on the Mitral. Ulceration of the Stomach.*—Elizabeth C—, æt. 18, was admitted March 28th, into Guy's Hospital. For two weeks before death she had very severe chorea, with constant jactitation, and sleeplessness; gradual exhaustion supervened. The mucous membrane of the stomach was softened and partially dissolved at the greater curvature. Near the lesser curvature were several small congested patches, in the centre of which the mucous membrane was destroyed. One of these had the appearance of a cicatrix.

On microscopical examination, the follicles were found to be full of granules, and cells containing highly refracting particles, somewhat resembling inflammatory cells. Similar cells, with mucus, were present on the surface, and the capillaries of the mucous membrane were much congested. The mucous membrane of the small intestine was similarly congested. The condition of the stomach appeared thus to indicate considerable irritation; but the severe nervous symptoms completely masked every other morbid state.

Numerous instances of superficial ulceration in connexion

with disease of the heart and liver, &c., might be quoted; but the following will suffice.

CASE XLIX.—*Chronic Catarrh and Superficial Ulceration of the Stomach, with intense Congestion. Disease of the Heart. Small Degenerated Kidneys.*—Joseph S—, æt. 41, was admitted into Guy's Hospital December 13th, 1854, and died January 13th, 1855. He was a large stout man, by trade a gas-pipe layer. When 15 years of age he had an attack of rheumatism, and again when he was 37 years of age; at the latter period he passed bloody urine. Two and a half years before his death he was again seized with rheumatism. He continued at his work till five months before admission, when his dyspnoea increased, and dropsy came on. The dyspnoea was urgent, and the dropsy was general; the urine was scanty and albuminous; and there was a bruit below the nipple and in the course of the aorta; the pulse was quick and irregular. On inspection, there was great congestion of all the viscera. The aortic valves were diseased, and the kidneys degenerated.

The *stomach* was intensely congested; its whole membrane, and especially the rugæ, were reddened. It was covered with patches of thick acid mucus, and at the cardiac extremity of the lesser curvature was a small ulcer, about $\frac{1}{2}$ of an inch in diameter; its edges were raised and intensely congested. Near the pylorus was a rather smaller, but similar ulcer. There were several points of actual ecchymosis. In the commencement of the duodenum were numerous minute points of ulceration, and the whole membrane was very much congested.

This case is imperfect in many of its details, but is especially mentioned as exemplifying the state of the stomach in obstructive disease of the heart. On examining the mucous membrane of the stomach, the follicles were found in their normal condition, but the capillaries were exceedingly distended; they were seen extending in a straight line between the follicles, and formed a close network immediately beneath the surface, apparently quite free from all cellular tissue.

CASE L.—*Catarrh, and Superficial Ulceration of the Stomach. Cystic Disease of the Ovary.*—Ann A—, æt. 23, was admitted October, 1854. She was a married woman, and with the exception of ague several years previously, she had enjoyed good health, till attacks of vomiting came on. Nine months before admission, after one of these attacks, she experienced pain in the side, and the abdomen became swollen. The enlargement increased for five months, and she was then tapped. One and a half pints of viscid fluid were thus evacuated three weeks before admission. The swelling reappeared and rapidly increased. On admission she was very ill. There was dyspnoea, with pain in the back and right leg. The bowels were constipated; the abdomen was large, and at the left side was a fluctuating

tumour, but on the right it was solid. Vomiting came on, and she gradually sank.

Inspection fifty-eight hours after death.—The peritoneum contained three to four quarts of bloody gray fluid, and also a large cystiform tumour formed by the right ovary. The stomach was large; its rugæ were reddened, and covered with a thick layer of mucus. The mucous membrane was thin, and presented, especially at the lesser curvature, numerous minute ulcers, which were found to extend to the submucous cellular tissue, and on their surface numerous spherical cells, containing highly refracting particles, were observed. The other parts of the intestine were much congested. The liver was fatty.

In this case the power of the patient was much reduced, and the abdominal tumour had exerted considerable pressure on the vessels. It appeared, however, that for some time before death, the mucous membrane of the stomach had been in an irritated, if not inflamed condition, as indicated by the repeated attacks of vomiting, before any mechanical pressure was exerted upon the viscus. The earlier attacks of vomiting were perhaps due to sympathy with the diseased ovary.

Follicular Ulceration of the Stomach.—Minute points of ulceration, varying in size from one sixteenth to one fourth of an inch in diameter, are sometimes found studding the whole surface of the stomach. They do not extend deeper than the mucous membrane, and are situated, not only at the lesser curvature, but over the greater part of the stomach; and they appear sufficiently distinct from the more common superficial ulcer to warrant separate mention. Dr. Brinton thinks that there is no proof of their origin in the gastric follicles, or in the lenticular glands when they are present; and suggests the term punctate ulceration as being more correct.

This form of ulceration has been observed in children with severe gastric symptoms; but it has been generally found after death when no indication of disease of the stomach had previously existed, excepting, perhaps, the vomiting of coffee-ground substance. A drawing, in the Museum of Guy's,* from an infant under the care of Dr. Lever, shows the mucous membrane of the stomach intensely congested, and covered with minute points of ulceration. The microscopical appearance of one of these minute ulcers presented irregular edges ex-

* Drawing, No. 2867^a.

tending into the gastric follicles ; its base consisted of the sub-mucous tissue, and on its surface were numerous cells, presenting changed secreting cells, and inflammatory granule cells. There was no proof that the disease originated in the solitary glands, but rather that it was follicular in its character. These ulcers, in some cases, are probably formed a short time before death ; and are due in part to irritating secretions, and to the depressed state of the nervous system. The disease is closely allied to the gastritis folliculosa of Cruveilhier, or to what is called hæmorrhagic erosion ; but the latter term is more applicable to some of the forms of superficial ulceration with great congestion.

In some fatal cases of hæmorrhage from the stomach, a minute ulcer, scarcely larger than those just described, has been found ; at the base of which the branch of an artery has been observed containing a small clot.* Sometimes there are seen numerous minute specks, each containing a small clot.

CASE LI.—*Follicular Ulceration of the Mucous Membrane of the Stomach, with Renal Anasarca and Diseased Heart.*—Susan K—, æt. 67, was admitted into Guy's, in June, 1854. She had general anasarca, with albuminous urine ; the pulse was irregular ; and there was dyspnoea, with palpitation of the heart. A short time before death, vomiting of a dark-coloured fluid took place.

On inspection, coarse congested kidneys were found, with a heart weighing 15 ounces, dilated and flaccid, and with some atheromatous deposit on the mitral and aortic valves. There were several small fibrous tumours beneath the peritoneum covering the uterus. In the stomach, above the line of gastric solution, were numerous minute ulcerations, about the size of a pin's head, studding over the whole of the membrane, and without any thickening of the submucous or muscular tissue. See Preparation, Museum, No. 1802⁷⁴.

CASE LII.—*Follicular Inflammation of the Stomach. Burn on the Leg. Amputation. Abscess in the Lungs and Spleen.*—George H—, æt. 15, was admitted into Guy's Hospital April 20th, and died June 23rd. He had scalded the arm and leg with hot tar. The left leg was principally injured, but was never disposed to heal, and the nerves became exposed ; the leg was amputated on account of his prostrate condition and the severity of the pain from which he suffered.

Inspection eight hours after death. The stump was sloughing, and the dry bone projected. The left arm was œdematous. There was lobular pneumonia of the lungs, with superficial pleurisy ; and the centre of the lobules were sloughing. In the substance of the heart were small white spots resembling pus. The upper surface of the spleen was softened, and

* See Preparation 1801³⁰, Museum.

presented suppurating points ; but the other parts were hard and filled with blood. The diaphragm on both sides was covered with purulent lymph.

The *stomach* contained coffee-ground fluid. Near the cardiac extremity were numerous minute follicular ulcers ; but gastric follicles were found in a normal state.

Chronic and Perforating Ulceration of the Stomach.—The form of ulceration which we have next to consider has been designated emphatically the ulcer of the stomach, by some also the chronic, and by others the perforating ulcer. Some cases of the latter description are not of a chronic character, and ought perhaps on that account to be considered apart ; many of those, however, which have been extended over considerable periods, terminate in perforation, so that we can scarcely separate the one from the other. If, however, the term perforating be meant to imply merely extension through the mucous, and also through the muscular and peritoneal coats, although adhesions prevent sudden fatal peritonitis, there is less objection to the term.

Chronic ulceration has probably in all cases been preceded by some of the conditions previously described. The ulcers are situated at the lesser curvature of the stomach, sometimes towards the anterior, but more frequently towards the posterior aspect, and near the pylorus ; they vary in size from a quarter of an inch to three inches or even more in diameter, and are round, oval, or reniform, the latter perhaps from the union of two ulcers. In the 'Dublin Hospital Gazette,' Dr. Law mentions an instance in which an ulcer was six inches in length in its long axis. Dr. Brinton,* in his investigations on 'Ulcer of the Stomach,' states that in 43 per cent. the ulcer was situated at the posterior surface, in 27 at the lesser curvature, in 16 at the pyloric extremity, in 6 at both the anterior and posterior surfaces, often at opposite places, in 5 at the anterior surface, in 2 at the greater curvature, and in 2 at the cardiac pouch ; those on the anterior surface being the most liable to lead to perforation and peritonitis.

The edges of the chronic ulcer are rounded and elevated ; but in those which are more emphatically termed the perforating ulcer the edges are frequently neither raised nor injected, but merely

* Brinton on 'Diseases of the Stomach.'

present a small punch-hole opening, perforating the peritoneal membrane. The opening is rather larger on the mucous surface, but there is no evidence of inflammatory action, and sometimes no adhesion whatever with adjoining viscera. More frequently, however, the edges of an ulcer in the stomach are considerably thickened by the infiltration of fibrous tissue in the mucous and submucous coat; the centre is therefore depressed, and a hollowed cavity is produced; and in many of these cases, also, the disease extends through the muscular and even through the peritoneal coat. The opening in the mucous membrane is larger than that of the muscular, and the muscular than the peritoneal, so that the ulcer has a bevelled appearance on its inner aspect. If the peritoneal ulcerate or slough before adhesions have formed, a round opening, as if a punch-hole had been made, is observed to extend into the serous sac, and to have led to rapidly fatal peritonitis. If, however, adhesions take place around the ulcer, its base is formed by the adjoining viscera, such as the pancreas, or the left lobe of the liver, or the spleen. In these cases the base of the ulcer, or cicatrix, is of a whitish colour, and consists of fibrinous effusion, and is smooth, or it has a minutely granular appearance; the edges become exceedingly firm, and are formed of dense fibrinous effusion into the mucous and submucous tissues. Glandular mucous membrane is not reformed in these cicatrices. The perforation into the peritoneum is sometimes found at the edge of a large ulcer which has been closed by adhesion, but has given way at the edge. Adhesions also take place between the anterior surface of the stomach and the abdominal parietes, so also with the parts in the lesser omentum, with the glands, &c. When the liver is invaded, we find that the adjoining hepatic tissue contains a considerable infiltration of white fibrous tissue. Ulceration occasionally leads to perforation of the coats of the adjoining vessels, either at an early stage, or when an ulcer has existed for some time. These hæmorrhages are sometimes rapidly fatal, or they become checked for a time, and often recur. Dr. Brinton describes three varieties of this hæmorrhage:—1st, The extension of ulceration into the minute vessels of the mucous membrane and submucous tissue, leading to a gradual discharge of blood, which becomes mixed with the secretions; 2nd, Greater hæmorrhage from sud-

den congestion of the ulcerated surface; and 3rdly, Very profuse bleeding from a large artery of the stomach. The perforated vessel is often seen closed by a small clot, or a drop of blood may be pressed from it, and in large ulcers may be sometimes seen like a small papillary eminence. This hæmorrhage, however, is not limited to the gastric arteries, but takes place from the arteries situated at the base of the ulcer, and belonging to adjoining viscera; thus, in one instance, Case LXVII, both the splenic artery and the pancreatic were perforated. (Preparation in the Museum of Guy's Hospital.) Dr. Lee, quoting Cruveilhier, states that ulceration into arteries and perforation take place more frequently in secondary ulceration of a cicatrix than with primary ulceration. Hæmorrhage, however, often occurs at an early period of the disease.

The form of the stomach may be greatly changed either by adhesions external to the viscus, or by contraction of the walls of the ulcer. When the ulcer is situated in the centre, the cavity may appear almost double, in the form of hour-glass contraction. It is exceedingly rare in simple ulceration, even when situated at the pyloric extremity, for the whole circumference of the part to be occupied by the ulcer and its contraction; the side is irregularly puckered rather than uniformly contracted. In cancerous disease it is more common to find one or other orifice surrounded. In Case LXXXIII the part which, on opening the stomach, was supposed to be the pylorus, was found to be a circular contraction and a large ulcer, about an inch and a half from the pylorus, and healthy mucous membrane intervened. But this was not simple ulceration; there was cancerous product in the contracted omentum at the part; it was doubtful whether the cicatrix of an ulcer had been followed by cancerous effusion in its neighbourhood, and it is probable that this is really the case in some instances. The thickening of the margins of the ulcer also encroaches upon the branches of the pneumogastric, and leads to intense pain, violent vomiting, and death from exhaustion. The second case related is of this character. The ulcer sometimes, also, extends into the sac of the lesser omentum, and forms there an abscess bounded by the spleen, diaphragm, pancreas, and liver; or it communicates with the colon, or even with the parietes; these, however, are generally

of a cancerous character;* or the diaphragm itself is perforated, and sudden acute pleurisy and empyema are produced, as in Case LXV.

A remarkable instance occurred in Guy's, in 1845, under Dr. Barlow's care, the full report of which, by Dr. Wilks, is found in the 'Medical Gazette' for May, 1845, but I have given a brief abstract of it here. A secondary cavity, partially filled with air, had given rise to the symptoms of pneumothorax. In another case which I have recorded, a sinuous ulcerated opening extended through the diaphragm into a sloughing cavity of the lung. A communication sometimes takes place from the colon, but this appears generally to extend from the intestine to the stomach, rather than from the latter to the former; and the ulceration in these cases is found more generally at the greater curvature. In a patient who died in 1847, Case LXIV, there was an ulcer opening from the colon into the greater curvature, and two others from the greater curvature into the sac of the lesser omentum, forming a large fæcal abscess, which extended through into the lung. Dr. H. Davies narrates a case in the 'Pathological Transactions,' of simple chronic ulcer extending into the colon. There had been dyspepsia and fæcal vomiting, whenever the bowels were constipated. The patient gradually sank.

Many facts of great interest have been brought forward in the valuable papers of Dr. Brinton. As to the sex,—that ulceration of the stomach is twice as frequent in females as in males; the reverse being the case in cancer. In 654 cases he found 440 were female, and 214 male; and that in one out of every five cases more than one ulcer was present; whilst in one out of every seven or eight cases perforation took place.

In the consideration of the age of those who are the subjects of this affection, the cases of cancer which have come under my own immediate notice have been more advanced in life than those who were the subjects of ulceration of the stomach. Dr. Brinton has collected a large number of cases, and shows that the ulcer generally "affects the periods of middle and advancing life with a frequency which gradually increases up to the extreme

* See Dr. Murchison, in 'Pathological Transactions,' vol. viii, &c.; and Dr. W. T. Gairdner, in 'Edin. Med. Journal,' 1855.

age allotted to man." But the cases of ulcer in which *perforation* happens "seem not only to select another period of life, but to exhibit a marked contrast of age in the different sexes, the period of life in which it is most liable to occur being quite a different epoch in the male and in the female,"—in the female being between the ages of 14 and 30, in the male from 50 to 60; the diminished risk of the female at the latter periods of life rendering the total risk in the same number of cases nearly equal.

The same author states that in the female one half the instances of perforation take place between the ages of 14 and 20, whilst in the male there is a constant proportion to 50, which diminishes but little to 70. The complete average age in the former, however, being 27 years, in the latter 42. Dr. Lees* mentions that he has seen perforation of the stomach from ulcer "in a girl of 8, and a boy of 9 years of age."

As to the situation of perforating ulcer of the stomach, Dr. Brinton gives the following as the relative situation in every 100 cases, but the term perforation here must be remembered as being applicable especially to those instances in which fatal peritonitis follows, not to those in which adhesion renders extravasation impossible, as we so often find in perforation on the posterior aspect.

Posterior surface	2
Pyloric sac	10
Middle	13
Lesser curvature	18
Anterior and posterior surface at once	28
Cardiac extremity	40
Anterior surface	85

The *symptoms* of chronic ulceration are at first those of ordinary dyspepsia, and are often very obscure, and imperfectly marked; thus slight uneasiness after food, and constipation, may be the only evidence of disease; afterwards the pain, with tenderness in the region of the stomach, especially at the *scrobiculus cordis*, attracts more attention from the patient; it is sometimes slight, at other times intense, and of a peculiar gnawing cha-

* Dr. Lees on 'Diseases of the Stomach and Indigestion.'

racter; the pain is generally increased by food, and relieved by the rejection of it; vomiting is, therefore, generally present, and pyrosis or water-brash. Other symptoms are, pain between the shoulders, more or less of abdominal uneasiness, constipation, emaciation, and a peculiar pallor and cachexia. Hæmorrhage and the rejection of blood by vomiting, hæmatemesis, or by the bowels, as black stools, melæna, are present in most cases at one or other stage of the disease.

The *pain* is not always of the same character, but may be considered as a symptom present in almost every case; it may be almost constant, but generally undergoes degrees of exacerbation, being increased by food; the patient often states that pain comes on as soon as the aliment reaches the stomach, and continues as long as it is retained; sometimes it is so intense that the patient is completely exhausted, as I have several times found when branches of the pneumogastric nerve have been involved in the dense edges of a chronic ulcer; at other times so slight, that it only amounts to tenderness on pressure at the scrobiculus cordis. In a case recently under my care, in which other signs of ulcer were present, the patient stated that the pain was sometimes relieved by firm pressure against the back of a chair. Position has, in not a few instances, a marked effect as to the severity of pain, and I have several times seen the observation of Dr. Osborne* confirmed, that the influence of the position of the patient on the pain serves as a guide to the seat of the ulcer, according as the contents of the stomach gravitate towards or away from the injured part; thus in an ulcer at the posterior part of the lesser curvature the patient has been most easy when leaning forward and towards the left side; on the contrary, I have seen the pain continue whatever position may be assumed. In young women suffering from well-marked ulceration of the stomach, with chlorosis, neuralgic pain in the side may be present at the same time as tenderness and pain at the scrobiculus cordis, and, in these, we have increase of pain during or prior to the menstrual periods. The pain in the back is rarely absent in chronic ulcer of the stomach; it is generally less severe, and comes on later than the gastric

* 'Dublin Journal of Medical Science,' 1st ser. vol. xxiv, p. 301.

pain, but is sometimes complained of more urgently than that at the stomach itself, the patient often stating that "the pain goes through to the back." In speaking of the diagnostic value of pain, Dr. Lees states, "the occurrence of pain will often be of great assistance in the diagnosis between simple ulcer and cancer; for it is an important and curious fact, that there is seldom pain in cancer of the stomach, unless great obstruction of the pyloric orifice prevents the passage of food out of this viscus." And again: "The mere fact of severe pain constantly occurring after food, should lead you to diagnose simple ulcer of the stomach rather than cancer. The pain, moreover, in simple ulcer, is often of a gnawing character, causing a sense of sickening depression; it is variable and remittent, sometimes being very severe and then ceasing for days or even weeks; but, in malignant disease, the pain, though not often severe or lancinating, yet is almost always constant after it has once commenced." Although all observers will bear testimony to the general variability and the gnawing character of the pain in ulcer of the stomach; still in some instances, as when branches of the pneumogastric nerve have been involved in the edges of the ulcer, or laid bare on its floor, the pain is terribly constant. Again, in cancer of the pylorus, as in Case LXXXIX, there was no pain acknowledged on repeated questioning of the patient as to present symptoms or previous history; and in other instances of cancer, when suffering is very severe, the relief from pain is often very great on the avoidance of solid and indigestible food, and during the use of anodyne remedies. The dragging pain in the epigastric region, produced by adherent omental hernia, and the pain often present at the scrobiculus cordis in chronic disease of the lungs and heart, must be remembered, as in Case LXIX, of chronic ulcer, in which the patient died from bronchitis and emphysema.

The period at which vomiting takes place in ulceration of the stomach is equally varied; sometimes the food is at once rejected, in other instances it is retained for many hours, or days. In the case previously alluded to, in which the thickened edge of the ulcer contained a large branch of the pneumogastric nerve, the stomach almost instantaneously rejected food, and the patient died exhausted. Fermentation and the development of the

sarcina ventriculi of Goodsir* take place in some cases of chronic ulcer, as well as in cancer and diseased pylorus. The sarcina can scarcely be considered as a proof of obstruction, for its development occurs without any impediment.

Vomiting, however, as a sign of gastric ulcer, must be regarded with great care, since in so many instances it is purely sympathetic in its origin. Pyrosis also is often present in disease of a less serious character, and is among the signs of functional disturbance.

Chronic disease of the abdominal viscera is often marked by an anxious and dejected countenance, with emaciation; in gastric ulcer this appearance is present, and is associated with pallor, arising from the condition of general nutrition.

In cancerous disease, a care-worn expression is found, with cachectic *sallowness*; and, in the anæmia of chlorosis and amenorrhœa, there is, in extreme cases, a waxen appearance, which is very peculiar; so also in the anæmia after considerable loss of blood. Again, in many instances of struma and of glandular disease, pallor is present; but in ulcer of the stomach, the anxious countenance of abdominal disease, conjoined with the emaciation and pallor of imperfect nutrition, afford a very characteristic morbid expression, and as a symptom is rarely absent.

As we have before said, hæmorrhage takes place in most cases of gastric ulcer; vomiting of several pints or even quarts of blood may be amongst the earliest symptoms of disease; in other instances the bleeding is slight, or entirely absent. The first hæmorrhage from the stomach is occasionally fatal; in ordinary cases the discharge of blood is preceded by a sense of weight and coldness, followed by faintness or actual syncope, and then rejection of dark-coloured blood takes place. The action of the gastric juice confers this deepened colour; but if the effusion be very rapid, the colour is more bright; a portion of blood passes onwards into the duodenum and intestines, and if life is prolonged, so that it may be discharged per rectum, a black, tarry evacuation is the result. Sometimes, however, the whole of the blood is thus discharged, and there is melæna without hæmatemesis; these two symptoms are generally combined. Instances have been recorded where sudden

* *Merismopodia ventriculi* (Robin).

hæmorrhage into the stomach was followed by fatal syncope, without the discharge of blood either by vomiting or purging; more frequently the hæmorrhage is oftentimes repeated.

Unless hæmorrhage, however, take place, we cannot with any certainty diagnose ulceration of the stomach; cachexia, emaciation, pallor, pain, and vomiting, all arise without ulceration, in cases of gastrodynia and irritability of the stomach, sympathetic or otherwise. But hæmorrhage is not in itself pathognomonic of ulceration; it often arises from over-distended capillaries in gorged portal circulation, and in cancerous disease: and although less frequent in cancer than in simple ulceration, it does occasionally arise. Disease of the œsophagus and aneurism sometimes produce the same symptom.

A form of ulceration, having much general as well as pathological interest, from its usual disastrous termination, is that class of cases which tends to fatal perforation into the peritoneal cavity. These instances often occur in young women affected with chlorosis and amenorrhœa, or with painful menstruation; the previous gastric symptoms are very slight, or altogether unnoticed, although there is generally impaired health, with leucorrhœa or chlorosis, neuralgic pain in the side, and symptoms of hysteria. The onset of the fatal attack is unexpected, and is generally after slight muscular exertion, or after a full meal; intense pain comes on, followed by rapid prostration and collapse. The skin becomes cold and clammy; the pulse fails; the pain in the abdomen becomes general; tympanitis follows; and occasionally vomiting supervenes; death ensues in from five to twenty-four hours, although life is sometimes prolonged for several days; and, in rare cases, the patient recovers.

Various suggestions or hypotheses have been made in reference to these cases of perforation.* The enfeebled nutrition, and inability of the coats of the stomach to resist the chemical action of the gastric juice is considered by some to be the cause of this terrible result; others refer it to the state of the nervous system; and we have ample proof of the close connexion of the gastric sympathetic nerve with the ovarian and uterine ganglia.

* Dr. Crisp and Dr. Brinton.

The pain below the mamma in leucorrhœa, arises probably from the connexion of the splanchnic with the dorsal nerves. The cause is equally obscure as to the part of the stomach usually chosen for ulceration. Why the lesser curvature, either at its posterior or anterior portion, should be so generally the part involved is not known. This is the least free in its movements, in fact almost stationary, the stomach in its general expansion and consequent movement turning upon its lesser curvature. This also is the region along which the pneumogastric nerve extends; but I am not aware that this portion of the stomach is more vascular than other parts.

Abercrombie distinguishes three modes of fatal termination of ulcer of the stomach. 1, Gradual exhaustion; 2, hæmorrhage; and 3, perforation into the peritoneal cavity. Another might also be mentioned: the production of inflammation by extension to adjoining viscera, as in the case related of extension through the diaphragm into the lung, and the production of acute pleurisy. The disease, however, sometimes remains in a passive condition, and the patient dies of some other complaint. It is not very rare to find cicatrices in the stomach; and in those cases where there has been extensive destruction of surface, and of the muscular and peritoneal tissue, the adjoining viscera are found covered with a smooth fibrous tissue.

The duration of life after the development of symptoms of ulcer of the stomach, as compared with cancer, is generally very different. Setting aside those cases in which perforation into the peritoneal sac takes place, the ulcer is more curable and extends over a longer period; it may be several years, and some have mentioned cases continuing even for twenty. I have several times observed patients in whom there were marked severe gastric symptoms: men of middle life, with sallow complexion, with pain at the scrobiculus cordis, vomiting of food, occasional hæmatemesis, loss of flesh, &c., who have lost their symptoms under treatment and care, have regained flesh and comfortable health, and had no return of symptoms for several years. In cancer, after the well-marked symptoms have occurred, we rarely find that a year passes, and frequently only three or four months, before a fatal termination takes place; and it is probable that many cases of supposed

cancer of the stomach, in which the patient survived for many years, were really chronic ulceration. It has yet to be shown whether the cicatrix of a chronic ulcer ever becomes the seat of cancerous deposition; but several instances warrant such a supposition.

CASE LIII.—Chronic Ulcer of the Stomach. Phthisis. Branches of the Pneumogastric Nerves involved.—Eliza B—, æt. 21, was admitted into Guy's Hospital October 13th, 1858. She had enjoyed good health till she was fifteen years of age, when her health failed. She was then obliged to leave her situation as nursery-maid, on account of weakness, and pain in the chest, &c. At sixteen menstruation commenced, and continued regularly till eighteen months before admission, but from that time there had been amenorrhœa. She was anæmic, emaciated, and had a dark areola around the eyes. The respiratory murmur was feeble at the apices; the heart sounds were normal. She complained of pain frequently recurring in the hypochondriac and lumbar regions; of a feeling of distension and pain after taking food, and of a sensation as of a weight rolling when she turned on her right side, the left being the side on which she could recline most easily. The appetite was bad; the bowels were regularly open; she slept but little; the urine was normal. Various remedies beside steel and cod-liver oil were given, and she left the hospital for a short time, but was readmitted on March 16th, 1859. She was then extremely blanched and much emaciated, and had a dejected countenance. She generally sat up in bed, inclining towards the left side, for it produced pain to turn on the right side. There was no evidence of disease in the chest found after careful examination; the respiration was 30; the chest was resonant; and the respiratory sounds were feeble, but otherwise healthy. She had vague pain in all parts of the body, especially in the shoulder; the spine was tender everywhere, and unusually prominent from the wasting of the muscles. The abdomen was rather distended, apparently from flatus. In the epigastric region there was a hard swelling, tender on pressure, and tympanitic. When in great pain at this distended part, if compression was made, the swelling disappeared, and the pain was relieved. On deeper pressure in the epigastric region, a hard, resisting sensation was communicated. The superficial abdominal veins were distinct. Opium, chloroform, chlorodyne, cod-liver oil, &c., were administered, with partial relief.

On May 5th she was transferred to my care, and was then greatly emaciated; she suffered constant severe pain in the epigastrium, and in the left shoulder; and the pain was worse after taking food, which also produced slight nausea. She had much flatulence, but no vomiting; the bowels were never acted upon without medicine or an enema. The heart's action was rapid; the rhythm regular; the pulse, 120, was compressible. The abdomen was firm and somewhat enlarged; the urine was normal. The pain increased in severity, and sometimes became of an agonizing character; it was somewhat relieved by medicines, as by opium, morphia, henbane, aconite, creasote, &c. The symptoms continued with varying severity till the end of July, but in August symptoms of phthisis began to develop themselves; the

right side became dull, and indubitable signs of disorganization at the apex were present. The patient began to cough and expectorate; became still thinner, so that she appeared to be a skeleton. Scarcely any nourishment was taken, but she lingered till October 10th.

On inspection, the right pleura was universally adherent; the left partially so. The right lung was extensively disorganized, and contained a large cavity in the upper lobe, and much tubercular deposit in the lower. The two lobes were adherent, and the cavity was situated at the lower part of the upper one. The apex of the lung was healthy and free from tubercle. The cavity was very large and ragged, and had apparently extended rapidly. The lower lobe contained numerous small tubercular masses. The left lung was nearly healthy; so also the heart. On opening the abdomen, the *stomach* was seen to be closely adherent to the left lobe of the liver at its under part. The stomach appeared to be thus hooked up in the central part, and presented an hour-glass contraction. This adhesion of the smaller curvature of the stomach and of the posterior surface of the liver was so firm that it was impossible to separate the two organs. The liver also contained much fibrous tissue. On attempting the removal of these viscera, it was found that the pancreas also was closely involved in the adhesions. By opening the stomach along its greater curvature, a large chronic ulcer was found to be situated at the middle of the lesser curvature. It was $2\frac{1}{2}$ inches in length, and 2 in breadth, and had completely healed, there being no sign of recent action about it. It had a raised, firm edge, and its floor was formed by tough, fibrous tissue upon the liver and pancreas, the coats of the stomach having been entirely destroyed at that part by the ulceration. The pneumogastric nerves on both sides terminated in the fibrous tissue which bounded the ulcer. The mucous membrane was elsewhere healthy.

The symptoms in this case advanced very slowly to a fatal termination; and nearly six years elapsed between the onset of the disease and the death of the patient. There was no hæmatemesis at any period, and the most prominent symptoms were anæmia, gradually increasing emaciation, and pain in the left side, which became exceedingly severe. The pain was at times agonizing, and was unrelieved by any form of anodyne medicine; the manner in which the branches of the pneumogastric nerve were involved in the dense fibrous tissue at the edges of the ulcer afforded an explanation of the suffering. The position of greatest ease, however, was that in which fluids gravitated from the ulcer, namely, the position assumed in leaning forwards and towards the left side; thus far confirming the statement of Dr. Osborne. Vomiting was not a prominent sign of disease. For some months the general symptoms were those of phthisis; and, in fact, towards the close, symptoms

of affection of the lungs were rapidly developed. In reference to the etiology of the phthisis, it is an interesting question whether irritation of the gastric branches of the pneumogastric nerve had anything to do with the production of disease at the termination of the pulmonary branches; or, whether the phthisical mischief was rather produced by the general impairment of nutrition.

CASE LIV.—*Chronic Ulceration of the Stomach, involving the Pneumogastric Nerve. Atrophy of the Left Lobe of the Liver. Death from Exhaustion.*—E. S.—, æt. 32, had been a widow for nine years. Four years before admission into Guy's Hospital she had an attack of hæmatemesis, but her health improved, and she continued in her situation as housemaid. Two years later she had a similar attack, and, six months later, severe pain at the region of the stomach came on. The pain continued for a few days, and was much relieved by vomiting of blood. A similar attack came on after admission. Great weakness, with severe pain in the stomach, were the principal symptoms; the pain extended to the back, and was of a paroxysmal character; and food was very quickly rejected. At the epigastrium, at first, fulness was felt, but afterwards, a distinct tumour. The vomiting was uncontrollable, but sometimes subsided for several days. Vomiting of grumous fluid then came on, and the pain at last became very severe; she gradually sank.

Inspection.—The body was much emaciated; there was no marked disease about the thoracic viscera; the stomach was distended, and reached nearly to the umbilicus. At the scrobiculus cordis the left lobe of the liver was contracted, and adherent to the lesser curvature of the stomach; and constituted the tumour which had been felt during life. The stomach contained a considerable quantity of grumous fluid, and at the lesser curvature, about half an inch from the pylorus, was an ulcer two inches and a half in diameter, with raised everted edges of mucous membrane, and dense fibrous tissue; the base of the ulcer was smooth, composed of an albuminous layer, and formed by the pancreas and by the inferior surface of the left lobe of the liver. The pylorus was not hypertrophied. The remaining part of the stomach appeared healthy. On carefully dissecting the pneumogastric nerve along the lesser curvature, it was found to pass to the margin of the ulcer, and its fibres were incorporated with the dense fibrous tissue of which the raised edges of the ulcer were composed. Plate I.

The continued suffering, and the daily rejection of food, at length completely exhausted the strength of the patient, and were the cause of her death; and although the pylorus was free from obstruction, and no hypertrophic change was evident in the muscular coat of the stomach after death, the symptoms were explained by the manner in which the branches of the

pneumogastric nerve were involved in the dense edges of the ulcer.

The attack of hæmatemesis four years before admission into the hospital, showed the chronic character of the disease; and, as the ulceration slowly extended into the surrounding vessels, the effusions of blood were repeated; but intervals of reparative action probably took place, for we find that there were periods during which the more severe symptoms were relieved, and the character of the ulcer, as shown by its general and microscopical appearances, warrants such a conclusion.

The left lobe of the liver was small and atrophied, and its section presented several large vessels surrounded by contractile tissue, without any intervening gland-structure. Chronic change had extended into this part of the gland, and the obliteration of the vessels had led to the atrophy of the whole left lobe.

CASE LV.—Chronic Ulceration of the Stomach, with Painter's Colic. Perforation.—George O—, æt. 28, was admitted into Guy's Hospital February 22nd, and died April 16th. He was a married man, of anæmic appearance, but with dark hair. For ten years he had been a painter, but previously he had employed himself as a publican, and he had then drank freely. Except an attack of fever several years before, he had been well till nine months before admission, when he was seized with colic, which came on with vomiting, obstinate constipation, and severe griping pain at the umbilicus; from that time he lost flesh, and had constant pain in various parts of the abdomen; the bowels became constipated, and he suffered from pain between the shoulders. A fortnight before admission he had hæmatemesis, and he became very anæmic; this condition persisted till his application at Guy's; when the tongue was pale and the respiration easy; there was griping pain, also, in the abdomen, and constipation; but the abdomen was supple, and no disease could be detected on palpation. Compound Soap pill, gr. v., was given every six hours.

On Feb. 29th—Vomiting of grumous matter came on; an enema was administered, and an electro-galvanic current applied to the spine and abdomen; the galvanism produced some uneasiness.

On March 3rd—There was great pain in the abdomen, vomiting of grumous coffee-ground substance; rapid prostration came on; and death took place in a few hours.

Inspection, twenty-nine hours after death.—On opening the peritoneal cavity it was found to contain a considerable quantity of gas; and grumous matter was extravasated from the stomach. Upon raising the left lobe of the liver an opening, about the size of a pea, was observed in the lesser curvature of the stomach; the mucous membrane was generally thickened, and about half an inch from the pylorus, near the lesser curvature, was an oval ulcer, about three inches by two in size, with raised and everted edges; the floor of

the ulcer was formed by the pancreas, covered by white fibrous tissue; the opening before mentioned was at the anterior part of this ulcer. The colon contained a large quantity of scybala.

There was considerable difficulty in the diagnosis of this case; the evident indication of colic, in a painter of intemperate habits, rendered the symptoms of ulceration of the stomach more than usually obscure. The attack of hæmatemesis might have been attributed either to ulceration or congested portal circulation; but the vomiting of coffee-grounds substance indicated the slow effusion of blood; and it afforded a very characteristic symptom of chronic organic disease. The immediate cause of the perforation was not apparent; but, although we could not affirm that the muscular contraction produced by the galvanism was in any manner the cause of the suddenly fatal issue, still, if the chronic ulcer of the stomach had been as prominently before the mind as the lead colic, it is probable that the use of galvanism would not have been directed.

CASE LVI.—Perforating Ulcer of the Stomach, with a second small Chronic Ulcer in the same organ.—Harriet B—, æt. 22, a single woman, a milliner, who worked in the city, but resided at Bethnal Green, was stated to have enjoyed good health, with the exception of occasional pain, and other uneasy sensations in the gastric region; but she had not any sickness, and she continued at work. The bowels were generally regular. On Friday, Oct. 20th, after eating at about 3 p.m. a full meal of anchovies and bread, she was seized with violent vomiting, followed by most intense pain, which commenced in the left hypochondriac region, and gradually extended over the abdomen. She said that the pain commenced by something giving way in her side. When seen, about 9 p.m., she was in a state of great prostration; and she died at 11 a.m. on the 21st, twenty hours from the commencement of the attack.

The body was well nourished. On opening the abdomen, the viscera were found distended, and covered with a coat of recent lymph, and with some castor oil, which had been administered by the friends shortly after the attack. In the stomach were two ulcers, one with raised and rounded edges, about half an inch in diameter, extending to the muscular coat; the other about the same size, but with a smaller opening in the muscular, and a round, smooth, small punch-hole opening perforating the peritoneal sac. Both were situated at the lesser curvature, and the latter, in which was the peritoneal perforation, was towards the posterior part. The remaining viscera appeared healthy.

This case was peculiarly interesting: 1st, in presenting two

ulcers in the stomach, but in different stages, the one resembling ordinary chronic ulcer with thickened edges, the other having the appearance of but slight action around it; 2nd, in the early age of the patient, twenty-two, the presence of but slight symptoms, pain but no sickness, although an ulcer existed in the stomach; for dyspepsia was the only precursor of the fatal perforation, and she considered herself in good health, and continued at work; 3rd, in the sedentary employment, constrained position, irregular meals, uncertain hours, and probably only scanty fare, all tending to impair nutrition, and act as causes of this fatal malady; 4th, in the occurrence of the perforation being, as is usually the case, after a meal; 5th, in the pain commencing at the hypochondriac region, for this is not always the case, and we cannot thereby ascertain at all times the seat of perforation; sometimes, even, the seat of pain suggests cæcal instead of gastric disease; and 6thly, in showing the folly of at once giving a purgative draught, for the castor-oil was here found floating in the peritoneal cavity.

CASE I.VII.—Perforation of the Stomach.—Henry F. B—, æt. 53, was admitted into Guy's, July 11th, 1860, in a dying state, and expired the same evening at 6 p.m. He had been ailing for several months, but no history was obtained.

On inspection, general peritonitis was found; and at the lesser curvature of the stomach was a chronic ulcer about the size of half-a-crown, with raised and thickened edges; its floor was formed in part by the pancreas; but at the anterior part was a small, round opening, extending into the peritoneum, which had caused the fatal attack; a branch of the pneumogastric nerve extended to the base of the ulcer.

Although the termination of this case in perforation, and peritonitis, was unexpected, there had been some indications of ill health for several months.

CASE LVIII.—Chronic Ulceration. Death from Perforation.—A man, æt. 37, had been subject to dyspepsia, constipation, and general abdominal uneasiness, but he had had no vomiting; intense pain came on suddenly, and he died in a few hours.

In the *stomach* a circular ulcer was found about the size of a five-shilling piece, the edges were rounded, and the mucous membrane was more extensively destroyed than the muscular; the base of the ulcer was formed by the pancreas and by condensed cellular tissue; and quite at its upper margin, below the left lobe of the liver, there was a small perforation, which had led

to general peritonitis. The microscopic examination of the ulcer showed it to be of a simple inflammatory character.

The absence of hæmatemesis, of vomiting, and of other symptoms of gastric ulcer, except dyspepsia, was remarkable in this case, for the ulcer was of large size, and was not of recent formation.

CASE LIX.—*Chronic Ulcer of the Stomach. Perforation.*—Mary C—, æt. 21, was admitted August 28th, 1858, about noon, and died in ten minutes. She had been in a situation near the hospital, and had suffered from vomiting occasionally for five years, but had never had hæmatemesis.

Inspection was made about three hours after death. There was general peritonitis. In the stomach there were two ulcers, each about three-eighths of an inch in diameter; both were situated at the lesser curvature, the one at the anterior, the other at the posterior part of the stomach. Their edges towards the stomach were thickened and bevelled; the anterior one presented a small punch-hole opening extending into the peritoneum, which had led to the fatal peritonitis. The posterior one had also perforated all the coats, but extravasation had been prevented by the adhesions. The other portions of the mucous membrane of the stomach were healthy, presenting only slight injection; the other viscera were healthy.

In this case there had been evidence of disorder of stomach for several years; and the appearance of the ulcer also indicated chronic action in its thickened edges, but there had been no vomiting of blood.

CASE LX.—*Perforating Ulcer of the Stomach.*—A servant girl, æt. 27, residing in the Borough, was suddenly seized with severe pain in the abdomen; collapse came on, and death followed in twenty-four hours. On *inspection* a round ulcer was found on the anterior surface of the stomach; on the mucous surface it was half an inch in diameter, but on the peritoneal only a quarter of an inch, and there was very little thickening. Its perforation into the peritoneum had set up fatal peritonitis. Other portions of the mucous membrane of the stomach were partially dissolved. Remaining viscera were healthy.

This was one of the instances in which there was very little indication of disease prior to the fatal attack, and the appearance of the stomach after death showed an absence of thickening, and of chronic action.

CASE LXI.—*Chronic Ulcer of the Stomach. Perforation except the Peritoneum. Fatal Peritonitis.*—(Reported by Mr. A. W. A. Evans.)—William M—, æt.

29, a policeman, was admitted into Guy's Hospital, October 28th, 1859, under Dr. Addison's care. He had been in the police force for six weeks on night duty, when he was seized with pain a little below the scrobiculus cordis, with vomiting about an hour after food. There was pain also at the right shoulder, and a little blood was found in the vomited matter, but no sarcinæ. He had suffered habitually from constipation. Three years previously he had had a similar attack, from which, however, he recovered in three months. He afterwards enjoyed good health, although he had vague and intermittent pains at the epigastrium. He was a tall, sallow, and anæmic man, and on admission suffered from rather severe pain at the epigastric region, but nothing could be felt on tactile examination. He stated that he had passed dark-coloured stools; the tongue was large and white; the pulse 88. Magnesia mixture, with hydrocyanic acid, was prescribed three times a day. A castor-oil injection was administered at night, and beef-tea and arrow-root allowed. A small blister was afterwards applied at the scrobiculus cordis, and zij of sherry wine, with two eggs, &c., were ordered. On November 2nd, nitrate of bismuth was given, with mucilage mixture. The bowels then became constipated, and the magnesia mixture was repeated with dilute hydrocyanic acid, tincture of opium, and of lavender. On the 4th he passed a black evacuation, but the constipation continued. On the 14th, gr. xv of colocynth and calomel were given, and a castor-oil injection administered, which acted very freely on the bowels. On the 16th, severe pain came on in the abdomen; he became collapsed, and had vomiting, with excessive tenderness of the abdomen; a grain of opium was then given every six hours; but on the 17th he was evidently sinking, and died at nine. On *inspection* the peritoneum was found to be universally inflamed, and the intestines were glued together by recent lymph; older adhesions were found between the stomach and the parietes at the superior part of the abdomen. The stomach was slightly hour-glass in form; and about one and a half inches from the pylorus on its posterior aspect was a chronic ulcer; all the coats were destroyed, and the pancreas formed the base of the ulcer, which was two inches in diameter. The mucous coat was more extensively destroyed than the muscular. A smaller ulcer, directed anteriorly, united with the posterior and larger one, its base was formed by delicate fibro-cellular tissue and peritoneum; this thinned, *almost perforated* portion, was a quarter of an inch in diameter, and at its centre was a transparent spot, only one eighth of an inch across; water, however, did not exude. The perforation was not complete; but there was no doubt that sufficient transudation had taken place to produce the fatal peritonitis. The thoracic and other viscera were healthy.

The condition of the ulcers in the stomach were peculiar; two ulcers had become united, but the posterior portion was the older formation, and that had a firm base of pancreatic tissue; not so, however, the anterior one; its base consisted of fibro-cellular tissue and peritoneum, but the serous membrane was still entire. In typhoid ulceration of the intestine we have

several times observed peritonitis produced when the peritoneum remained entire, transudation having taken place; and it is probable that a similar occurrence ensued in this instance. It illustrates, also, the caution that should be used in the administration of purgative medicines.

CASE LXII.—Perforation of the Stomach. Local Suppuration in the Peritoneum. Pleuro-pneumonia.—M. A. B—, æt. 33, was admitted into Guy's Hospital, under my care, September 20th, 1860. She was a rather large woman, who stated that till the 18th she had enjoyed good health; at two a.m. of that day, she had been seized with severe pain in the left side, which continued till admission. I did not see her at the time of admission; the pain was then very severe, and she was very ill.

On the 21st, she was in less pain, and the countenance was less distressed; the tongue had white fur on it, and was rather dry; deep inspiration produced severe pain in the left side, but no friction sound could be heard; air entered the lower lobe of the left lung very imperfectly; and in fact respiration was scarcely audible either in front or behind on the left side. There was no abnormal sound with the heart; the pulse was compressible, and the skin hot. The resident medical officer had ordered gr. xv of colocynth and calomel, and pills twice a day, containing calomel, antimony, and opium.

On the 22nd, she still complained of severe pain in the side; there was dulness at the base of the left lung, and scarcely any air entered that lung; the skin was hot; the tongue was dry and furred; the pulse was compressible; the bowels were opened freely. She was ordered to be cupped on the left side to 3vj; to take soap and opium gr. v every night; and julep of acetate of ammonia three times a day.

On the 24th, the pain in the side was much less severe, but she was still very ill, complaining of tenderness of the abdomen; the tongue had a whitish fur; the bowels had not been again acted upon. The abdomen was supple, not tympanitic, but there was tenderness at the lower part. She stated, on the previous week, immediately after menstruation, she had been exposed to cold, and that then the pain came on. Julep of ammonia was given three times a day, and the opium pill continued.

On the 25th, she complained of an offensive taste in the mouth, and of increased pain in the left hypochondriac region, but she bore considerable pressure without aggravation of her suffering; the tongue was furred; the mouth was clammy; the pulse compressible, and slightly excited. She very much desired to have soda water, and was ordered effervescing mixture, with carbonate of magnesia, gr. x every four hours, and wine 3vj.

On the 29th, she was still distressed with an offensive taste in the mouth; the tongue was furred; there was no tenderness of the abdomen, nor distress in breathing. Opium was given night and morning, and soda water occasionally.

On the 30th, she was seized with eructation, and vomiting of thin offensive pus; to use the words of the sister of the ward as if an abscess had broken; she became exceedingly prostrate, and died on the 2nd October at 9.50 a.m.

Inspection on the same day at 2:30 p.m. The pleura was adherent at the lower lobes of both lungs. The lungs were consolidated in lobules at each of the lower lobes; some of these were red and dense, others gray and softening down; and these pneumonic lobules were becoming continuous the one with the other. The heart was healthy, and the left ventricle was contracted. On opening the abdomen the intestines were found to be moderately distended; the parietes were fatty, as was also the omentum. The liver was elongated and pale, and it extended nearly to the crest of the ileum; it was very fatty, and weighed 6 lbs. On the left side there was a large abscess in the peritoneum bounded by the spleen, the left lobe of the liver, the stomach, &c. The walls of the abscess were covered with a layer of greenish-yellow lymph, and the abscess contained about two pints of gray offensive pus. The abscess communicated with the stomach by a small punch-hole perforation, situated about the middle of the lesser curvature of the stomach on the anterior aspect. The edges of the ulcer were bevelled internally, but they were not thickened. The mucous membrane of the stomach was thickened and ecchymosed in patches; there was some contraction of the mucous membrane of the stomach opposite the ulcer over the pancreas, as if from the cicatrix of a former ulcer. The kidneys were pale; the intestines and ovaries were healthy.

The diagnosis of this case was obscure: on admission, it appeared to be diaphragmatic pleurisy; but, afterwards, it was evident that the disease was below, not above, the diaphragm. There was an absence of the ordinary signs of peritonitis, as shown by tenderness and distension of the abdomen; still, there was pain of moderate degree in the lower part of the abdomen, but especially on the left side. There cannot be a greater proof of the absence of the ordinary signs of peritonitis, than that a gentleman of great experience ordered, in my absence, free purgatives of colocynth and calomel, &c. The patient was very restless, but her principal complaint was of the offensive taste in the mouth, and she incessantly begged to have soda water. The adhesions had localised the inflammation, and there was none of the collapse which marks perforation into the general cavity of the peritoneum. For several days the suppuration increased in the abscess, but it evidently communicated with the stomach; and the decomposition of its contents led to the offensive character of the breath which distressed the patient so much. It is probable that the pneumonia arose from pyæmia; and it did not come on till a short time before death. The sudden discharge of pus only a few hours before death indicated the rupture of an abscess; but it was even then very doubtful whether it proceeded from a thoracic or abdominal source.

CASE LXIII.—*Chronic Ulceration of the Stomach, extending to the Diaphragm, and simulating Pneumothorax.*—Barbara —, æt. 39, was a married woman, who for eighteen months had suffered from symptoms of dyspepsia or chronic gastritis, from pain between the shoulders and at the epigastrium, and from vomiting. Two days before admission she was seized with intense pain in the left side and shoulder, and she had urgent dyspnœa. On examination at the base of the left lung there was resonance, amphoric breathing, metallic tinkling, and ægophony. She died twenty days after the attack of dyspnœa.

Dr. Barlow's diagnosis was confirmed; there was pleurisy on the left side, and a large peritoneal abscess, which communicated by two openings with the lesser curvature of the stomach; one near the œsophagus, capable of admitting the middle finger, and another, smaller, near the anterior wall. The abscess was bounded by the ribs, spleen, liver, and diaphragm; and by inflammatory adhesions in a partially sloughing condition. The pleura in this case was not perforated, so that the symptoms of pneumothorax were produced by the air in the peritoneal abscess.

The post-mortem examination in the following remarkable case is here introduced, as an illustration of perforation of the stomach, associated with perforation of the colon, as well as of the diaphragm. The disease, however, in the colon, appeared to be of longest duration, and to have led to the formation of the fæcal abscess; whilst the perforation of the stomach was, perhaps, secondary, and from the external to the internal surface.

CASE LXIV.—*Fæcal Abscess, connected with the Stomach, the Lung, the Spleen, and the Transverse Colon.*—Ellen R—, æt. 25, was admitted in July, and died August 5th, 1847. History not known, except that she had fæcal vomiting.

On inspection, the left lung was found adherent by old adhesions; a vomica was found at the upper lobe, and gray hepatization existed at its lower part. *Abdomen.* The intestines were matted together by old adhesions; the liver also was adherent to the stomach. On the left side was an abscess of some extent, bounded in front and to the outer side by the ribs, to the inner side by the stomach and by the spleen, below by the transverse arch of the colon, above by the diaphragm and right lung. This abscess communicated with the chest by an opening through the diaphragm, and was there bounded by the lower surface of the right lung and thickened pleura. The abscess communicated also by two separate openings with the greater curvature of the stomach, and by one opening with the transverse colon; it was filled with partially coagulated blood; the upper portion of the spleen was found sloughing in the cavity of the abscess. The stomach also, and the transverse colon

contained each of them a considerable quantity of blood. The openings in the stomach were round holes, having tolerably smooth edges, and the mucous coat was not thickened; the peritoneal coat appeared as if it had been ruptured. There were several ulcers in the transverse colon, which communicated with the abscess, and had considerable thickening of the gut around them. The liver was large and fatty. Some of the mesenteric glands were calcareous. The contents of the pelvis were all bound together by old adhesions, and there was a considerable quantity of recent lymph at this part. Between the rectum and the bladder was an abscess communicating with the rectum; there were several other ulcers in the rectum, and the whole mucous membrane was intensely injected, and of a deep purple colour. There was strumous ulceration of the mucous membrane of the uterus.

CASE LXV.—*Chronic Ulcer of the Stomach. Peritoneal Abscess. Perforation of the Diaphragm. Empyema.*—J. F. L., æt. 64, a basket maker by trade, was admitted into Guy's Hospital, under Dr. Gull's care, May 14th, 1860. He had inflammation of the lungs when thirty years of age, and a year afterwards he had rheumatic fever. Seven years before admission he had a blow by a capstan, which threw him into the hold of the vessel, and from that time he had always suffered pain on taking cold. Seventeen or eighteen months previous to admission, he had also noticed a swelling at the epigastric and umbilical regions, and he had nausea and vomiting after taking food, but no blood was vomited. Severe tearing pain at the stomach was produced by retching, and he also experienced at those times a burning sensation, which lasted for twenty-four hours. The swelling had not increased, but had extended more generally over the abdomen; and for a few days the pain had reached the loins. For two weeks he had felt severe pain above the crest of the ileum on each side; it came on after attempting to work, and it passed up each side behind the clavicle. The painful part began to swell, and continued swollen for two days, but the swelling afterwards disappeared. From the commencement of his illness, he stated, that he had had a sense of tearing whenever he lifted his left arm; and he had frequently felt severe pain below the last rib on the left side; the pain was also felt in the lumbar region. He had suffered much from flatulence, but not from sickness; the bowels were generally constipated. He was a spare man, with an anxious expression. The chest on admission was normal. The liver could be felt in the epigastric region; the spleen was also enlarged. The abdomen was increased in size; it was tender and tympanitic; but no fluctuation could be felt. The urine was not albuminous, its sp. gr. was 1.020. He was ordered iodide of potassium, with the bicarbonate of potash in mint julep. On the 19th, the pain in the abdomen was aggravated so that he could scarcely bear the pressure of the bedclothes; the bowels were confined. Half a grain of opium was given twice a day, and a simple injection was administered. During the afternoon of the same day intense pain came on, and extended from the crest of the ileum to the side of the thorax. On the 20th he seemed to be in severe pain, and he suffered from great dyspnoea; the face was pallid; there was an anxious expression of countenance; and he had slight cough and scanty expectoration; the abdominal muscles did not move; the base of the left lung was dull, both anteriorly and

posteriorly, and there was an absence of any respiratory sound ; but there were tactile vibration and vocal resonance. The abdomen was tense ; the skin was hot, but the extremities cold ; and he complained of great thirst. On the 21st there was still great pain on the left side of the chest, the left side was dull on percussion except at the apex posteriorly ; there was an entire absence of respiratory murmur and vocal resonance at the base of the left lung, but passing upwards the breathing became bronchial ; and higher still, it became amphoric, with almost an ægophonic voice and great resonance on coughing. The heart sounds were distinct on the right side of the sternum ; and the dulness extended as far as that part. On the right side generally the respiration was puerile, and accompanied with bronchial râles ; the pulse was 104 ; the tongue moist and furred ; the bowels were open ; there was no appetite, but there was great thirst, and the breath was sweet. A blister was applied ; acetate of potash with nitric ether, and compound tincture of camphor were prescribed, and four ounces of gin, afterwards increased to eight, were ordered. The patient, however, rapidly sank, and died the following day at half-past four. *On inspection*, the abdomen was enlarged. The left pleura was full of pus, and was lined with a flocculent membrane ; the heart was pushed over beyond the median line, and the left lung was compressed. The right pleura and lung were healthy, so also the pericardium and the heart. On opening the abdomen, the left lobe of the liver was found to be adherent at the scrobiculus cordis by firm tissue, and it had formed the hard mass felt during life ; the peritoneum generally was healthy. On drawing down the stomach, and removing the left lobe of the liver, a large abscess was found between the stomach and the diaphragm ; and it contained nearly a pint of pus. The abscess was bounded by the liver in front and on the right, by the spleen on the left, and by the pancreas posteriorly. In the diaphragm was a small opening about the size of a crow quill, and thus a communication had been formed between the abscess in the abdomen and the pleura. At the lesser curvature of the stomach, and on its posterior aspect, was a chronic ulcer two and a half inches long, bounded behind by dense cicatrized tissue on the pancreas, but no perforation could be detected. The liver was fatty, the other viscera were healthy.

Dr. Gull correctly diagnosed this case as one of chronic disease of the stomach, leading to perforation of the diaphragm, and causing the fatal attack of pleurisy. The swelling at the epigastric region, accompanied with nausea, vomiting, and "tearing pain," were strongly indicative of gastric disease ; but it was very doubtful whether the malady was of a cancerous character. It was doubtful, also, whether the blow he received from a capstan had anything to do with his subsequent disease. Chronic ulceration took place, perforation of the walls of the stomach followed ; but the mischief was localised by adhesions ; suppuration then took place, beneath the diaphragm ; this

muscle at last itself became perforated, and intense pleurisy was at once set up by the extravasation of pus into the pleural serous membrane.

CASE LXVI.—Chronic Ulceration of the Stomach. Perforation. A Sinus extending into the left Lung. Gangrene. Empyema. Second Chronic Ulcer.—Eliz. F—, æt. 36, had been treated as an out-patient for dyspepsia, and it was supposed that ulceration of the stomach existed; the prominent symptom was vomiting of coffee-ground matter. After admission into the hospital she became extremely low and emaciated, and gradually sank. It was then believed that she had cancerous disease. She died October 13th, and was examined twenty-six hours after death.

Chest.—The left pleura contained purulent effusion. The lower lobe of the left lung was pneumonic and adherent to the diaphragm; a vertical section of this lobe exhibited an excavation, filled with dark-gray and tenacious matter, which exhaled a gangrenous odour. The cavity was traversed by pulmonary vessels, which, when placed under water, had a curious flocculent appearance; a sinus passed from this cavity, through several fistulous openings in the diaphragm, into the stomach. The heart and pericardium were normal, but the foramen ovale was open. In the abdomen there were chronic and vascular adhesions between the viscera and parietes, more particularly about the right hepatic lobe; the liver was situated unusually low in the abdomen. The small intestine appeared perfectly healthy; the kidneys were coarse, and the tunic adherent. The liver and gall-bladder were healthy.

On opening the stomach, along the greater curvature, an opening of a circular figure was discovered in its walls, the circumference of which, with the exception of a small aperture at its upper border, was very firmly adherent to the under surface of the left lobe of the liver. This appearance, the remains of old ulceration, was situated in the region of the lesser curvature of the stomach. From the perforation in the ulcerated walls of the stomach a sinus passed upwards, bounded upon the left by the spleen, on the right by the left lobe of the liver, and behind by the pancreas and small omentum; above, it extended to the diaphragm, which was perforated by several foramina, and the sinus communicated with a cavity in the inferior lobe of the left lung; the surfaces of the organs bounding this sinus were tinged with a dark-gray hue. The opening from the ulcer in the stomach was valvular, and was situated under its superior border. The stomach contained dark, almost black, thick viscid fluid; there was also a second chronic ulcer near the pyloric extremity of the stomach.

The diagnosis of this case was obscure; the earlier symptoms indicated ulceration of the stomach, but the unusual prostration led to the idea that the disease was of a cancerous character. The disease commenced in the stomach, and the ulcer at the lesser curvature led to perforation; the aperture, however, was

at the posterior aspect, and it was also surrounded by adhesions, so that it passed into the structures between the diaphragm and the stomach without leading to general peritonitis. Circumscribed suppuration then took place, and ulceration extended through the diaphragm. Here, also, adhesions between the pleural surface of the lower lobe of the left lung, and that of the diaphragm, prevented acute pleurisy being at once produced; the lung tissue was perforated, and a sloughy cavity was formed. The pleura subsequently became acutely diseased, and effusion of pus took place in the non-adherent part of the serous membrane. The gangrene of the lung and the empyema led to the excessive prostration in this case. A second ulcer was found in the stomach.

CASE LXVII.—Chronic Ulceration of the Stomach. Fatal Hæmorrhage. Perforation of the Splenic and of the Pancreatic Arteries.—Charlotte T—, æt. 55, was admitted March 4th, 1857, into Guy's Hospital, and died March 5th, at 9.40 p.m.

She had previously been admitted under Dr. Oldham's care, in a very blanched condition, and complained of severe pain in the left side; she had had no vomiting nor spitting of blood, and her appetite had failed; whilst in the hospital, however, she took food well. She was in a week's time transferred to Dr. Wilks's care, and was then evidently suffering from internal hæmorrhage; she had great pain and uneasiness in the left side, with nausea, but did not vomit. During the night she vomited a cupful of blood, and shortly afterwards died. She had been a charwoman, and of intemperate habits, and six years before death she had vomited blood.

On inspection, the pleura was found adherent, and the lungs healthy. The left ventricle was contracted and empty, as in death from loss of blood.

In the abdomen the peritoneum was healthy, except adhesion at the upper part, where the anterior wall was firmly united to the stomach and liver. These structures could be separated with care, except at the left hypochondriac region, where the adhesions were exceedingly firm. The whole of the liver, stomach and spleen were removed together; the stomach was found to be contracted at its centre by a large oval ulcer placed transversely; two pouches were formed, the pyloric being the smaller of the two, and the cardiac one formed a large cavity, capable of holding at least a quart of fluid; each part contained a large quantity of coagulated blood, partly digested. At the posterior part of the stomach, near the lesser curvature, was a large chronic ulcer, with raised dense rounded edges, and with a depressed slightly granular centre; the ulcer was oval or rather reniform in shape, and appeared to be formed by two ulcers which had coalesced; it was at least three inches in length, and one and a half to three in breadth. Its floor was formed partly by the left lobe of the liver, to which it was firmly

muscle at last itself became perforated, and intense pleurisy was at once set up by the extravasation of pus into the pleural serous membrane.

CASE LXVI.—Chronic Ulceration of the Stomach. Perforation. A Sinus extending into the left Lung. Gangrene. Empyema. Second Chronic Ulcer.—Eliz. F—, æt. 36, had been treated as an out-patient for dyspepsia, and it was supposed that ulceration of the stomach existed; the prominent symptom was vomiting of coffee-ground matter. After admission into the hospital she became extremely low and emaciated, and gradually sank. It was then believed that she had cancerous disease. She died October 13th, and was examined twenty-six hours after death.

Chest.—The left pleura contained purulent effusion. The lower lobe of the left lung was pneumonic and adherent to the diaphragm; a vertical section of this lobe exhibited an excavation, filled with dark-gray and tenacious matter, which exhaled a gangrenous odour. The cavity was traversed by pulmonary vessels, which, when placed under water, had a curious flocculent appearance; a sinus passed from this cavity, through several fistulous openings in the diaphragm, into the stomach. The heart and pericardium were normal, but the foramen ovale was open. In the abdomen there were chronic and vascular adhesions between the viscera and parietes, more particularly about the right hepatic lobe; the liver was situated unusually low in the abdomen. The small intestine appeared perfectly healthy; the kidneys were coarse, and the tunic adherent. The liver and gall-bladder were healthy.

On opening the stomach, along the greater curvature, an opening of a circular figure was discovered in its walls, the circumference of which, with the exception of a small aperture at its upper border, was very firmly adherent to the under surface of the left lobe of the liver. This appearance, the remains of old ulceration, was situated in the region of the lesser curvature of the stomach. From the perforation in the ulcerated walls of the stomach a sinus passed upwards, bounded upon the left by the spleen, on the right by the left lobe of the liver, and behind by the pancreas and small omentum; above, it extended to the diaphragm, which was perforated by several foramina, and the sinus communicated with a cavity in the inferior lobe of the left lung; the surfaces of the organs bounding this sinus were tinged with a dark-gray hue. The opening from the ulcer in the stomach was valvular, and was situated under its superior border. The stomach contained dark, almost black, thick viscid fluid; there was also a second chronic ulcer near the pyloric extremity of the stomach.

The diagnosis of this case was obscure; the earlier symptoms indicated ulceration of the stomach, but the unusual prostration led to the idea that the disease was of a cancerous character. The disease commenced in the stomach, and the ulcer at the lesser curvature led to perforation; the aperture, however, was

at the posterior aspect, and it was also surrounded by adhesions, so that it passed into the structures between the diaphragm and the stomach without leading to general peritonitis. Circumscribed suppuration then took place, and ulceration extended through the diaphragm. Here, also, adhesions between the pleural surface of the lower lobe of the left lung, and that of the diaphragm, prevented acute pleurisy being at once produced; the lung tissue was perforated, and a sloughy cavity was formed. The pleura subsequently became acutely diseased, and effusion of pus took place in the non-adherent part of the serous membrane. The gangrene of the lung and the empyema led to the excessive prostration in this case. A second ulcer was found in the stomach.

CASE LXVII.—*Chronic Ulceration of the Stomach. Fatal Hæmorrhage. Perforation of the Splenic and of the Pancreatic Arteries.*—Charlotte T—, æt. 55, was admitted March 4th, 1857, into Guy's Hospital, and died March 5th, at 9.40 p.m.

She had previously been admitted under Dr. Oldham's care, in a very blanched condition, and complained of severe pain in the left side; she had had no vomiting nor spitting of blood, and her appetite had failed; whilst in the hospital, however, she took food well. She was in a week's time transferred to Dr. Wilks's care, and was then evidently suffering from internal hæmorrhage; she had great pain and uneasiness in the left side, with nausea, but did not vomit. During the night she vomited a cupful of blood, and shortly afterwards died. She had been a charwoman, and of intemperate habits, and six years before death she had vomited blood.

On inspection, the pleura was found adherent, and the lungs healthy. The left ventricle was contracted and empty, as in death from loss of blood.

In the abdomen the peritoneum was healthy, except adhesion at the upper part, where the anterior wall was firmly united to the stomach and liver. These structures could be separated with care, except at the left hypochondriac region, where the adhesions were exceedingly firm. The whole of the liver, stomach and spleen were removed together; the stomach was found to be contracted at its centre by a large oval ulcer placed transversely; two pouches were formed, the pyloric being the smaller of the two, and the cardiac one formed a large cavity, capable of holding at least a quart of fluid; each part contained a large quantity of coagulated blood, partly digested. At the posterior part of the stomach, near the lesser curvature, was a large chronic ulcer, with raised dense rounded edges, and with a depressed slightly granular centre; the ulcer was oval or rather reniform in shape, and appeared to be formed by two ulcers which had coalesced; it was at least three inches in length, and one and a half to three in breadth. Its floor was formed partly by the left lobe of the liver, to which it was firmly

adherent, and by the pancreas. Two small papilliform eminences were found on careful examination, and a bristle could easily be passed into open vessels; one opening was found to communicate directly with the splenic artery, on the upper margin of the pancreas, and a second with the artery in the centre of the pancreas. Each of the perforations in these vessels had a small quantity of blood at their orifices, but did not contain any clot or blood. The pylorus and the rest of the stomach were healthy. The intestines contained a considerable quantity of blood, but were otherwise healthy. The portion of the left lobe of the liver in connexion with the stomach, was atrophied, and it presented fibroid degeneration; the other part of the liver was fatty. The kidneys were granular and degenerated.

This case presents us with an unusual mode of termination of gastric ulcer. The ulceration had been slow in its progress, and it had apparently extended over at least six years, or more: there had been some hæmorrhage, which had probably come from some of the branches of the gastric arteries; as the ulceration extended the walls of the stomach were perforated, but adhesions prevented peritonitis. In this state her health had become impaired by disease of the kidneys, which were found after death in a state of advanced degeneration; slow ulceration had reached the vessels at the base of the ulcer, and the perforation of two large vessels led to the fatal hæmorrhage. These vessels were healthy, but the ulceration had destroyed the surrounding structures more extensively than the arteries, and the contraction of the vessels was also prevented by fibro-elastic tissue, so that minute papillary eminences were formed; the disease of the kidneys, and the condition of the blood also tended to increase the hæmorrhage. It is remarkable that so little blood was vomited, although the stomach was full, and the intestines contained a considerable quantity. The absence of this symptom arose partly, perhaps, from the adhesions of the stomach to the parietes, as well as from the prostrate condition of the patient.

As to the cause of the complaint, we are led to suppose that the intemperate habits of the patient produced the disease of the stomach, as well as that of the kidneys; the one tended to increase the other, and at last hastened the fatal termination.

CASE LXVIII.—*Ulceration of the Stomach. Fatal Hæmorrhage.*—

Joseph G—, æt. 53, was admitted into Guy's, February 28th, and died March 6th.

This patient was admitted after hæmatemesis had taken place; it came on suddenly, and there were no premonitory symptoms; he died on the sixth day, completely blanched.

On inspection, forty hours after death, a small ulcer about the size of a fourpenny-piece, was found at the lesser curvature of the stomach; it was round, depressed in the centre, and the edges of the mucous membrane were raised; in its centre was an opening from which exuded a drop of blood, and a probe could be passed into a large vessel beneath, apparently the gastric. The stomach was of normal size, and free from blood, but the large intestines contained blood, as shown by their dark colour. A partial inspection only was allowed.

CASE LXIX.—*Chronic Bronchitis. Emphysema. Chronic Ulcer of the Stomach.*—Samuel W—, æt. 60, admitted January 16th, 1861, and died January 19th. He had been of very intemperate habits, and several months before admission he had passed more than a quart of blood from the bowels. He was very ill from chronic bronchitis, which was the cause of his death.

Inspection.—The lungs were excessively distended with air; they were emphysematous, and dark carbonaceous patches were everywhere spread over their surface. The heart was hypertrophied; the aortic and pulmonary valves were thinned, and the double crescents of the valves were perforated. The kidneys were congested. The whole tract of the alimentary canal from the stomach was healthy; there were no piles nor any cicatrices in the rectum or colon. At the lesser curvature of the stomach, on the anterior surface, and midway between their pyloric and œsophageal orifices were two chronic ulcers, about the size of half-a-crown. Their edges were raised and bevelled, and the mucous membrane was entirely destroyed. They were pale in colour, and tenacious mucus adhered to their surfaces. The greater curvature presented considerable gastric solution. The muscular coat of the rectum and sigmoid plexure were hypertrophic. In the centre of the larger ulcer was a small projecting eminence, and on section of the ulcer it was seen to be the truncated extremity of a small vessel, filled up by a clot. The base and margins of the ulcer presented much firm, fibrous tissue.

In this case the severe bronchitis masked the gastric symptoms, and the patient was exceedingly ill from the thoracic disease when brought to the hospital. The source of the previous hæmorrhage into the stomach was apparent in the minute vessel obstructed by a coagulum of blood.

A most interesting and rare case of recovery after apparent perforation, is recorded by Dr. Hughes and Mr. Hilton, in the Guy's 'Reports.' The young woman left the hospital, and appeared convalescent; subsequent indiscretion in diet produced

a return of the symptoms, and a fatal result. A cicatrix of previous ulcer and adhesions were found, but with new perforation. The opiate plan of treatment of Drs. Stokes and Graves was adopted with unusual success.

CASE LXX.—*Chronic Ulceration, with Villous Growth. Stomach exceedingly contracted, simulating Cancer.*—Thomas F—, æt. 34 years, a married man, who resided at Dover, and followed the occupation of a fruiterer, was admitted into Guy's, June 30th, 1854, under my care, in the Clinical Ward, in a pale and exceedingly emaciated condition. With the exception of an attack of rheumatism fifteen years before, his health had been good till eight months prior to admission. He stated that eight months previously he took cold, and experienced pain in the chest, at the lower part of the sternum, accompanied with difficulty of deglutition. He obtained no relief, but the pain gradually increased in severity, and was accompanied with vomiting after food; his food was brought up directly after being swallowed, his own description being that it never seemed to reach the stomach, but was brought up unchanged; the vomiting sometimes subsided for several days, and he was thus able occasionally to retain fluid food; when this occurred he experienced relief from the sense of painful exhaustion. Emaciation had slowly increased. On admission, his exhaustion appeared extreme, but still he experienced no pain; the abdomen was collapsed; no tumour could be felt; the distress on swallowing was localised at the lower part of the sternum. At the base of the right lung there was dulness on percussion, and some tubular breathing, but no cough nor dyspnoea. He sank on the fourth day.

Inspection.—The lower lobe of the right lung was consolidated, granular, and very readily broke down. The heart was healthy. The peritoneum also was healthy. The stomach was so small and concealed that it was not at first perceived; it was exceedingly contracted and lobulated externally, resembling a portion of large intestine; it was about six inches in length and two in breadth. On laying it open, from the œsophageal to the pyloric orifice, it presented a very unusual appearance. At the pylorus, and extending along the greater curvature, was a deep excavation or ulcer, bounded by a sharp, slightly ulcerated border, the surface of which was smooth, and of a grayish colour. This ulcerated surface extended about half an inch beyond the pylorus; passing towards the cardiac extremity and along the lesser curvature, the mucous membrane appeared smooth, shining, and glazed; and towards the cardiac extremity presented several raised, circular patches; the largest of these, very near to the ulcer, was about one eighth of an inch in elevation, and about one inch in diameter, and was composed of villous folds, which appeared to radiate from the centre; floated under water, this growth from the mucous membrane had a very beautiful appearance; nearer to the œsophagus was another circular patch of a similar description; and on either side there were slight folds, but less elevated, and having a longitudinal arrangement. On taking a small portion of this villous growth, it was found to consist of very delicate plicated folds; scarcely any epithe-

lium was found on the surface, but numerous crystals, resembling triple-phosphates, were observed upon it; the growth was composed of cells of large size, from $\frac{1}{16}$ th to $\frac{1}{8}$ th of an inch in size, many oval, some angular; they contained granules, and large nuclei from the $\frac{1}{16}$ th to $\frac{1}{8}$ th of an inch. These cells were very similar to those found on the mucous membrane of a healthy stomach, or in connexion with the gastric follicles; and although they bore some resemblance to the cells observed in epithelial cancer, they were, I doubt not, merely changed secreting cells of the gastric follicles. A section of the growth rendered this more probable; immediately beneath the surface of the mucous membrane was a thick layer of these secreting cells, reaching to distended gastric follicles, which were lobulated and much distended by similar cells; beyond these enormously enlarged gastric follicles was a stratum of white fibrous tissue, from one sixteenth to one eighth of an inch in thickness; and similar tissue extended between the follicles themselves. All the growths from the membrane had a like structure. On the surface of the apparently smooth portion were several small isolated dendritic or imperfect villi, containing cells, as before described. Beneath the mucous membrane was a dense fibrous layer, and then hypertrophied muscular fibre. The hypertrophy of the muscular fibre was more marked towards the pylorus, but even there did not exist in an extreme degree. The examination of the ulcerated surface did not show any structure which indicated the disease to be of a carcinomatous character. The liver, pancreas, and the remaining abdominal viscera and glands, were healthy. One kidney was large and healthy; the other appeared atrophied. See Plate II.

The pathology of the case just detailed is of great interest; it could not be ascertained, from minute enquiries from the patient, that he had taken any poisonous or corrosive substance. There had apparently been inflammation of the mucous and submucous tissues, leading to very slow ulceration in one part; in another, to the development of contractile tissue in the substance of the membrane, and producing contraction of the whole organ. The villous growths at first gave the idea of epithelial cancer; but the presence of gland follicles in their normal arrangement, though much hypertrophied, and the absence of every other indication of cancer, lead me to the belief that these parts were merely portions of changed or hypertrophied mucous membrane. There was no glandular enlargement nor disease resembling carcinoma in any part of the body. The disease during life was believed to be carcinomatous, and located at the cardiac extremity of the stomach; the manner in which the food was at once regurgitated or rejected from the stomach, the unrelieved pain, and steady emaciation, seemed to warrant such a suppo-

sition. The acute disease at the base of the right lung was interesting, as illustrating the manner in which such disease in an exhausted subject may take place without general symptoms. There were no cough, dyspnœa, nor febrile symptoms; the pulse was quiet, and the tongue clean; still there were dulness and tubular breathing at that part, and the lung was found, on inspection, in the second stage of pneumonia.

Fatal Cases of Chronic Ulcer of the Stomach.

Eliza B—, æt. 21. Perforation at the posterior surface, limited by adhesions; death from exhaustion; nerves involved.

Elizabeth S—, æt. 32. Perforation at the posterior surface, limited by adhesions; death from exhaustion; nerves involved.

George O—, æt. 28. Perforation at the posterior part, and limited by adhesions; at the anterior margin perforation into the peritoneum; acute peritonitis.

Harriet B—, æt. 22. Perforation; two ulcers at the lesser curvature, anterior one perforated, and causing fatal peritonitis; the second ulcer posteriorly placed, and with thickened edges.

Henry B—, æt. 52. Perforation; chronic ulcer at lesser curvature, limited by adhesions with pancreas; at the anterior margin perforation into the peritoneum and fatal peritonitis.

Man, æt. 37. Perforation—as in the last case.

Mary C—, æt. 21. Perforation; two ulcers at the lesser curvature, both perforated, one anteriorly placed, causing fatal peritonitis; the other placed posteriorly.

Female, æt. 27. Perforation; ulcer on the anterior surface; fatal peritonitis.

William M—, æt. 29. Perforation; chronic reniform ulcer at the lesser curvature; the posterior part perforated, but adherent to the pancreas; at the anterior part perforation, except the peritoneum, causing fatal peritonitis.

Mary A—, æt. 38. Perforation; ulcer at the anterior part of the lesser curvature; local suppuration; pleuro-pneumonia.

Barbara —, æt. 39. Perforation; chronic ulcer; local suppuration; simulating pneumothorax.

Ellen B—, æt. 25. Perforation; communication with the colon; faecal abscess.

Eliza F—, æt. 36. Perforation; ulcer at the posterior part of the lesser curvature; sinus extending through the diaphragm; empyema; a second chronic ulcer.

John L—, æt. 64. Perforation; ulcer at the posterior part of the lesser curvature; peritoneal abscess; perforation of the diaphragm; empyema.

Charlotte T—, æt. 55. Perforation; chronic ulcer on the posterior aspect; extravasation prevented by adhesions; fatal hæmorrhage.

Joseph P—, æt. 53. Fatal hæmorrhage from ulcer at the lesser curvature.

Samuel W—, æt. 60. Ulcer at the lesser curvature; hæmorrhage from divided artery; death from chronic bronchitis.

Thomas F—, æt. 34. Chronic ulcer; contraction; villous growth; exhaustion.

It will be seen that out of the 18 cases just mentioned the walls of the stomach were perforated in 15 instances, including the one in which the peritoneum only was left; but in only 7 out of the 15 did the perforation cause fatal peritonitis; in 8 others adhesions formed, and prevented extravasation.

Causes.—There is much obscurity as to the cause of ulceration of the stomach. Some cases are preceded by a state of chronic inflammation of the whole mucous membrane, produced by intemperance or irregularity in diet. In others it appears probable that the general state of nutrition and of the nervous system act as predisposing causes. Mental depression or anxiety, scanty food, late hours at night, and insufficient exercise, pressure upon the scrobiculus cordis, either by direct girthing of the abdomen, or by constant and constrained position, as in milliners and shoemakers, or the striking of the epigastrium by the shuttle of the weaver, are also causes of gastric ulcer.

Treatment.—There are several objects to be sought for in the treatment of ulceration of the stomach:

1. The promotion of reparative action by sustaining and increasing general nutritive power.

2. The relief of distressing symptoms, pain, vomiting, hæmorrhage, pyrosis, constipation, &c.

3. The prevention of the extension of the disease.

4. The removal of its complications.

1. Almost the first consideration, and certainly one of the most important, is the administration of proper food. If absolute rest could be afforded to the stomach, the ulceration affecting its surface would probably in many cases rapidly heal; but, since this is almost impossible, it must be our object to give such forms of nutriment as will spare the stomach; and in seeking to accomplish this purpose, it must be borne in mind that the especial office of the stomach, and its peculiar secretion, is the solution of nitrogenous compounds. These elements are found in the flesh of animals, in beef and mutton, &c.; hence we generally find that solid animal food produces pain and vomiting, and must in most cases be avoided.* If, however, these elements of food be given, they must be in an unirritating form, as the less oleaginous kinds of fish, the sole, whiting, cod, &c., or poultry; or in a fluid state, as veal and mutton broth, clear soups, &c.; beef-tea often creates nausea and vomiting. Still more must hard and indigestible meats, preserved meats, and cheese be avoided. Oysters, sweetbread, can often be taken when more irritating diet would be rejected.

Starchy food is converted into sugar by the saliva and by the secretions in the intestine, and in that state is readily absorbed. So also, oleaginous substances are converted into an emulsion by the alkalies in the secretions of the mouth and intestine, and in the bile; so that these forms of diet, whilst they are demulcent and soothing to the diseased gastric surface, do not require the stomach in order to place them in a state ready for absorption. Good stale bread, biscuits, milk, starchy substances—as arrow-root, tapioca, maize, or Indian corn, flour, rice, &c., may thus be given to the patient; eggs often disagree, but may be taken in the form of light puddings; milk, also, when refused in its simple character, may be better tolerated by combination with isinglass, as in blanc-mange, or with soda water or lime water; and even cream and bacon are occasionally well borne.

* Corvisart believes that the pancreatic fluid promotes the solution of nitrogenous substances.

Rich soups, highly-seasoned dishes, peppers, mustard, &c., are better abstained from ; so also pastries, and food containing much insoluble material, as salads, unripe raw fruit, green vegetables, &c. It is, however, undesirable altogether to abstain from vegetables, for we may thus defeat our object, by inducing cachexia ; oranges, lemons, &c., may be often taken with benefit.

Again, it is most important that food should be partaken of slowly, and thoroughly masticated ; and it is better to take small quantities at a time, and to repeat the allowance more frequently, than to distend the stomach by a large and bulky meal ; about three to four hours should intervene in ordinary cases, but where there is great exhaustion, with irritability of the stomach, food may be required more frequently, and in very small quantities. Exertion, both mental and physical, should be avoided directly after meals ; in fact, every thing should be done to facilitate the process of digestion. It is well to abstain from alcoholic liquors if possible ; they tend to aggravate the disease, and should not, I think, be given, unless the circulation be failing, and there be tendency to syncope ; but, when required, brandy in small quantity, and well diluted, or the forms of sherry which contain the least quantity of sugar, are best. New wines, port, and imperfectly fermented malt liquors, generally disturb and distress the patient.

It is desirable to use every means in our power to improve the health, as exercise in the open air ; but over fatigue, or constrained positions, should be avoided. Moderate horse exercise, and bracing air, will sometimes afford more relief than medicinal agents, even when long continued ; but violent shaking is injurious. When a chlorotic or anæmic state has been produced, the preparations of steel, by restoring a more healthy condition of the blood, greatly facilitate reparative changes. We prefer the milder preparations, as the ammonio-tartrate or citrate ; the compound steel pill, with aloes and myrrh, or quinine with iron, as the sulphates or citrates conjoined, may also be beneficially prescribed.

It is obviously most desirable to administer that form of aliment which will nourish the body, so that healing may be favoured, but without irritating and disturbing the process

which is going on towards recovery. The difficulty is still more increased by the occasional irritability of the stomach itself; and this leads us to the consideration of the means we possess for the mitigation of distressing symptoms—pain, vomiting, hæmorrhage, pyrosis, constipation. For the relief of *pain*, opium or its alkaloid morphia is often the best remedy, in doses of $\frac{1}{4}$ to 1 grain of the former two or three times a day, or a few minims of the solution of the latter. Chloric ether, in doses of 10, 15, to 20 minims, will be found very efficacious, especially when combined with nitrate of bismuth in 10 to 20 grain doses. Chlorodyne is stated as being a valuable substitute, but I have no experience in its use. Dilute hydrocyanic acid, in doses of 3 to 5m, is also a useful adjunct in some cases, especially when given with alkalies. Both potash, soda, lime, and magnesia, have been used; they neutralize acid secretion, and oftentimes increase the anodyne power of the remedies previously mentioned—opium, morphia, chloric ether, &c. If, however, there be constipation, dryness of the tongue, and opium is not well borne, conium or henbane may be used as substitutes. The nitrate and oxide of silver in doses of $\frac{1}{4}$ to 1 gr., in some instances diminish the pain and irritability of the stomach, especially when the gastric symptoms are associated with pyrosis. Creasote in 1m doses we have found very effectual in relieving pain, when accompanied with irritability of the stomach, vomiting, and fermentative changes in the food. Again, carbonic acid, as in ordinary soda water, is effective in relieving pain as well as vomiting. So also the use of cold water and ice, which are often very grateful to the patient.

Vomiting is a very distressing symptom in many cases of ulcer of the stomach. It is best combated by only partaking of fluid diet, and of that in moderate quantities. The remedies we have already mentioned are of service, but especially bismuth, hydrocyanic acid, creasote, ice, and the alkalies.

Dr. Jenner has pointed out the value of the sulphite of soda in checking the fermentative action, and the development of sarcinæ in obstruction from chronic ulcer, as in cancerous or pyloric disease. It may be given in ʒj doses, alone or conjoined with other agents; the hyposulphite is also given in similar cases.

Counter-irritants are often of service for the relief of pain and vomiting in these cases. A small blister applied to the scrobiculus cordis, or the croton oil rubbed in so as to produce pustular eruption. Some even use a seton; but I think that milder remedies may attain the same beneficial result with less suffering and distress to the patient.

If there be *excessive secretion or hæmorrhage*, astringents may be given; thus, mineral acids, as the sulphuric alone, or with Epsom salts; acetate of lead, tannin, and alum, are also available; and when we have hæmorrhage without great irritability, small doses of turpentine with mucilage or yolk of egg are sometimes prescribed; but this is a remedy in which I have little experience; I have sometimes seen it aggravate the symptoms, besides being very offensive to the patient.

When hæmorrhage has recently taken place, it is well to avoid the use of anything likely to distend or mechanically disturb the stomach, as carbonic acid. Ice, however, tends to produce contraction of bleeding vessels, and should be allowed to the patient.

Pyrosis may be checked by the astringents just mentioned; but we have found the greatest benefit from nitrate or oxide of silver with opium, from creasote, from the compound kino powder, and, when other symptoms would bear it, from the astringent preparations of iron.

The bowels should be acted upon either by agents which are not retained in the stomach, or irritating to it, as the aloes or colocynth pill, with henbane; the effervescing citrate, the carbonate or Dinneford's fluid magnesia; in other instances enemata are useful, consisting of simple water or castor oil, or of turpentine; and occasionally a mercurial purgative will be found beneficial in thoroughly emptying the canal without increasing gastric irritability, as a few grains of grey powder, one or two of calomel, or of blue pill, with henbane, &c.; but to continue this form of medicine is, we think, injurious and prejudicial to the patient. In many cases of constipation with gastric disease, minute doses of strychnia, or of the extract of nux vomica with aloes, act very beneficially.

In order to carry out the *third* indication of treatment, namely, in preventing the extension of the disease, sudden and

violent exertion should be guarded against; and also the distension of the stomach by large meals, or by the formation of gaseous products, as the result of fermentative changes.

4th. In the treatment of the complications of gastric ulcer, arising from its extension to neighbouring parts, as when perforation has taken place, and the symptoms of peritonitis have been suddenly produced, there is still a slight chance that life may be prolonged: the patient should not be moved, nor anything introduced into the stomach, except a teaspoonful of water, or milk, to assuage thirst. Opium must be given freely, as recommended by Dr. Stokes and Dr. Graves, so that the patient may be entirely under its influence—a grain every three or four hours—by this means peristaltic action is checked, nervous shock diminished, extravasation prevented, adhesions promoted, and life may be thus preserved. For many days aperient remedies should be avoided, and food only taken in the most cautious manner.

If local suppuration have taken place, opium is still the best remedy, in order to diminish irritative fever, to relieve pain, and to place the patient in the most favorable condition for reparative action. If the disease have extended into the chest, the prospects of recovery are still less; for sudden acute pleurisy and empyema, or asthenic pneumonia are almost certain to follow. Life may be prolonged by sustaining the patient, and the severity of the symptoms of acute disease of the chest may be partially relieved by ammonia and opium.

The two following cases illustrate the relief that may be afforded by the treatment just recommended.

CASE LXXI.—*Chronic Ulcer of the Stomach. Relieved.*—Jane H—, æt. 34, was admitted under my care into Guy's Hospital, May 1st, 1861. She was a married woman, but had had no family. For eleven years she had had pain at the stomach, with frequent attacks of vomiting of clear fluid. In 1858 she vomited a large quantity of blood, and eighteen months later had a second attack of hæmatemesis. During three months prior to admission, she had continued pain at the scrobiculus cordis, extending to the spine, and increased by food; there was tenderness at the scrobiculus cordis; and she became so weak as to be obliged to keep her bed. For two months she had had vomiting, sometimes *directly* after food, at other times an hour afterwards; the bowels were constipated. On admission she was emaciated, and rather sallow; there was tenderness and increased pulsation at the scrobiculus cordis. She was most free from pain

when lying partly on the back and towards the left side. The pulse was very compressible, 74; the tongue was very red in the centre; the respiration was coarse at the apices of both lungs; menstruation was scanty and irregular, and had disappeared for two months. She was ordered colocynth pill, with henbane at night; and nitrate of bismuth with chloric ether three times a day, and food in a fluid form.

8th.—Symptoms relieved; no vomiting, and less pain. She continued to improve greatly, and left the hospital convalescent on May 27th.

There appeared to be little doubt, in this instance, as to the presence of ulcer at the stomach; and we have very rarely observed a greater measure of relief than she experienced.

CASE LXXII.—*Ulceration (cancerous?) of the Stomach. Relieved.*—David H—, *set.* 36 years, was admitted into Guy's Hospital under my care, April 8th, 1861. He was a married man, a printer, who had resided at St. Luke's. Fifteen months previously he had drunk gin immoderately, and violent vomiting was produced; no blood, however, was ejected. After that time he had suffered from pain across the abdomen and in the back. The pain was greatly increased by food, and vomiting came on directly after it had been taken; but for six weeks the pain had been less severe. His diet had consisted of beef-tea, eggs, milk, &c. He was greatly emaciated, sallow, and had an anxious expression of countenance. A hard tumour, about two inches across, could be felt at the scrobiculus cordis. The action of the heart was feeble, and the pulse was very compressible. Extract of nux vomica, with disulphate of quinine and aloes, and myrrh, were given three times a day.

April 18th.—He complained of intense pain at the scrobiculus cordis, unrelieved by position, and increased by food. There were spots of purpura on the legs. He craved for meat, but refused vegetables. Soap and opium pill were prescribed night and morning; and lemon juice, with infusion of calumba, three times a day.

20th.—Although he had no vomiting, the pain was very severe at the stomach; he was more prostrate and distressed, the bowels were constipated, and he was unable to take solid food. The opium was continued, and he was directed to take chloric ether with bismuth.

22nd.—He was very much relieved.

30th.—Again suffered severe pain. He was in the habit of taking a small quantity of food, and after a short time, because the pain became very severe, he endeavoured to excite vomiting, as affording the only means of relief. He was very much emaciated and prostrate; but he said that he was rather easier when sitting up. Opium was added to his medicine.

May 28th.—Very prostrate, and emaciated to an extreme degree; but he said that he was well, because the pain and vomiting had subsided. The tumour at the scrobiculus cordis was less distinct.

June 8th.—The gastric symptoms continued in abeyance, and he left the hospital in improved health.

The pain produced by the reception of food into the stomach, and increasing till vomiting had taken place, was very characteristic of organic gastric disease. The prostration was excessive; but when almost pulseless, and when the pain and vomiting had subsided, the patient stated that he was well, and he insisted on endeavouring to walk about. It was a good illustration of the benefit of avoiding fresh causes of irritation, and opium with chloric ether afforded great relief; but, although he left Guy's free from pain, we fear the disease was advancing, and would lead to a fatal result.

Sloughing of the Mucous Membrane of the Stomach.—The action of caustic poisons is the ordinary cause of sloughing of the mucous membrane of the stomach; but in the two following cases the appearance was peculiar, and very different from that produced by a clot of blood covering an ulcer. At the lesser curvature of the stomach there were several black patches, the largest being about an inch in length; and other smaller patches were placed in the same direction along the lesser curvature. The black central portion could not be removed from the tissue beneath; but, on section, it was found that a cup had been formed of fibrous tissue surrounding the base and on either side of the slough, showing either that an inflammatory condition had preceded the loss of vitality in this isolated portion of membrane, or that having sloughed, this new action had been set up around it. The appearance presented was very similar to an ordinary bed-sore on the sacrum. A slight, unusual irritation, with depressed vital power, appeared sufficient to cause total loss of vitality. Effusion of blood into the substance of the mucous membrane probably preceded this change, and it closely corresponded with a condition sometimes found in the lung, namely, that observed when hæmorrhage into the substance of the lung is followed by loss of vitality in the part; and one or more lobules of the lung are found detached by an attempt at reparative action.

The cases here recorded, confirm the opinion expressed by Dr. Copland, and are in accordance with the experiments of others, that the condition of the nervous system has an important influence on the mucous membrane of the stomach. In

both cases there was acute pneumonia; in the one, with renal anasarca, in the other, with paraplegia. The effect of division, or disease implicating the pneumogastric nerve, on the nutrition of the lung, is shown by great congestion, and often by acute pneumonia, as we have remarked in the consideration of diseases of the œsophagus; instances also are very frequent of functional disease of the stomach arising from irritation of the pulmonary branches, and cerebral centre of the pneumogastric nerve; but the production of organic change in the stomach by division or disease of the nerve has not been established.

CASE LXXIII.—*Ulceration of the Stomach; Sloughing. Paraplegia. Softening of the Spinal Cord. Disease of the Vertebrae.*—Elizabeth G—, æt. 33, was admitted February 23d, 1855. She had been ill for six weeks with paraplegia. Sloughing of the hips, &c., followed, and she gradually sank.

Inspection thirty-six hours after death.—Opposite the eleventh dorsal vertebra the cord was quite diffuent; and this softening, though in rather less degree, extended to the upper part of the dorsal region. The softening was more marked upon the posterior column. *Chest.*—The bronchi were congested, and were full of tenacious mucus; the lower lobes of the lungs were in a state of red hepatization, being red, fleshy, and very soft. The mitral valve was thickened. *Abdomen.*—Omentum attached to the bladder; the stomach was placed vertically, and was distended; it was pulled down to the pelvis, and occupied half the abdomen. *Stomach.*—Much enlarged, containing grumous fluid; its greater curvature presented post-mortem solution, and the mucous membrane was partially destroyed. Above the line of solution there were several ulcers about the size of a shilling piece. The mucous membrane at the margin of the ulcers was pale and slightly raised, and the floor of the ulcers was covered by a black slough. The intestines were much congested. The liver was very fatty. The spleen was healthy. The bladder was sloughing, as also were the vagina and os uteri, so that there was free communication between them. The uterus contained a decomposing fœtus of about two months.

CASE LXXIV.—*Mottled Kidney. Anasarca. Pneumonia. Sloughing Mucous Membrane of the Stomach.*—Stephen F—, æt. 51, was admitted April 10th, and died April 20th, 1855, from chest disease. Nine years before he had had scarlet fever, and for the last eighteen months he had not been well. On admission the urine was very albuminous.

Inspection fourteen hours after death.—The body was generally anasarcaous. The lower lobe of the left lung was red, consolidated, and almost breaking down. The rest of the lung was very œdematous. The bronchi were full of frothy mucus. The left ventricle was much hypertrophied. The weight of the heart was 17 oz. At the lesser curvature of the stomach were several sloughs; the largest two inches in length, and about one in breadth, black, and slightly raised; a section showed that the slough was situated in a cup

of slightly thickened tissue. Two smaller sloughs were situated near to it. On microscopical examination of the adjoining portions of mucous membrane, the gland follicles were not distinct; and on the surface were columnar epithelium and crystals, &c. The small intestines were healthy. The spleen was small, firm, and lardaceous. The kidneys were mottled, and the Malpighian bodies were degenerated and lardaceous.

Fibroid Degeneration of the Pylorus.—The condition of the pyloric valve in which degeneration of a fibroid character is found to exist, has, by some pathologists, been considered as a form of cancerous disease, by others as hypertrophy of the normal constituents of the affected part. If, however, the diseased structure be carefully examined, no evidence of cancer will be found in it, or in the adjoining parts. The disease apparently commences in the submucous cellular tissue, which undergoes fibrous thickening, whilst the mucous coat is in many cases unacted upon. This fibroid deposit leads to obstruction at the valve; the muscular coat then becomes hypertrophied, and the amount of the hypertrophy is an indication of the degree of obstruction.

The growth beneath the mucous membrane is whitish in colour, firm, and without any juice, as in cancer, sometimes cartilaginous in hardness; it consists of elongated or wavy fibres, resembling a fibroid tumour, and with acetic acid it presents numerous elongated nuclei; bands of similar tissue pass between portions of involuntary muscular fibre; and externally the omentum may be contracted, and adhesions may have been formed with adjoining structures.

The *symptoms* closely resemble those of cancerous obstruction; and they consist in chronic dyspepsia, followed by emaciation, vomiting occurring several hours after food, pain, distension of the stomach, with eructations, fermentation, and the development of *sarcina ventriculi*, constipation and gradual exhaustion, till at last the patient sinks from inanition. The abdominal walls are wasted and collapsed, and a tumour is often felt at the epigastric region, consisting of the thickened tissues at the pylorus. If, however, the stomach be free from adhesions, the thickened pylorus is often pushed downwards, so as to be felt near the umbilicus, or even near to the pubes. Pain is not generally a *marked* symptom of this form of pyloric

disease; but tenderness on pressure is sometimes present, and this, perhaps, arises from peritoneal adhesions.

After death, in some cases, we find evidence of chronic change, and a gray and thickened appearance of the mucous membrane of the stomach, and chronic ulcer or cicatrix are occasionally present. At the pylorus the mucous membrane may be quite healthy, having distinct, or even hypertrophied gastric follicles; but the irritation at the part may have excited secondary disease and ulceration. The glands near the pancreas are not usually affected.

The *diagnosis* is sometimes obscure, and the presence of other more acute disease, as in Case LXXV, may entirely mask the complaint. The patient, in that instance, was emaciated, anæmic, and exhausted from loss of blood, and from disease of the large intestine; vomiting also was absent during his residence in the hospital, nor was it mentioned as a symptom. The duration of life after fibroid degeneration has taken place is greater than in the ordinary forms of cancer, especially medullary and epithelial cancer; and in simple fibroid disease the cachexia has not the expression peculiar to cancerous affections; but it will be often found extremely difficult to distinguish these diseases during life, especially when only observed at their earlier stages.

We are not acquainted with the predisposing, nor with the *exciting causes* of this fibroid disease; but it is probable that long-continued irritation, as indicated by dyspepsia, generally precedes it. The intemperate do not appear to be more liable, and one sex is equally the subject of it as the other; it occurs, also, in early and middle, as well as in advanced life.

As to treatment, we can afford relief, but cannot remove the obstruction. The change from solid and irritating food to fluid and unirritating nutriment is often followed by much benefit; and we may use with advantage those agents and means which have been recommended in chronic ulceration of the stomach.

CASE LXXV.—*Thickened Pylorus. Cicatrix of the Mucous Membrane, with Hypertrophy. Ulceration of the Cæcum and Colon. Fatty Degeneration of the Heart.*—G— G—, a silk weaver, æt. 62, was admitted in a prostrate and

anæmic condition. He had had hemorrhoids for twenty years; and he had occasionally lost a considerable quantity of blood. Four months before admission, he had violent pain from the hip to the foot, and his legs swelled. Violent pain also came on in the region of the stomach. Diarrhœa followed, and continued till death.

On inspection, the heart was found to be fatty, the colon and cæcum ulcerated. The stomach was somewhat enlarged, and its mucous membrane pale; at the greater curvature, for a space of about two inches in circumference, the mucous membrane was thickened, and was a little puckered; and at the upper border of this patch was a small growth, consisting of thickened and prominent mucous membrane, about one eighth of an inch above the remaining part. On examining the raised portion, it was found to consist on the surface of columnar epithelium, and beneath of cell-structure. The nuclei of the cells were very distinct, and gave the idea of cancer, but they were identical with the secreting cells ordinarily observed in a healthy organ. The pylorus was much thickened, and consisted of dense fibrous tissue, passing between bundles of involuntary muscular fibre. There were no true cancerous structures, and the diseased condition of the pylorus arose from fibroid degeneration of the submucous and submuscular tissues, which had been followed by hypertrophy of the muscular coat. See Drawing, in Museum, No. 298⁶⁰, Prep. 1806⁷.

This fibroid degeneration, with hypertrophy, contrasted remarkably with true scirrhus disease. It had not led to the ordinary symptoms of obstructed pylorus, and was not diagnosed during life; the cause of death was exhaustion from diarrhœa in an anæmic subject.

CASE LXXXVI.—*Diseased Pylorus. Phthisis.*—Mary W—, æt. 22, admitted into Guy's Hospital December, 1856. She stated that she had worked at the fur trade, and was nearly always in a stooping posture; three years previously vomiting had come on, preceded by pain across the chest; the symptoms, however, were much relieved, and she married; in a short time she became pregnant, and the symptoms returned; they were, however, regarded as sympathetic from the uterine condition. After her confinement she nursed for seven months; and for four months prior to admission she had constant vomiting, which came on several hours after taking food; she suffered from constipation, and gradually emaciated.

On admission into Guy's she was exceedingly wasted, and had a strumous appearance; her complexion was dark, and she was anæmic; she suffered from flatulent distension, which was easily dispersed, and complained of burning pain at the stomach; the vomiting often came on about six o'clock in the evening; on examination of the abdomen, a prominent tumour could be felt at the region of the pylorus. The vomiting after food and emaciation continued, and medicine afforded very temporary relief; a few weeks before death cough came on and expectoration; she died March 10th; her death had been expected week after week, but still she lingered on, and at last the

emaciation became extreme. A short time before admission she had slight hæmoptysis, but there was no evidence of disease of the chest at that time.

Inspection was made on March 12th. On opening the abdomen scarcely anything but the stomach enormously distended could be seen; the pylorus was somewhat depressed, but the greater curvature reached nearly to the pubes. The tumour consisted of the diseased pylorus. The interior of the stomach presented a growth at the pylorus which completely surrounded the valve, so that the little finger could not pass; it extended nearly two inches into the stomach; the disease was of the character which has been described as hypertrophy, and was manifested in a very marked degree. The semi-transparent muscular layer was more than a quarter of an inch in thickness, and was traversed by delicate lines; upon it was placed a very dense, whitish substance, nearly half an inch in thickness, firm and tough in texture, and could be cut with difficulty; on pressure no juice exuded; upon this, again, was thickened mucous membrane; the surface was not ulcerated, but was whitish in colour, and irregularly tuberculated. The disease did not terminate so abruptly in the duodenum as we often find, but gradually subsided to the natural thickness of the intestine. The duodenum was otherwise healthy; so also the mucous membrane of the rest of the stomach. The intestines, liver, kidney, and glands were healthy.

On microscopical examination, the non-malignant character of the growth was well shown. The mucous membrane at the pylorus covering it was thickened, but presented normal structure; the gastric follicles were beautifully distinct, elongated, and filled with nuclei, apparently quite healthy. The white submucous substance was composed of dense fibre, and with acetic acid presented elongated nuclei, arranged as in fibrous tissue; there was no evidence of cancerous deposit. The muscular tissue had the usual involuntary fibre, but firmer bands intersected it. There was no cancerous disease in any part of the body.

At both apices of the lungs there was disorganization; there were several small vomicæ filled with pus, and surrounded with iron-gray pneumonia, and with some white, granular deposit, resembling tubercles; but no cancerous disease could be found on microscopical examination.

The disease apparently commenced in the submucous cellular tissue, and consisted of abnormal development of the ordinary fibrous tissue, and closely resembled the fibrous growths of other parts. The hypertrophy of the muscular and mucous coats were probably secondary, and were the result of the obstruction.

The history of the case, the disappearance of the symptoms, and their recurrence after an interval of more than two years, are more allied to fibrous degeneration than to cancer.

The occurrence of phthisis with the diseased pylorus is rare; the patient was a strumous subject, and during the exhaustion

consequent on the disease of the stomach pneumonia was set up, and low organized product effused.

The age of the patient was less than that in which we generally find this disease, only twenty-two; and it is doubtful how far her employment induced the complaint.

Polypoid Growth of the Stomach.—The mucous membrane of the stomach not unfrequently presents polypi attached to its surface; several of those which I have examined have presented the appearance of healthy mucous membrane, and they had not produced any symptoms; sometimes smaller growths of this character appear incorporated together, and closely resemble the appearance of commencing carcinoma; it would seem that a cicatrix or some irritating cause has in some cases induced these growths.

Cancer of the Stomach.—The stomach is one of the organs most frequently affected with cancer, and in this frequency a remarkable contrast is presented when compared with the rarity of strumous disease of the same organ. Every form of cancer is found to occur in the stomach, but instances of medullary and scirrhus cancer are the most numerous, whilst epithelial, villous, colloid, and melanoid are rarely found. It is seen, however, that these varieties frequently pass the one into the other, and thus, while one part has almost the firmness and structure of scirrhus, another has the characteristics of medullary growth; and again, the surface also of a medullary cancer may have the appearance of a villous structure. The disease originates in the mucous membrane of the stomach, or its submucous tissue, or it is propagated to the stomach by the affection of the glands in the neighbourhood of the pancreas; and the pylorus, lesser curvature, and cardiac extremity are the parts generally affected. It is not necessary for me to describe the ordinary characters of the several forms of cancer; the cases I have briefly given show the general appearance of the structures found in them. The first case is one in which scirrhus disease was well marked; but cells closely resembling those of epithelial cancer were observed. The second and third cases are medullary; but the surface of the growths had a villous character, which led me to regard them as almost of an intermediate type. The fourth and fifth are cases of col-

loid cancer, the one with similar disease of the ascending colon, the other of the omentum, peritoneum, and rectum; in both the mucous membrane of the stomach was primarily affected.

We are not acquainted with the determining cause of the forms of cancer, or whether the opinion, which is maintained by some pathologists, can be established, that scirrhus is connected, in its origin, with the fibrous tissues of the part; medullary with the mucous surface or gland-structure, and colloid especially with the latter; or whether they are rather indications of the intensity of the morbid action. The part affected has a modifying influence on the character of the disease: the epithelial cancer of a surface covered by squamous epithelium is different from the same disease, where the epithelium is columnar; an instance of differentiation as applied to morbid changes. It would seem that scirrhus disease is less removed from normal nutritive change than medullary cancer; in the one there is a greater disposition to form fibroid tissue, in the other the growth is cellular or even nucleolar.

The *symptoms* of cancerous disease of the stomach, when a tumour cannot be detected on manipulative examination of the abdomen, are often exceedingly obscure, especially in the earlier stages of the disease. It may be convenient to divide the symptoms into three classes, as they are manifested in different stages of the complaint. First, before the formation of any perceptible tumour; secondly, during the development of a growth; and thirdly, the last stage, that of disintegration, by ulceration or sloughing. The first symptoms are those of dyspepsia; and with it there is often a peculiar sallow and anxious expression of the countenance; pain at the stomach may be entirely absent, or there may be severe gastrodynia; pyrosis is frequently present. In the second stage of the disease vomiting is generally the most prominent symptom, especially when the disease is situated at the pylorus or cardia, and the rejection of food takes place according to the seat of obstruction or irritation of the gastric surface, either a short time or several hours after a meal. In some instances the vomiting so quickly follows deglutition as to lead to the supposition of œsophageal disease, as in Case LXXXVIII. The pain also becomes more severe, and is generally of a more lancinating

character, than that experienced in chronic ulcer of the stomach. The vomited matters are often frothy and fermenting, and present us with abundant *sarcina ventriculi*. Hæmatemesis is occasionally present. Flatulence distresses the patient, and eructations are frequent; the bowels become constipated; emaciation steadily advances, and the countenance becomes haggard and cachectic. On careful examination, a tumour may generally be felt at the region of the stomach, or of the pylorus; it often increases rapidly, and, on account of the wasted condition of the parietes, becomes very apparent. In the third stage of the disease the symptoms are more severe and the emaciation extreme, and the vomiting of coffee-ground substance often precedes a fatal termination. The vomiting sometimes ceases, on account of the sloughing of the growth; the obstruction thereby being removed, or the branches of the pneumogastric nerve being destroyed, there may be cessation of consequent irritation; the pain also, sometimes diminishes from similar causes, and as the exhaustion becomes typhoid in its character, the pain may entirely cease, or, as in Case LXXIX, pain may be almost absent throughout the course of the disease. Again, as it has been shown by Dr. Kennedy,* the size of the tumour may actually lessen from the sloughing process.

The immediate cause of the death in cancer of the stomach differs exceedingly; the fatal termination results from exhaustion consequent on the interference with the absorption of nutriment and the completion of the digestive function; or, after sloughing has taken place, the patient rapidly becomes prostrate; the breath is often very offensive; he is seized with hiccough, and, in many instances, the absorption of septic matter takes place, typhoid prostration ensues, and lobular pneumonia is the result; or the ulcerative process may produce, in rare cases, fatal hæmorrhage. Again, the extension of disease to adjoining parts may materially modify the later symptoms of disease, by extension to the liver, by pressure of enlarged glands on the bile-ducts, thus producing jaundice, or ascites follows the implication of the peritoneum. The cancerous ulceration sometimes extends to the actual destruction of adjoining tissues. In Case LXXXIV the transverse colon was

* 'Dublin Quarterly,' 1851.

adherent, and communicated with the stomach by a valvular and sloughy opening; and, if the opening had been enlarged, fæces would, probably, have passed. Case LXXXVIII was of a similar kind. The adhesions and sloughing occasionally reach the external parietes; and, if a communication also exist with the colon, an artificial anus is the result. We have, however, more frequently found this perforation of the skin, in instances of cancerous disease affecting primarily the transverse colon, as after blows. (See Disease of Colon.) In the cases of cancer of the stomach just referred to there was no evidence that fæces passed into the stomach, but merely gas, which greatly distressed the patient by the fæcal odour of the eructation. Dr. Gardner* states that fæcal vomiting is more likely to take place when the pylorus is free; but Dr. Murchison, on the contrary, and we think correctly, remarks, that fæcal vomiting is regulated by the size of the communication between the stomach and colon.

The coffee-ground substance to which we have referred consists of blood which has oozed from the diseased surface, and has become darkened by the action of the gastric juice.

The cancer is generally found to have involved the glands in the small omentum at the lesser curvature. Next in frequency we find the liver also involved, sometimes to a very great extent, so that it is difficult to state in which structure the disease commenced; then the glands in the anterior or posterior mediastinum and tubercles are found on the pleura or in the lungs. And lastly, other abdominal viscera, the peritoneum, kidneys, spleen, contain cancerous growths. In cancerous disease, also, we observe that the coats of the stomach themselves become infiltrated, and in this respect contrast with the condition which we find in fibroid degeneration of the pylorus. In any form of obstructive disease at the pylorus the muscular walls become hypertrophied; but, if there have been ulceration at the pylorus, and the obstruction has subsequently been removed, the hypertrophy may be exceedingly slight; so also when the central portions of the stomach or the cardia are affected.

It will be found that the symptoms of cancer closely resemble those of chronic ulcer of the stomach; both are preceded by a

* 'Edinburgh Medical Journal,' July, 1855.

period of dyspeptic suffering, during which the diagnosis is exceedingly obscure. The expression of the countenance in both is indicative of distress, but in chronic ulcer there is pallor, in cancer cachectic sallowness. Vomiting of blood is more frequently observed in ulceration than in cancer; but in the closing stages of cancerous disease the rejection of coffee-ground substance is of very frequent occurrence. The pain of chronic ulceration is often very intense, even more so sometimes than in cancer; but it is of a gnawing character in the former, more acute and lancinating in the latter; again, the vomiting is often more severe in ulceration than in cancer. The tumour of cancer is generally much larger and more perceptible than the thickening around an ulcer. The emaciation in both may be gradual, progressive, and extreme; but the termination in ulcer is more frequently by hæmorrhage or perforation, whilst in cancer it generally arises from the typhoid exhaustion consequent on the degeneration or sloughing of the growth, the absorption of decomposing material into the blood, or the extension of disease to adjoining structures. Both diseases may occur at the same age, but it is more common to find chronic ulceration at an earlier period than cancer. This is shown by contrasting the ages of the cases we record of chronic ulcer of the stomach with those of cancerous disease of the same organ; the average of the former being 37, and of the latter 52. From 40 to 60 years is the age at which we are most likely to have cancerous disease of other organs, and the law holds good with the stomach. The age will in some measure assist us in the diagnosis even at the later stages, but still more in the earlier; for the varied forms of dyspepsia, gastrodynia, pyrosis, &c., are very frequent at a period long antecedent to the age at which cancer generally manifests itself; dyspepsia being exceedingly common among young females, whilst cancer is almost unknown.

The investigations of Dr. Brinton* on this subject are very interesting and important; he has collected from varied sources a considerable number of cases of cancerous disease of the stomach; and he shows that males are more subject to the disease than females in the proportion of 2 to 1; out of 223 cases, 151 were males, and 72 were females; as to the age of

* 'Med.-Chir. Review.'

those affected, the period does not coincide with those which have come under my own observation as compared with ulceration of the stomach. The following table is also from his researches.

Liability to cancer and ulceration of the stomach.

Age	0	10	20	30	40	50	60	70	80	90
<i>Cancer</i>		$\frac{1}{2}$.	11 $\frac{1}{2}$.	31 $\frac{1}{2}$.	63.	88.	100.	52 $\frac{1}{2}$.	60.	
<i>Ulcer</i>		20.	51.	49.	47.	56.	80.	75.	100.	

The average age of those affected with cancer he mentions to be 51 in the male, and 40 $\frac{1}{2}$ in the female; and in reference to the position, the well-known fact that whilst the pyloric portion is the most frequent seat of cancer, the lesser curvature and posterior surface are the positions of ulcer he shows in the following proportions:

Position.—In 360 cases.

Cancer.—219 Pylorus. 38 Lesser Curv. 36 Cardia. 13 Stomach generally. 11 Greater Curv. 11 Posterior Surface. 11 Anterior Surface. 4 Middle.

Ulcer.—52 Pylorus. 98 Lesser Curv. 5 Cardia. 0 Stomach generally. 8 Greater Curv. 177 Posterior Surface. 18 Anterior Surface. 0 Middle.

As to the duration of chronic ulcer compared with cancer, the former disease extends over a longer period of time; in cancer the duration may be from three to six to twelve months, or even two years; but, in ulceration, the disease will be found continuing three, four, or even seven years, with varied accessions of severe symptoms; and instances have occurred in which twenty years have intervened between the commencement of the symptoms and their fatal termination. Moreover, ulceration is more amenable to treatment.

The evidence of cancer is most marked when the pylorus is affected, and obstructive disease set up. Where this is not the case, cancer is sometimes, however, found after death, without having led to any marked symptom, the patient having died from another disease; the third case is an illustration of this fact, for the symptoms of ascites and cirrhosis obscured those of cancer of the stomach. In Case LXXX the regurgitation of

glairy, gelatinous fluid, and gradual emaciation, were the most marked symptoms. It sometimes happens that cancerous disease of the liver is followed by infiltration of glands at the head of the pancreas, which become united to the pylorus; and, without having infiltrated the mucous membrane, these glands lead to obstruction at this part, causing hypertrophy of the muscular coat, and, by this obstruction, they stimulate primary cancer of the stomach itself.

When the cardiac extremity is diseased, the vomiting frequently occurs so immediately after taking food that the symptoms resemble cancerous disease, or obstruction of the œsophagus.

In some instances the pneumogastric nerves may be traced through the medullary tumours of the stomach; and either the nerve-fibres may be found to present their ordinary microscopical appearance or be entirely destroyed. It is this destruction of the nerve-fibres which sometimes leads to a cessation of pain and of the extreme irritability of the stomach.

In the *treatment* of cancer of the stomach the same remedies which have been mentioned in chronic ulceration may afford great comfort to the patient, although they are ineffectual to the cure; but it will sometimes be found that life is prolonged and ease obtained by administering nutrient enemata. The distressing flatulence and fermentation and vomiting of food are thus avoided, and the sufferer is nourished by this imperfect means, more than by ineffectual attempts to induce normal digestion.

CASE LXXXVII.—Scirrhus Pylorus. Carcinomatous Tubercles in the Liver, Spleen, and Kidney, and on the Diaphragm.—Edgar C—, æt. 40, was admitted January, 1855. He was a cooper, and till the attack, for which he applied to the hospital, he had enjoyed good health. Four months previously sickness had come on, and it took place generally a few hours after taking food; but sometimes he was able to retain three or four meals in succession. A tumour could be felt at the region of the pylorus; there was great emaciation, and he slowly sank.

Inspection twenty-six hours after death.—The thoracic viscera were quite free from disease. There were several white, firm tubercles in the abdomen, about the size of peas, on the under surface of the diaphragm, opposed to the liver; similar tubercles were found on the sheath of the right kidney, and a rather larger one on the surface of the spleen. In the liver, on its under surface, were several tubera, about half an inch in diameter, with

raised and well-defined edges; the remaining portion of the viscus was healthy. The stomach was very much distended with air and dark, reddish-coloured fluid; at its lesser curvature a small tubercle was observed on the peritoneal surface; several of the nerves at the lesser curvature were involved in this growth. On opening the stomach, it was found to contain fluid, as before mentioned, smelling very strongly of lactic acid. At the pylorus was found a hard mass, composed principally of glands, and on the inferior surface the pylorus itself was infiltrated with dense cancerous deposit. The valve was contracted, so as only to admit a large sized catheter, and its mucous membrane was destroyed by ulceration; the ulcer extended into the stomach; its edges were raised, and in some parts were vascular. The muscular coat could be traced nearly to the pylorus, somewhat thickened, but in a healthy condition; it then became involved in the cancerous infiltration, and was of a whitish colour; at the pylorus both muscular and mucous coats were destroyed, and semi-cartilaginous tissue only remained for about three quarters of an inch. The mucous membrane and the infiltrated tissue presented well-marked cancer-cells, with very large nuclei, and aggregated cells, as in epithelial cancer; the dense tissue beneath was gland-tissue, infiltrated with scirrhus product. The duodenum contained bilious matter and a considerable number of white grains, which were at first supposed to be Brunner's glands, but they were found to consist of solitary glands. The pancreas and the remaining portion of the intestine were healthy; there were some infiltrated glands at the commencement of the rectum, but the mucous membrane was sound.

The symptoms in this case were well marked, and it was evident that there was obstructive disease at the pylorus. The examination of the growth at that part showed great resemblance to epithelial cancer; the glands, however, in the neighbourhood of the pancreas, which were infiltrated, and the cancerous tubera found in the liver and on the peritoneum, presented the characters of ordinary scirrhus.

It was an interesting fact to find at the rectum, a frequent seat of cancer, the glands infiltrated; but the lumbar and mesenteric glands were free from disease.

CASE LXXVIII.—*Medullary Cancer of the Stomach, having a villous character.*—Thomas G—, æt. 62, was admitted December 13th, 1854. He had been a shepherd at Shoreham, and eight months before admission experienced flatulence, loss of appetite, and dyspepsia. For six weeks he had been very ill, and he had suffered occasionally from vomiting. He had no pain or uneasiness at the stomach; but he was emaciated, and there was a tumour, about the size of an orange, situated just above the umbilicus, but separable from the liver, and slightly moveable on respiration. Slight œdema of the ancles came on before death, which took place in January, 1855.

Inspection seventeen hours after death.—There was slight arcus senilis. The thoracic viscera were healthy. On opening the abdomen, a tumour, about the size of an orange, was found to be situated at the pyloric end of the stomach; the gall-bladder above was adherent to it, accounting for the movements of the tumour with the liver; and below, the transverse colon and omentum were inseparably united with it. The pylorus appeared embraced by the growth extending from above and below, and on opening the stomach the whole of its circumference was found affected. The intestines were collapsed; the liver was healthy, but its peritoneal coat was thickened at its lower margin. The gall-bladder was empty, the pancreas not at all affected, though in close contact with the tumour; the kidneys were small, atrophied, and contained cysts. There were several gastric glands in the neighbourhood of the lesser omentum, which were infiltrated with cancer, but the lumbar and bronchial glands were not affected. On opening the stomach, it presented a large medullary growth, extending about two inches from the pylorus into the stomach, involving the whole of the valve, and forming a projecting, soft, tubercular ring, vascular, and extending into the duodenum. The pylorus itself would admit the tip of the little finger. This growth was soft, of a yellowish-white colour, and about one inch in thickness. At the margin the muscular coat could be traced into it, forming a semi-transparent layer, about a quarter of an inch in thickness, but evidently infiltrated with cancer; at the edge of the cancer the muscular coat suddenly became of its usual thickness, showing that there had not been great obstruction, so as to lead to much hypertrophy of that layer. Near to the lesser curvature was another growth, projecting from the mucous membrane, soft, irregular on its surface, and covering about a square inch in extent; it was about half an inch in thickness, and at its edges presented small, soft, tubercular growths, projecting from the membrane. The mucous coat, involved in carcinomatous disease, could be dissected off, away from the muscular, till near the centre of the growth, where the whole were firmly united together, and large vessels could be seen passing into the cancerous mass. Near this part, vessels full of blood passed to its circumference, giving it, in some parts, a red and vascular appearance.

Microscopic examination showed that the mass consisted of cells and nuclei, varying in size; some cells were about the size of healthy epithelium. The nuclei were large, very distinct, and some had double nucleoli. On taking a portion of the surface of the tumour, and floating it in water, numerous rod-like processes were observed, extending for a considerable distance from it, having the character of villi; and they gave to the margin of the growth a flocculent appearance. These villi were found to contain numerous nuclei.

At the margin of the growth the gastric follicles were much degenerated, and they were in some parts distended, but without

cells; in other parts, only the termination of the follicles could be seen; again, some of the follicles had an irregular outline, and presented crystals on the surface of the membrane. Around the former portion of atrophied follicles there was fibrous tissue, arranged in meshes, and with acetic acid the fibres appeared minutely granular.

The whole appearance of this structure was that of medullary cancer; it was composed principally of nuclei, and had affected the pyloric extremity, leading to symptoms of obstruction. There was some infiltration of the adjoining glands; but the remaining viscera were healthy. The growth appeared to have commenced in the mucous membrane. It was on the examination of the surface, however, that the resemblance to villous cancer was manifested; the surface had a flocculent appearance, and microscopical examination showed that this arose from villous processes extending from the surface.

This case appeared to stand in an intermediate position between medullary and villous cancer; and it confirms the opinion expressed by Mr. Paget, that the latter may be merely a variety of the more common form.

As to the symptoms, several months of dyspepsia were passed; the health then rapidly failed, and prostration of strength, emaciation, and occasional vomiting, were the principal indications of disease. It was remarkable to observe the absence of pain or even uneasiness at the stomach.

CASE LXXIX.—*Villous Cancer of the Stomach. Cirrhosis. Ascites.*—Isabella D—, æt. 65, admitted July 11th, 1855, and died July 26th. She was a married woman, who had been accustomed to take spirits, but she stated that she had been in good health till the previous January, when she caught cold, which was succeeded by cough, by shortness of breath, and by burning pain at the scrobiculus cordis. Seven weeks before admission, her legs, and afterwards the abdomen, began to swell; diarrhoea, great prostration, and syncope came on, and before death she became partially comatose.

Inspection nineteen hours after death.—The mitral and aortic valves were opaque. There were adhesions between the liver, colon, and stomach; and the peritoneum contained about a gallon of serum. The liver was in a state of advanced cirrhosis. The stomach was moderate in size, flaccid, and on the inner aspect of its anterior wall presented a large villous growth, about three inches in diameter, the edges of which were raised, and the centre ulcerated. On floating in water, it presented beautiful villous processes; these, under the microscope, were found to consist of long, delicate growths,

some terminating in points, and filled with granules. The base of the growth presented nuclei. There was no hypertrophy of the pylorus, nor of any portion of the muscular coat. The other portions of the mucous membrane presented gastric follicles, containing fat and nuclei. The kidneys were atrophied.

In this case, with the exception of burning pain at the stomach two months before death, which is a not unfrequent symptom of dyspepsia, there was no sign observed of disease of the stomach. This is partly explained by the disease affecting the anterior surface of the organ, leaving the pylorus perfectly free. This absence of obstruction was further shown by the atrophic rather than hypertrophic condition of the muscular coat. The advanced disease of the liver producing dropsy, appeared sufficient to explain all the symptoms, and the distension of the stomach entirely prevented any tumour being felt at that region. The appearance of the growth, under the microscope, gave less positive proof of a cancerous origin than in the preceding case; some granules at the base of the villi, but none of the ordinary cancer-cells or nuclei, were present.

CASE LXXX.—*Colloid Cancer of Stomach and of the Colon*.—Elizabeth T—, æt. 37, was admitted July, 1850. She had been a servant, and had been out of health for four months, but twelve months previous to admission she had had jaundice. She was somewhat emaciated, and had a sallow, aged, and very haggard expression of countenance. She complained much of flatulent distension of the abdomen, with a sensation of sinking at the scrobiculus cordis; after eating she suffered much pain at the stomach, but this pain was most severe after taking fluids. There was occasional vomiting, or rather regurgitation of thin, glairy, and gelatinous fluid; this came up into the throat, especially at night. The bowels were constipated, and she was troubled with hæmorrhoids. The abdomen was moderately distended, but no tumour could be felt on manipulation. The pulse was feeble. She was in a semi-jaundiced and drowsy condition, complained of a sense of fulness in the head, and of muscæ volitantes; she became more and more exhausted, and gradually sank.

Inspection was made August 5th. The body was not extremely emaciated. The intestines were much distended with flatus, and the peritoneal sac contained several pints of fluid. The stomach was very much contracted, and its walls were three fourths of an inch in thickness. The outer or muscular layer was a quarter of an inch in thickness, semi-transparent, and it was divided by white bands which were continuous with the submucous tissue. The mucous membrane had a pulpy, honey-comb appearance, and it was replaced by minute colloid cysts, containing clear, gelatinous fluid; these cysts were most distinctly observed on the internal surface of the stomach; there

were also some ulceration and congestion of the vessels. The pylorus was not thicker than the rest of the stomach; but the hypertrophy of the muscular coat extended the whole length of the œsophagus. Some of the glands of the curvature of the stomach were hard and thickened. The fluid from the colloid cysts contained large cells filled with several nuclei, and were surrounded by very delicate areolar tissue. The vessels of the stomach were rendered quite patulous by the tissue placed around them. The small intestines were free, but the large intestine was much thickened; immediately above the cæcum there was a portion of colon affected with colloid growth, and from the hypertrophy of the muscular coat it had the appearance of a pyloric valve; the submucous coat also was much thickened. Some of the solitary glands in the colon were enlarged. The liver, kidneys, and spleen were healthy; so also the thoracic viscera. The heart was contracted.

In this case the symptoms at first were not at all more severe than those often observed in pyrosis, with flatulent distension of the abdomen; nor was the serious nature of the disease for some time anticipated.

The stomach is preserved in the Museum, No. 1813²⁹, and shows in a very beautiful manner the structure of colloid cancer. The hypertrophy of the muscular coat was remarkably extensive, reaching into and passing along the whole length of the œsophagus. The small intestine was free; but the mucous membrane of the colon was affected with similar disease to that of the stomach. Of this there was no evidence during life, although the constipation of the bowels was, perhaps, rather more obstinate than in cases of ordinary cancer of the stomach; but not more than is observed in many cases of dyspepsia. The most marked symptom was the regurgitation, and the filling of the mouth during sleep with watery, gelatinous fluid; this, unfortunately, was not examined microscopically during life; it might have afforded clear evidence of the nature of the disease. The semi-jaundiced condition arose from slight pressure by diseased glands on the common bile-duct, and the colloid disease gradually extended through the whole of the mucous membrane of the stomach by continuity of structure. It would seem probable that it originated in the glandular structure of the stomach rather than in the submucous areolar tissue, the ordinary seat of carcinoma medullare.

CASE LXXXI.—*Colloid Cancer of the Stomach, the Omentum, the Peritoneum, and of the Rectum.*—John C—, æt. 47, was a pensioner, and one month before

his admission began to experience pain at the scrobiculus cordis. Vomiting came on, with costiveness and gradual emaciation. A tumour could be felt extending across the abdomen, and it was doubtful whether this was the margin of the liver, or a tumour involving the pylorus, or merely thickened omentum. He died September 4th, and was examined thirty-one hours after death.

The parietes of the abdomen were very thin. The peritoneal cavity contained several gallons of fluid, of very deep colour, almost sanguineous; the serum presented shreds of lymph, and other delicate bands of lymph passed between the coils of the intestine. The omentum was found to be contracted into a thick, yellowish mass, about half an inch in breadth, which projected towards the abdominal parietes. The margin was irregularly notched, and situated immediately above the transverse colon. The surface of the liver was roughened by small gelatinous tubercles, and a thick layer covered the whole opposed surface of the diaphragm, which at this part was much thickened, and the pleural surface was also encroached upon. The lesser omentum was also much thickened, and a white, hard mass, about the size of a hen's egg, was situated at the lesser curvature of the stomach, near the pylorus. The small intestines were contracted, the large were distended; the peritoneal surface was everywhere studded with small tubercles, from the size of a millet-seed to that of a bean; these were soft, gelatinous, and of a red colour. The sac of the lesser omentum contained tubercles similar to those in the general cavity of the peritoneum, and it was distended with fluid. The cavity of the stomach was small, its parietes were thickened; and at the lesser curvature from the œsophagus to the pylorus the mucous membrane was irregularly raised, and presented an appearance of cells distended with clear, gelatinous fluid. The larger curvature was healthy; the liver was small, and of a deep, bilious colour; the hepatic cells contained very little fat. The pancreas and the small and large intestines were healthy, but at the commencement of the rectum was a small nodule of cancerous growth; this had led to thickening of the mucous and muscular coat, and the intestine at that part would scarcely admit the index-finger. Preparation No. 1813⁹⁰.

The microscopical examination showed well-marked characters of colloid cancer. The growths on the peritoneum consisted of large, compound, nucleated cells, and of delicate, intervening, fibrous tissue. In the omentum there was a greater quantity of fibrous tissue between the cells; and some of the cells contained four or five large nuclei, which were rendered very distinct by acetic acid. The mucous membrane of the stomach presented similar structural elements. The affection of the rectum in this case was an interesting association of disease.

CASE LXXXII.—*Medullary Cancer of the Stomach, of the Liver, and of the Lungs, &c.*—William C—, æt. 60, admitted into Guy's Hospital February 12th, and died March 7th. He was a patten-maker at Woolwich, and of

temperate habits. Three months before his death he began to experience pain at the scrobiculus cordis and in the loins, but had no vomiting till a short time before his death, when he brought up coffee-ground substance. On admission he was feeble, emaciated, and anæmic; his pallor, however, increased, and it was evident that he was rapidly sinking. At the right hypochondriac and epigastric regions a large tumour could be felt.

Inspection was made eleven hours after death. On removing the liver and bringing the stomach into view a large cluster of malignant glands were seen at the lesser curvature, one being the size of a hen's egg. From the great amount of disease in the lymphatic glands and the enlargement of the liver, the cancerous tubera in the two parts came into contact. When the stomach was opened a large tumour was found within it, occupying the lesser curvature. The tumour was sloughing; it was of a greenish-brown colour, and very offensive, and its tissue was broken down; it was very vascular, and it had no doubt been the source of hæmorrhage. The growth in the stomach was nearer to the œsophageal opening than the pylorus, and it occupied about half the length of the lesser curvature. The stomach was elsewhere healthy, and contained a light-brownish fluid. At the seat of the growth the walls of the stomach were beginning to slough.

The liver was much enlarged by carcinomatous growths throughout its substance, the tubera being soft, red, and very vascular. The intestines and other abdominal organs were healthy. There were numerous small cancerous nodules scattered throughout the lungs; they were soft, red, and vascular; the largest about half an inch in diameter.

In this case there was no obstruction at the pylorus, and hence an absence of some of the more common signs of cancer, as vomiting after food, &c. The pallor and exhaustion, with evidence of an abnormal growth at the stomach and liver, were the more marked signs, and it was shown after death that the exceedingly vascular character of the cancer had led to the oozing of blood and to the anæmic condition of the patient.

CASE LXXXIII.—*Chronic Ulceration of the Stomach—Cancer.*—James T—, æt. 46, a weaver, who had been living at Spitalfields, was a regular and sober man, but he had been a great smoker. His father and mother both died of phthisis; for thirty-four years he had been employed at the loom, and he had suffered much from the shuttle striking the scrobiculus cordis; these blows at first produced nausea and faintness, which continued for several hours. Five years before admission the same unpleasant symptoms returned, obliging him to discontinue his work; they were accompanied with vomiting, although at first only his breakfast was rejected; these symptoms continued for four years, and then left him for three months, during which time he rapidly gained flesh, and continued his employment.

Six months prior to his appearance at Guy's he was again attacked with pain and vomiting, and he began to lose flesh; he suffered great pain if he

fasted, but on taking food the pain very soon returned, and it was only relieved by vomiting; the vomiting sometimes came on immediately after a meal, or it was delayed for about six hours; he had never vomited blood; the bowels were constipated, and the urine scanty.

He was a small man, of light complexion, and had a diabetic appearance. The chest was healthy; the tongue was moist and clean; the abdomen soft, flattened, and contracted; the integuments dry. Fluid magnesia ʒss , and dilute hydrocyanic acid ʒiij , were ordered three times a day; and a soap enema.

The vomiting continued very severe, and he became increasingly prostrate; hiccough came on, coffee-ground vomiting, and death.

On inspection, the body was extremely emaciated; the lungs were collapsed; much black pigment was found upon them, but they were otherwise healthy. The heart was healthy. *Abdomen*.—The intestines were collapsed; at the duodenum there was much contraction from puckering of the omentum and stomach. *Stomach*.—The walls were exceedingly thin and atrophied; about two inches from the pylorus was a contraction, which at first was mistaken for the pylorus; there was considerable contraction also of the omentum at that part, and firm semi-cartilaginous hardness of the structure. On opening the stomach, an oval ulcer, about two and a half inches in length, and one in breadth, was observed surrounding the constriction; its edges were rounded and elevated; its base quite smooth. On section, the mucous membrane appeared to be continuous with the upper layer of the ulcer; its deeper layers were very firm, white, and fibrous; beyond the ulcer and its contraction, was a portion of healthy mucous membrane, then the pylorus, which was also perfectly healthy. The first part of the duodenum was congested, and there was pigment in the mucous membrane; in the omentum were several hard tumours, and the omentum itself formed a firm, contracted mass, about the size of the middle finger. On section, these structures were firm, and contained whitish juice, and under the microscope showed large cells containing large, very distinct nuclei, evidently cancerous. In the stomach no follicles could be detected on the smooth surface of the ulcer; and in the structures beneath, none of the cancerous cells found in the omentum and glands were present, but abundant fibrous tissue; there was also much fibrous tissue in the omentum, &c. The rest of the intestine was healthy, the colon was contracted, and contained some scybala. The liver was healthy; the spleen was enlarged and firm; the kidneys also were healthy.

The history and appearances after death in this case warranted the belief that ulceration of the stomach had existed for a considerable time; and although we found evidence of cancerous tubercles in the omentum, I think it probable that the development of cancerous growth only took place during the latter stage of the disease; the growth closely resembled chronic ulcer in its general and microscopical ap-

pearance, except that it nearly surrounded the pyloric extremity. In some instances, cancerous deposit takes place at the edges of a chronic ulcer; such, however, was not the case here.

CASE LXXXIV.—*Cancer of the Stomach. Communication with the Colon. Ulceration of the Cæcum and Ileum. Chronic Phthisis.*—John T—, æt. 67, was admitted August 15th, 1855, he was a married man, who had resided at Greenwich; he had been out of health for twelve months, complaining of dyspepsia, and pain at the scrobiculus cordis; the food appeared to remain at the end of the œsophagus, and not to reach the stomach; there had been no vomiting either before or after admission, but a hard defined growth could be felt at the scrobiculus cordis, which left no doubt as to the nature of the complaint; the abdomen was collapsed. On September 15th, he was greatly emaciated, and able to take but very little food; his mind wandered much; the feet and hands were œdematous; and numerous spots of purpura were found on the hands and forearms. He gradually sank.

Inspection.—The body was much emaciated. *Chest.*—Very strong pleuritic adhesions were found, especially on the right apex; the right lung was puckered, exceedingly dense, and on section presented iron gray consolidation, occupying nearly the whole upper lobe; in the centre it was firm and calcareous; the lower portion of the upper lobe contained numerous miliary tubercles, some surrounded by dense, others with crepitant lung; at the lower lobe were scattered isolated miliary tubercles, semi-transparent in colour. On the surface of the lobe were several lobules which were broken down in the centre into thin pus, and surrounded by a tolerably defined margin; the extreme edge of the lung was emphysematous; the left lung was in a similar condition; but the bronchial glands were healthy. The tubercles in the lungs consisted of molecular matter, of small irregular cells and nuclei; some of the cells were the size of the ordinary ones in the pulmonary structure, but none were like those in the stomach.

The abdomen was collapsed; the stomach was firmly adherent to the transverse colon. On opening the former, along the lesser curvature, a large growth nearly four inches in circumference was found at the pyloric extremity, involving the whole of the pylorus, and surrounding the stomach at that part; the edge was thick, rounded, and raised an inch above the surrounding mucous membrane, so that the growth formed a sort of cup; the margin was of a deep purplish hue; the centre presented an irregular ragged slough of a brown colour; it was deeply excavated, and had a feculent odour. At the pylorus the muscular coat was about four lines in thickness, of a whitish colour, with small intersecting semi-transparent bands. Nearer to the cardiac extremity were two small raised growths, one about half an inch in diameter, red and prominent; the other about a quarter of an inch in diameter. The rest of the mucous membrane was pale. The stomach near the pylorus was firmly adherent to the transverse colon, and from the centre of the slough a probe could be passed into the colon; the opening in the colon was valvular, gray, and about a line in diameter. The

pancreas and omentum were healthy; several mesenteric glands in the neighbourhood were infiltrated with soft cancerous product. The duodenum was gray, but its mucous membrane healthy; at the ilio-cæcal valve were the remains of an ulcer occupying nearly the whole of the last Peyer's gland, and extending to the cæcum; its margin was raised, and presented several congested nodules. Externally the cellular coat was firm, hard, and contracted; the remaining part of the intestines was healthy.

On examining the stomach with the microscope, the growth presented on the surface columnar epithelium, and consisted of nucleated cells, with very large and distinct single or double nuclei, and of delicate intervening fibrous tissue; there was no doubt of their cancerous character. The adjoining mucous membrane presented numerous fat particles in the follicles; the glands contained similar cancerous nuclei. The ulcer in the ileum and cæcum appeared partially cicatrized, it did not present any cancerous product, but only fibrous tissue. At the pylorus, bands of involuntary muscular fibre were found to extend between the cancerous elements.

- As far as it could be decided by microscopical examination, the disease in the lung was of a non-cancerous character; it appeared to consist in a chronic and almost quiescent state of phthisis; but besides this chronic disease, there was evidence of acute lobular pneumonia, which had probably come on a short time before death. The condition of the ileum was that of a healing ulcer. It was difficult to obtain a full history from the patient, and the evident cancer of the stomach obscured the signs of pulmonary disease. The existence of a communication with the colon was not known during life.

CASE LXXXV.—*Cancer of the Stomach*.—Martin F—, æt. 65, was admitted into Guy's Hospital, August 8th, 1856, and died November 23rd. He suffered from the ordinary symptoms of cancerous disease of the stomach, and gradually sank.

Inspection was made thirty-two hours after death. On opening the abdomen, the intestines and stomach were found to be collapsed; at the lesser curvature of the stomach were several hard glands, some adherent to the stomach, others to the pylorus; some of these were softening in the centre. On opening the stomach, the pylorus was found to be very much contracted, its walls of a yellowish-white colour, and soft consistency, replacing the ordinary muscular and mucous coats. The growth was evidently cancerous. The muscular coat was not hypertrophied as we generally find in obstruction at the pylorus, showing that the passage had remained tolerably free. Near the lesser curvature one of the tumours was beginning to protrude.

At the upper part of the œsophagus the mucous membrane appeared as if it were affected with cancer, but this was not the case; numerous oval

slightly raised spaces were of a whitish colour, and firm in consistency, but they did not contain the elements of cancer; the apices of the lungs were adherent, and presented some gray induration at those parts; other tissues were healthy, except that the liver contained two small cancerous masses.

I have introduced this case to show by the inspection after death, that the hypertrophy of the muscular coat is sometimes very slight, although cancerous disease exist at the pylorus.

CASE LXXXVI.—*Struma. Cancer of the Stomach.*—Hannah W—, æt. 33, was admitted, under Dr. Rees' care, October, 1857. She was a thin, emaciated young woman, pale, with a dejected and somewhat melancholic expression; she had always been delicate, and had been subject for a long time to vomiting. For twelve months she had suffered constantly from this symptom, which generally came on in the evening; menstruation had also ceased during that period; but there was no evidence of disease of the ovary. *January, 1858.*—The abdomen was moderately rounded, and there was considerable tenderness at the scrobiculus cordis; and at the region of the greater curvature towards the pylorus, a considerable tumour or induration could be felt. At the angle of the jaw, on the left side, there was enlargement of the glands of the neck; no disease could be detected in the chest, but the respiration was very coarse at the apices. The heart was normal. Various remedies had been tried to relieve the irritability of the stomach, and magnesia mixture, creasote, calomel, &c., had been given. Steel pill and some lemon-juice were occasionally taken, but afforded only slight and temporary relief. Bismuth, and afterwards the injection of nutrient enemata were then tried, but without relief; solid food produced excessive pain, and fluids had the same effect, although in less degree. *February 1st.*—She was unable to bear the injection; and complained much of pain. The tumour was more distinct, oblong, hard, and tender; the pulse compressible, the face flushed. One drachm of fluid pepsine with mucilage was tried, with a small quantity of chop, and milk diet night and morning. *4th.*—She was rather more comfortable. *6th.*—She complained of nausea after the medicine. Opium, 1 grain, was given every night. *April 8th.*—Continued in the same anæmic condition, but the emaciation became greater; the abdomen was collapsed; the tumour in the lower part of the epigastric region was distinct, but not enlarged. There was greater prostration. The pain and vomiting remained as before. She gradually sank, and died July 9th, 1858.

Inspection nine hours after death.—There was consolidation of the lower lobe of the left lung. The bronchial glands were enlarged, and filled with firm and cheesy deposit (strumous). The glands on the left side of the neck were enlarged from strumous deposit. The heart and pericardium were healthy. The peritoneum was healthy. The stomach was reduced in size; the walls were infiltrated with cancer, extending from the pylorus to the œsophagus, along the lesser curvature, and, in some parts, were three quarters of an inch in thickness; whitish, firm, and consisting of cancerous cellu-

lar deposit. The muscular coat was much hypertrophied. The lesser curvature was adherent to the pancreas, and the neighbouring glands were infiltrated. The mesenteric glands contained white softened deposit. The liver presented vascular nodules of cancerous deposit, their central parts being depressed; some nodules were firmer, and white in colour. The right ovary was enlarged.

In this case the association of sturmoid disease of the glands of the neck and of the chest, with cancerous disease of the stomach, was unusual. The patient was younger than those usually affected with cancer, and the duration of the disease, which lasted two years, was greater than in ordinary cases. The pain was severe throughout the disease, and this suffering was, perhaps, due to the infiltration of the glands about the ganglia, or to the implication of branches of the pneumogastric nerve in the diseased walls of the stomach.

CASE LXXXVII.—*Cancer of the Stomach. Disease of the Supra-Renal Capsule.*—John S—, æt. 42, a sailor, was admitted into Guy's Hospital, under Dr. Barlow's care, December 21st, 1859. Six years before he had had dysentery in the Black Sea. In November, 1859, he was taken ill somewhat suddenly at Malta, he fell down on deck, and suffered from severe pain at the scrobiculus cordis. He remained in hospital at Malta for a month; vomiting then came on, and afterwards continued; but he had occasional intervals of cessation of several days' duration. On admission he was emaciated, and had a haggard, distressed appearance; he had severe pain across the abdomen. In January, 1860, severe vomiting returned, generally after every meal, or at night; the bowels were much constipated; he was prostrate, and on placing the hand on the abdomen a hard growth could be felt in the region of the pylorus and gall-bladder. The forehead was discoloured, as in Melasma Addisonii. He slowly sank, and died February, 1860. There was inspection on the 20th, and a cancerous ulcer, about three inches in diameter, was found at the pylorus; its edges were slightly raised, and its surface sloughy; near to the liver was some infiltration of medullary matter. The supra-renal capsule was enlarged and infiltrated. No other part was diseased.

The symptoms in this case were marked, but the sudden onset of the disease, causing him to fall whilst walking and carrying a tray on deck, was unusual, and might have led to an incorrect diagnosis. The patient rapidly became very prostrate, and it is probable that the affection of the supra-renal capsules tended to increase the exhaustion, and hastened the fatal termination.

CASE LXXXVIII.—*Cancer at the Pylorus, simulating disease of the Œsophagus.*—James E—, æt. 61, was under my care in Guy's Hospital in 1859. He was a hawker, who had resided at Kensington; his habits of life had been intemperate. He had been in St. George's Hospital with fractured ribs; hæmoptysis followed, and from 1852 he had suffered from bronchitis. In 1857 he was in the Consumption Hospital for bronchitis; but at the same time he had dyspepsia, with occasional nausea. In October, 1858, he first noticed a small hard swelling in the left hypochondriac region; it was moveable, and did not cause any pain. After that time he gradually emaciated, and had daily increase of pain at the affected part; and vomiting, previously seldom, became almost constant, although it occasionally ceased for several days. On admission, on March 19th, he had a sallow, slightly jaundiced, appearance; he was of dark complexion; his skin was of normal temperature; the tongue was furred, and brown in the centre, but whitish at the edges. The chest presented the signs of old bronchitis; the heart-sounds were normal, but the pulse was feeble, 75. The abdomen was tense and resisting, and was resonant, excepting at the epigastric and left hypochondriac regions, where a defined tumour was present; the tumour was painful on pressure, and had slight pulsation, and on taking solids they were *instantly rejected*, as if there were obstruction at the extremity of the œsophagus; but fluids were retained for three or four hours, and were then rejected as a chymefied mass. There was interscapular pain. The bowels were much confined; the urine was high coloured, of sp. gr. 1020, and free from albumen and sugar. On April 14th, he vomited dark offensive matter, of almost faecal odour. Local erysipelas of the forehead and right eyelid came on, and slight abscess followed. This, however, subsided; but he gradually sank, and died June 19th. The abdomen only could be examined. There was a tumour at the pylorus, about the size of a closed fist, firmly adherent to the anterior abdominal parietes and to the edge of the liver. On removing it from the parietes, a small abscess was found to have been produced anteriorly from the extension of the cancer to the surface, and the consequent local peritoneal inflammation. On opening the stomach, the pyloric end was found to be occupied by a large cancerous growth, which extended about three inches into the stomach and surrounded the orifice, and it reached nearly to the œsophagenal opening at the lesser curvature. The growth was sprouting and fungating; it had a green, sloughy appearance, and was situated in the mucous and submucous tissues. Posteriorly the muscular coat was hypertrophied, but anteriorly this coat was invaded by the cancerous disease, and there was adhesion to the anterior abdominal parietes. There was adhesion also with the colon, and a communication existed, through which a probe could be passed. The pancreatic glands were involved, but the other abdominal viscera were healthy. The growth was soft, vascular, and of an encephaloid character.

The vomiting, in this instance, during the early stage of the disease, took place so quickly after deglutition, that the disease was at first referred to the œsophagus, and, till nearly the close

of life, the cardiac extremity rather than the pylorus was supposed to be the seat of organic mischief. The communication with the colon gave a faecal odour to the ejected matters; and the extension of disease to the skin would, in a few days, if life had been prolonged, have led to the formation of an artificial anus.

CASE LXXXIX.—*Cancer of the Pylorus; Hydatid disease of the Cellular Tissue of the Bladder.*—W. A—, æt. 52, a clerk, who had resided at Woolwich, was admitted into Guy's Hospital under my care, March 28th, 1860. He had enjoyed good health till eight months previously, when loss of appetite and vomiting after food came on; the rejection of food took place either at once or after long intervals. The bowels were constipated. Emaciation had gradually become extreme, and when brought to the hospital it was thought that he would scarcely reach the ward. He, however, rallied, and survived for three weeks. Vomiting did not recur till two days before death. There were no signs of disease of the chest. The abdomen was much contracted; an ill-defined tumour could be felt in the region of the pylorus; and there appeared to be no doubt that he suffered from chronic disease of the stomach. He had not suffered from hæmatemesis, neither did he complain of any pain at the stomach. The diagnosis of cancerous disease was confirmed at the inspection. In the hypogastric region was a tumour reaching as high as the umbilicus, precisely resembling in form a distended urinary bladder; it was dull on percussion, rounded in form, and fluctuation was distinct; it was also readily felt in the rectum. The patient stated that he never experienced any difficulty in passing water, nor had he any pain at the part. A catheter was passed without difficulty, and a few ounces of healthy urine were drawn off. On inspection, the thoracic viscera were found to be in a healthy state; the peritoneum also was healthy. The stomach was slightly distended, and on drawing it aside, a marked constriction was observed at the pylorus; and several of the glands at the lesser omentum and near the pancreas were enlarged and infiltrated with cancerous product. A firm growth was found to exist at the pylorus, extending into the stomach for about one inch and a half, where it terminated by a rounded, raised, and vascular edge; the valve was quite surrounded by the growth, and the surface was partially ulcerated. The growth had a similar vascular and raised edge on the duodenal aspect, but was there less prominent. The little finger could be passed through the pylorus. The growth was of a yellowish-gray colour, moderately firm, containing succulent fluid; it was composed of cells with large nuclei, free nuclei, elongated cells, &c., and was evidently cancerous. The liver, kidneys, and spleen, were healthy, so also the intestine. The ureters were not distended, the right one was spread out on the cyst, which occupied the usual position of the bladder; whilst the bladder was itself flaccid and situated on the left side of the hypogastric tumour. The peritoneum was smooth and healthy. The mucous membrane of the bladder, the prostate and urethra, the vesiculæ seminales, and vasa defe-

rentia were all normal. To the right of the bladder, in the median line, and apparently developed in the loose cellular tissue of the bladder was a large hydatid cyst, holding nearly three pints of small cysts, varying in size from a line to an inch in diameter, and full of clear fluid. At the base of the cyst was firm yellowish-gray substance, containing plates of cholesterine. The cysts beautifully showed their lineated structure, and numerous hooklets of the echinococcus were observed. The cyst had apparently commenced in the neighbourhood of the prostate.

The diagnosis of cancerous disease at the pylorus in this case was made out without difficulty; but there was much obscurity as to the character of the hypogastric cyst. There was no difficulty in micturition, and the patient was scarcely aware of the presence of the tumour; in some instances the ureters are compressed; this is shown in one of the instances recorded by Dr. Bright, the pelves of the kidneys became distended, and suppuration took place in them; in that case some of the hydatids were discharged with the urine. Occasionally the pressure is upon the urethra, when the cyst may still more easily be confounded with a distended urinary bladder. The vomiting in this case subsided to an unusual extent, ceasing for a fortnight.

The preceding cases indicate several important facts in relation to the symptoms and course of cancer of the stomach:— 1st. That the symptoms may be exceedingly slight, and the disease easily overlooked. 2nd. That the indications are more marked when the orifices are affected. 3rd. That the cachexia, the pain, the vomiting, &c., vary in almost every case, being sometimes slight, or altogether absent; in other cases intensely severe. 4th. That the onset of the severer symptoms may be very sudden; but it is generally preceded by a period of dyspeptic symptoms. 5th. That the disease is not limited to persons in advanced life. 6th. That it is sometimes associated with struma. 7th. That the occurrence of cancer with chronic ulcer of the stomach tends to explain some cases in which the disease extends over many years. 8th. That cancerous disease generally terminates within a year after a tumour has formed. 9th. That the mode of termination is greatly modified by the extension of disease to adjoining structures. 10th. That in most cases death takes

place from exhaustion or asthenia, and that fatal hæmorrhage and peritoneal perforation are more rare than in ulceration of the stomach. 11th. That the absorption of degenerating cancer-structure sometimes leads to symptoms resembling pyæmia. 12th. That some of the distressing symptoms may be alleviated; but that over-active treatment appears to hasten the fatal termination.

Although in several instances which we have recorded strumous disease was coincident with cancer, the former appeared to be in a quiescent state, and the two morbid processes were not, therefore, in active operation together; and they were so far antagonistic in character. Still there may be some unrevealed connection between these morbid states; for phthisical parents have children who die from cancer, and some who are apparently strumous in early life are affected at a later period with cancerous disease. We must hesitate, however, to consider as causative and connected what *may* be merely a coincident relationship.

CHAPTER IV.

FUNCTIONAL DISEASES OF THE STOMACH.

THE imperfect performance of the digestive process constitutes dyspepsia; but, this general term is the expression of an effect which arises from numerous causes, and it associates maladies which differ in their origin, in their course, and in their termination. Thus, at the commencement and throughout the course of organic disease of the stomach the food is imperfectly assimilated; but in the greater number of instances of gastric disease the dyspepsia is a transient symptom, and it entirely ceases after a longer or shorter period; and where other diseases are the immediate cause of death, we are often unable to find any structural change in the stomach, either in its secretions or component parts, although dyspepsia may have existed for some time; these cases, then, constitute what are ordinarily regarded as *functional* diseases of the organ, the conditions being either transient or of such a character as to be beyond our sphere of observation.

Dyspepsia arises:—1, from a disordered condition of the mucous membrane of the stomach and of its secretion; 2, from the muscular movements of the stomach being impeded; 3, from an abnormal state of the vascular supply; 4, from changes in the condition of the nervous system; and, lastly, from improper diet, or from chemical decomposition taking place during the digestive process. Several of these causes of dyspepsia may be combined; some lead to very transient maladies, others to irremediable disease.

1. Dyspepsia from an abnormal condition of the mucous membrane and its secretion.

The experiments and observations of Dr. Beaumont on Alexis St. Martin have pointed out the state of the mucous membrane which sometimes exists after improper food or stimulants have been taken; the surface of the stomach he found in such cases much injected, or erythematous. The secretion was diminished, and during this period more or less discomfort was generally produced; this condition entirely ceased in a short time, and the surface presented its usual appearance; but if death had taken place from some other cause during that condition of dyspepsia, the abnormal state would have disappeared, and no structural lesion have been discovered on careful or even microscopical inspection. This form of dyspepsia is, perhaps, one of the simplest degrees of acute catarrh or erythema, of which mention has been made in the last chapter.

The deficiency of gastric juice may be a relative rather than an actual one. A greater amount of food having been taken than is needed for the system, or than can be dissolved, it remains in its undigested state and acts as an irritant; the crude substance not only disturbs the stomach and its secretions, but if it be allowed to pass through the pylorus it acts upon the whole canal; or fermentative changes are set up, which we shall afterwards have to notice. Where excess of this kind is habitual more permanent results follow, and the symptoms resemble those found in some cases of diminished secretion.

In describing the forms of dyspepsia which have their origin in an abnormal condition of the gastric juice, we may divide them as follows:—1st, Dyspepsia from a deficiency of gastric juice; 2nd, from its excess; 3rd, from its irregular secretion; 4th, from its changed character, as in pyrosis, in gout, in rheumatism, and in albuminuria.

1st. The *deficient secretion of gastric juice* produces characteristic symptoms, and arises from many causes. After intemperance, either in eating or drinking, the gastric mucous membrane becomes over-stimulated, the portal system is at the same time engorged, and the liver is congested and disordered; in this state natural secretion does not take place in the stomach, and dyspepsia is produced.

In this hyperæmia or erythema of the gastric mucous mem-

brane the complexion becomes slightly sallow, and the patient complains of mental depression, lassitude, or headache; the tongue is furred, the appetite is impaired, and the condition of the bowels is disordered, as shown by a confined state, or by irregular action, with more or less pain. In some cases pain comes on at the scrobiculus cordis and between the shoulders, with thirst, nausea, and vomiting, and often with the rejection of green bilious fluid.

When excess is habitual the same symptoms are produced in a modified degree; the patient is hypochondriacal; he often believes himself to be the subject of serious disease of the liver; the bowels are constipated or irregular; flatulence, spasmodic pain or cramp in the abdomen, pain across the chest, tenderness at the scrobiculus cordis are produced; the tongue is furred, or its papillæ are distinct and injected, the appetite is lost, stimulants are longed for, and a bitter or nauseous taste in the mouth, especially in the morning, distresses the patient; the pulse is compressible, and a sense of exhaustion and of physical fatigue is attributed to actual loss of power. Sometimes also there are severe headaches, vomiting, disturbed special sensibility, as indicated by double vision, *muscæ volitantes*, noises in the ears, disturbed general sensibility, as manifested by numbness, formication, loss of sleep, or sleep disquieted by frightful dreams. In this condition food taken into the stomach remains undigested, and there is a sense of weight or "load at the chest;" the thick mucus covering the congested membrane prevents the action of the gastric juice on the alimentary mass, as in the instances of chronic catarrh previously described.

The deficient secretion of gastric juice does not, however, always arise from vascular plethora or active hyperæmia; but even from an opposite condition, namely, an anæmic state of the mucous membrane, and it may be produced by sedentary occupation, want of exercise, mental distress, over-excitement, anxiety, and insufficient food. The nervous energy of, and the vascular supply to the digestive organs, appear to be inadequate to their requirements, and digestion becomes oftentimes a painful process, from which the patient shrinks. This state is marked by pallor, and by an anxious expression of the countenance; the appetite is lost, or it is a fastidious one; the pulse is sharp,

irritable, but compressible; palpitation of the heart, throbbing sensations, and often pain in the head, are produced; the tongue is slightly injected in its papillæ, and has a whitish fur upon it, though in many cases the tongue is clean, large, and indented; there is sometimes nausea, or actual vomiting; the bowels are constipated or irregular; a sense of oppression or weight comes on after eating, sometimes followed by a throbbing sensation in the abdomen, and almost over the whole body, with languor or drowsiness; at other times there is faintness; and where undigested food passes into the pylorus and duodenum, violent cramp or spasmodic pain is produced.

Ingesta may be retained in the stomach many hours, and in some cases even days in a crude state; the secretion is not sufficient to dissolve what is placed in the viscus; the irritation produced by the retained food aggravates the ailment, and fermentation or decomposition is set up, with flatulence, pain, heartburn, or severe gastralgia. This, however, may arise from excess of food rather than diminished solvent power, as we have previously noticed.

In the *treatment* of dyspepsia following *excess* an emetic may be advisable, and not unfrequently this natural mode of relief takes place spontaneously, and the vomiting is preceded by pallor and faintness. If the irritability of the stomach continue, soda-water, or effervescent salines, as the carbonates of potash, soda, or magnesia with citric acid, may be administered, with or without the addition of hydrocyanic acid. The carbonic acid acts as an anodyne and sedative to the mucous membrane, and the saline compound which is produced relieves the portal congestion.

In this condition of great irritability of the stomach, in which the nausea and vomiting are sometimes excessive, and the disrelish for food well marked, even the sight or smell of it being distressing to the patient, the administration of alkalies is more beneficial than that of mineral acids, the former act as sedatives, rendering the abnormal as well as scanty secretion less irritating, and enable the diseased membrane more quickly to recover itself, and to pour forth its proper secretion; the latter act as astringents and tònics to a relaxed mucous membrane. Cold drinks and ice are often craved for, and when given in

moderation tend to relieve the congested state of the gastric membrane.

If more chronic effects have been produced, small doses of blue pill with rhubarb, and with magnesian purgatives, may be useful; for, by these means the portal system becomes freed from engorgement, and proper secretion takes place; should a sense of exhaustion then continue, it is well to give hydrochloric or nitro-hydrochloric acids with infusion of gentian or calumba.

The diet should be plain and easy of digestion, not rich or highly seasoned, and without stimulants; and three to four hours should be allowed to intervene between each meal. The character of the diet, and the quantity of the food taken are most important considerations; for meals taken too frequently or in excess, may, equally with the administration of improper and indigestible substances, be the cause of the malady. Before the stomach can empty itself it is often again irritated by a fresh supply; numerous dishes may prompt to intemperance, and excess is especially injurious when associated with late hours and deficient exercise. The function of the stomach is especially connected with the solution and preparation for absorption of the nitrogenous articles of diet, as they are present in our ordinary animal food; and in the conditions of active congestion and great irritability bland demulcent and starchy substances, as milk, arrow-root, &c., are to be preferred.

Again, imperfect mastication increases the difficulty of digestion; for the secretions of the stomach are then unaided by the division of the food, and the action of the saliva in changing the starchy portions into saccharine matter is not duly performed. This defective division of food may arise not only from the hurry of business and the force of habit; but also, because the agents of mastication are destroyed; and the dentist, by restoring the teeth, may afford the most effectual means of removing this form of dyspepsia.

But, in the class of cases where great congestion does not exist, constipation should be relieved by aloes and myrrh, or by colocynth with henbane; a dose of blue pill or of oxide of mercury is sometimes of advantage; and to rectify the condition of the stomach, capsicum with small quantities of ipecacuanha may be given to increase the secretion; dilute hydrochloric

acid to add to its solvent power, or carbonate of ammonia with bitter infusions to stimulate the vascular and nervous systems of the abdominal organs. These means, however, will be useless unless the diet be regulated, unless it be well masticated, and be followed by proper exercise.

In the dyspepsia produced by over-fatigue, anxiety, excessive grief, deficient food, or exhaustion from any cause, we find that stimulants are of great value; a small quantity of brandy, or a glass of wine acts as a healthy stimulus to the mucous membrane, and to the nervous system. In many cases where food could not be taken without the production of severe pain, or vomiting, the stimulant acts with the best effect.

When there is anæmia, as after parturition, miscarriage, or loss of blood, this form of dyspepsia is not unfrequent; and may be relieved by small doses of steel, often advantageously combined with quinine when there is no irritability of the stomach. The milder preparations of steel are, however, to be preferred, as the ammonio-citrate, tartrate or phosphate, the compound steel pill; and the liquor cinchonæ is a more elegant preparation than decoction of cinchona, and is often more easily borne than quinine itself; the bitter infusions, gentian, orange, cascarrilla, may also be prescribed with advantage with hydrochloric or nitric acids.

In imperfect secretion of gastric juice Dr. Ballard* has introduced into English practice the suggestion of M. Corvisart, to employ an artificial digestive fluid, in the form of pepsine, prepared from the stomach of ruminants. This is mixed with starch, and constitutes "Poudre Nutrimentive," and is prescribed in doses of 15 grains; to some of this lactic acid, hydrochlorate of morphia, or strychnine is added. In several cases in which I have tried the fluid pepsine (prepared by Mr. Squire), I have not yet met with the success expected from the high encomiums of Dr. Ballard. Rightly to estimate the value of pepsine as a remedy, the morphia and strichnia must be omitted, otherwise beneficial results may be improperly interpreted. Both these remedies are themselves valuable in atonic dyspepsia. In all cases it is desirable to remove the

* Ballard, 'On Artificial Digestion.'

causes of the imperfect secretion, if possible, rather than to supply a very imperfect artificial substitute.

The stimulant effects of coffee, ammonia, &c., are not so effective as those of alcohol in these cases, and brandy or wine is often better than malt liquor. In saying this we are very far from recommending the habitual use of such stimulants.

The habit of smoking, or snuff-taking, produces a relaxed and enfeebled condition of the mucous membrane, its secretions are not sufficient to ensure solution of the food ; stimulants are often resorted to, to counteract the effect, and many suffer severe dyspepsia from this cause.

In *advanced life* we sometimes find that the powers of digestion are greatly enfeebled and the gastric juice is imperfectly secreted ; there is pain at the scrobiculus cordis, and neither appetite nor relish for food ; the pulse is compressible ; the tongue is partially furred, and the bowels are inactive. It appears probable, that from the feebleness of the circulation, and from the degenerated condition of the vessels, the mucous membrane does not receive its proper supply of blood ; it is true that in age, repair takes place in a less degree than the destruction of tissue, and that emaciation is the result : but this waste may become so great, and the circulation be so enfeebled, that the patient may suffer from anæsthesia ; the brain may be unable to carry on its function, and syncope, vertigo, and even ramollissement of the brain may be produced.

In this state it is well to give food in its most nutritious forms, and more frequently than usual. Well-seasoned dishes, mustard, capsicum, salt, &c., are useful ; stimulants in small quantities are beneficial, and coffee, ammonia, with tonics are also called for.

In *very stout* persons, or those in whom the appetite has previously been pampered, we find feebleness of digestion, with a sense of weight or exhaustion ; spasmodic pain also, and irregular action of the heart, are easily induced. These symptoms arise in part from the feeble condition of the heart and circulation ; and are increased by an inactive state of the liver. The appetite is often small in stout persons ; and the hydrocarbons are stored up, instead of being removed in the ordinary changes of respiration, &c. ; but the mischief is still

further increased when the heart is irregular from an excess of fat about it, or when the feeble circulation of the brain manifests itself in vertigo and disordered sensations. Much relief is afforded by occasional alteratives, by aloes, rhubarb, and taraxacum, or by nitro-hydrochloric acid with bitter infusions; stimulants should be avoided if possible, and out-door exercise gradually increased.

2nd. Dyspepsia is also occasioned by an *excessive secretion* of gastric juice, which is apparently, in some cases, poured out in unusual quantity from a slight stimulant. A burning sensation at the stomach two or three hours after a meal, heartburn, pain in the back, are the usual symptoms of this state. It may be associated with hepatic or with cerebral disease, and it is then best combated by remedial agents calculated to relieve the exciting cause of the disease; imperfect secondary assimilation, such as exists in gout and in rheumatism, produces symptoms similar to those just mentioned, and they are probably due to a changed character of the secretion rather than to mere excess.

In the treatment of this form of dyspepsia the diet should not contain an excess of nitrogenous substances; and food should be slowly and thoroughly masticated; it is also important not to limit the patient to fluid forms of food, which are rapidly absorbed, and often leave an undigested sedimentary deposit; the evil of this form of dietary we have often seen in chronic disease, in which dyspepsia has apparently been produced, or, at least, aggravated by this cause. Stimulants are better avoided, or they should be taken in great moderation; whilst ardent spirits, and the stronger wines are better abstained from altogether, for a temporary relief does not compensate for the injury they perpetuate. Exercise, and the maintenance of good action from the skin are very important, especially when this gastric disturbance is combined with hepatic derangement.

As to other remedies, the carbonate or caustic alkalies with bitter infusions often afford almost immediate relief; but they do not remove the cause of the malady, and are ineffective unless the diet be regulated, and right exercise maintained. Creasote, combined with sedatives and aperients, greatly miti-

gates the distressing symptoms ; and in all these cases it is well carefully to watch that the bowels are not confined, and that the liver performs its functions. It is not necessary, however, to resort continually to blue pill or calomel to rectify any deviations from healthy action in the hepatic secretion.

3rd. Dyspepsia from an *irregular* secretion of the gastric juice is closely allied to the state just described. Such irregularity Dr. Budd has mentioned as one cause of the varied degrees of gastric solution observed after death, where other conditions have previously been the same. He states, very truly, that whilst in health the stimulus of food leads to the effusion of gastric juice, in disease it may be poured out without this stimulus. In cerebral disease, a perverted and craving appetite, and desire for food at unusual periods, may be due to this cause ; but, it is not uncommon to find those who suffer at irregular periods from burning pain at the stomach, which is at once relieved by partaking of a mouthful of biscuit or dry bread. When the complaint arises from gastric causes, the directions we have briefly given for the relief of the excessive secretion of gastric juice, may be found of service ; but where it arises from the former, namely, from cerebral disease, other symptoms will be generally present, such as pain in the head, a slow, labouring pulse, disturbed special or general sensibility. In this case, a spare and unstimulating diet is called for ; the free action of the bowels is desirable, and the avoidance of all cause of mental excitement is most important. Too frequently in children is a ravenous appetite found to be the precursor of organic disease of the brain ; and in mania the sufferer is oftentimes prompted to swallow the most extraordinary substances, as large quantities of gravel, possibly from a sense of gastric distress, and of unsatisfied morbid appetite.

4th. Dyspepsia produced by *morbid changes* in the gastric secretion leads us to other forms of disease ; and the first that we have to notice is *Pyrosis or water-brash*. This is a disease of frequent occurrence, and it receives its appellation from the fact of its principal symptom being the rejection of a thin watery mucus. The poorer classes of society, and especially its female portion, are the ordinary subjects of the complaint. Half a pint of thin watery fluid, somewhat resembling the white

of an egg, is sometimes vomited or regurgitated at once; it is generally neutral in its chemical reaction, and often tasteless, but sometimes it is found to be slightly alkaline, and the patient complains of its saltness. The period at which the discharge of fluid takes place varies both as to the hour of the day, and the frequency of the recurrence of the attack. The vomiting, however, generally occurs when the stomach is empty; and it is accompanied with a sense of contraction and of pain at the epigastric region and at the spine; with some patients the attack comes on in the forenoon, with others during the night, as at one or two in the morning, and even several hours after retiring to rest. As to other symptoms, the tongue may be clean, the pulse normal, the patient tolerably nourished, or anæmic and enfeebled; headache is often present, and in some instances the water-brash alternates with more severe gastralgia.

It is the opinion of Dr. Handfield Jones* that pyrosis is a chronic catarrh of the mucous membrane of the stomach, similar to blennorrhœa from the bronchi; and there is much to warrant this supposition. Dr. Chambers,† however, favours the idea that the œsophagus is the source of the discharge. The disease comes on after the continued use of oatmeal, and hence it is more common in the north; it may follow symptoms of chronic gastritis; and it is produced by great anxiety of mind, by exhausting disease, by over-fatigue, or by an overworked frame; it also occurs during pregnancy, and it is met with amongst the symptoms of commencing cancerous disease of the stomach. With such causes, it is not surprising that numerous instances of this disease are found among the out-patients of dispensaries and large hospitals.

The remedies which relieve pyrosis are astringents and tonics, as the sulphate of iron with the extract of logwood; quinine with aloe and myrrh; nitrate of bismuth alone or with conium and nux vomica; an alterative of blue pill, with rhubarb, is sometimes beneficial. Solution of potash, with hydrocyanic acid or with henbane and bitter effusions, is of great service when there is much pain. Other astringents may be advanta-

* Handfield Jones, 'On the Mucous Membrane of the Stomach.'

† Chambers, 'On Digestion.'

geously employed with sedatives, anodynes, and tonics, as the compound Kino powder, catechu with morphia or opium, oxide of silver, sulphate of copper, strychnia, or the infusion, tincture or extract of nux vomica.

CASE XC.—*Pyrosis*.—A Scotch woman, married and in comfortable circumstances, came under my care at Guy's Hospital, with pyrosis of several months' duration. The attacks of pain, followed by vomiting of watery fluid, recurred several times during the day. Bismuth, with other remedies, failed to afford relief, but the oxide of silver, gr. $\frac{1}{4}$, with extract of logwood, gr. ij, three times a day, produced very marked improvement. I have seen equal benefit result from sulphate of iron. Strychnia and nux vomica have been less serviceable in my experience of these cases than in those of atonic dyspepsia previously referred to.

A form of pyrosis is found to arise in connection with colloid cancer, watery fluid being regurgitated into the mouth; and it is important to bear this fact in mind in the diagnosis of colloid disease; and in ordinary pyrosis the symptoms are sometimes so severe and persistent as to cause hesitancy in our prognosis, and to suggest the presence of carcinomatous disease. The following case was one of that kind.

CASE XCI.—*Dyspepsia, Pyrosis*.—A gentleman, æt. 53, applied to me some months ago; his mind had been much overworked, and for more than two years he had suffered exceedingly, as much or more from mental depression as from actual disease. Whilst his energies were being overtaken he experienced sudden vomiting, and nearly every day water was regurgitated into the mouth; after two months he applied to a surgeon, but his symptoms increased in severity, and they were associated with languor and exhaustion. He went on the Continent, but the ailment continued; the sudden, severe pain at the stomach was only relieved by lying on the back; he could take but little sleep, and suffered occasional vertigo. He returned home, and afterwards went to Somersetshire, Brighton, &c., but without relief.

The fluid ejected was tasteless, clear, and like water; it was generally brought up three hours after eating, and at night

sometimes twice ; his nights were wretched. Various forms of medicine and diet had been tried, Prussic acid, bismuth, silver, nitric and other acids, gentian, soda, calumba, &c., but without apparent benefit. His countenance was natural, but his mind dejected ; the conjunctiva was watery, the tongue clean, the circulation feeble, the pulse compressible, the urine acid, sp. gr. 1020, not albuminous, neither did it contain crystals nor deposit ; nothing could be detected on palpation of the abdomen, but slight pain was produced at the scrobiculus cordis.

Steel and quinine were prescribed with capsicum and conium, and a sedative draught at night ; regularity of diet was enjoined, and he was encouraged to regard his case hopefully. These means, with a subsequent change of air, produced considerable improvement.

I looked upon this case as functional rather than organic disease of the stomach, and the subsequent history has so far confirmed the diagnosis.

Beside the abnormal conditions of the gastric juice already mentioned, there are two others which must be considered, namely, the dyspepsia occurring in what has been termed lithic acid diathesis, and the dyspepsia found in albuminuria. The former is especially observed in those who are the subject of rheumatism and of gout, and the following symptoms mark its presence :—A fastidious appetite, heartburn, flushes of heat, pain at the scrobiculus cordis and in the left hypochondriac region, a constipated or irregular condition of the bowels, a furred tongue, pain in the head, mental depression or unusual excitement, and sometimes severe vomiting and intense pain at the stomach.

The disease appears to be produced by imperfect secondary assimilation, as explained by Dr. Prout. The functions of other viscera are disordered, as of the liver and kidneys ; the motions become pale, the urine high-coloured, and it deposits lithates, or contains excess of uric acid. The heart and sympathetic nerve are affected ; there is often irregularity of the pulse, and there may also be vertigo or transient anæsthesia. The blood contains lithic acid, as shown by Dr. Garrod, or other elements from the decomposition of tissue ; and in this state the gastric juice is secreted in an abnormal state ; it becomes preternaturally

acid from lactic or hydrochloric acids, or is excessive in quantity, and may be otherwise changed.

This form of dyspepsia is easily produced when hereditary tendency exists; but, even where this is not the case, it may arise from over-stimulating diet, excess, and other irregularities.

In few dyspeptic conditions is the regulation of the diet more important, both as to its quality and its quantity; and it should consist of well-cooked, plain, animal food, with vegetables, the latter being in excess. Stimulants should be avoided, or the lighter wines taken; for although the immediate distress is relieved by ardent spirits, the disease is subsequently aggravated.

As to medical treatment, the first object is to mitigate present distress and pain; if the suffering be severe, chloric ether, chlorodyne, opium, or morphia may be resorted to; the salts of potash, soda, and magnesia afford relief to the heartburn and distress, and may be combined with carminative or antispasmodic remedies. It is, however, most desirable to remove, if possible, the cause of the dyspepsia, namely, the secondary mal-assimilation; and to correct an inactive condition of the liver, if such a state exists, small doses of blue pill may be combined with colchicum, rhubarb, aloes, and sometimes, also, with quinine. *Taraxacum* with bitter infusions, and with the alkaline bicarbonates, acts sometimes as a useful laxative. If, however, there be exhaustion and general feebleness of power, ammoniacal stimulants must be given in combination, and wine allowed. The saline mineral waters are in some of these instances justly recommended; those of Bath, Bristol, Buxton, and of Cheltenham are most likely to be of service amongst British springs; and on the Continent, the springs at Ems, Karlsbad, and Vichy may be resorted to.

Nothing will avail effectually, however, unless strict dietetic rules be observed, accompanied by exercise in the open air. If the meals be daily hurried, the mind constantly on the stretch from business occupations, the hours of rest shortened, and the consequent exhaustion removed by stimulating potions, the physician has no chance of affording relief. His advice, however, is frequently followed in a form that he does not direct; a visit to some lovely spot is tried (with globules or hydropathy);

cessation from business and anxiety is allowed ; stimulants, and late irregular hours are discontinued ; simple and unstimulating food is taken in moderation ; the patient quickly recovers, and the globules or wet sheets are lauded at almost every breath.

In *albuminuria*, the vomiting and nausea, which are amongst its most common symptoms, are generally considered as sympathetic ; and the renal plexus of nerves, in its connexion with the semilunar ganglion, with the pneumogastric nerves, and with the gastric plexus, is regarded as the exciting cause of the vomiting and nausea. This is probably in great measure the case ; but another cause exists, namely, the altered condition of the blood, and the excess of urea which it contains ; the urea is poured out with the normal gastric juice, and acts as an irritant to the stomach, and tends to neutralize the gastric juice. Urea has been demonstrated in the secretion from the bronchi, and, in these cases, it exists in all the secretions ; but, perhaps, in none, to a greater degree than in that from the stomach.* It is in vain to expect much relief from remedies directly applied to the stomach ; attention must rather be given to the disease of the kidney, and means employed calculated to restore the blood to its normal state, or to free it from poisonous excreta.

Diaphoretics, as antimony with acetate of ammonia, and salines ; purgatives, as jalap or elaterium ; warm baths or vapour baths will afford more relief than hydrocyanic acid or creasote ; and cupping from the loins will sometimes remove, or, at least, greatly mitigate the nausea and vomiting ; counter irritation may be applied to the loins or to the scrobiculus cordis ; and in some instances of extreme general anasarca, the gastric symptoms and the distress of the patient are greatly diminished by puncturing the thighs and thus allowing the serum gradually to exude.

There are other forms of mal-assimilation which occasion dyspepsia, and we find indications of this in some of the varieties of cutaneous disease. No organ sympathises more closely with the stomach than the skin ; in every period of life this fact is noticed ; in infants we have strophulus, and eczema from gastric irritation ; in adults some of the forms of urticaria

* Bernard, also Goodfellow, 'On Disease of the Kidney.'

and roseola, eczema, and lepra; in advanced life, eczema and prurigo, &c.

It not unfrequently happens that flatulence is produced by the formation of gas in the stomach, irrespective of the decomposition of food, to which reference will subsequently be made; and in cases of hysteria, or in prolonged abstinence from food, &c., the stomach sometimes becomes painfully distended, eructations take place, and the power of digestion is diminished. It has been supposed that gas is effused from the capillaries, but of this we have no proof; and equally hypothetical is the opinion that it arises from mucus becoming decomposed by gastric juice, and thus evolving gaseous products; the flatulence is generally preceded by slight pain, or by a gnawing sensation at the *scrobiculus cordis*; a full meal in this condition will probably not be digested, but the flatulence will be prolonged, and colic be produced. The better method is to take a small quantity of nourishment, with some stimulant—a cup of coffee, or a glass of wine—and afterwards a more substantial repast, giving time for thorough mastication.

II. The *impeded movements of the stomach* are not sufficiently considered as causes of dyspepsia. In hernia when the omentum is fixed and the stomach is dragged from its position, pain in the hypochondrium is produced; and the habit of tight lacing, which few young ladies are willing to admit, is a fertile source of the same suffering; in most cases the mischief is done very early in life, the ribs are scarcely allowed to expand, and the stomach is gradually tilted into a vertical position whilst development is taking place. Neuralgic pain in the side, flatulent distension of the stomach, pain after food, spasm, borborygmi, hysteria, are the usual sequences of this folly. Digestion requires that the nutriment should slowly revolve round the stomach, and as it is converted into chyme, that it should pass into the duodenum. When the stomach is placed vertically, its semi-digested contents are more likely to be impelled at once into the pylorus.

The dyspepsia which is so common in those who spend many hours over the desk, in writing, or in reading, or in constrained position, is of the same kind; and amongst tailors, shoemakers, dressmakers, &c., this unnatural and long continued posture is

productive of severe indigestion, which is increased, in many cases, by irregular and intemperate habits.

Constant pain at the *scrobiculus cordis* and between the shoulders is complained of; eructations sometimes distress the patient, the bowels are often constipated, the tongue is furred, and the mind depressed. We may often do much to remove the disease by enforcing an erect posture during the hours of occupation, by strict attention to diet, by well regulating the bowels, by relieving torpor of the liver, and, if needful, by administering mild alteratives, or by nitric acid with *taraxacum*.

In tumours developed at the lesser omentum, or at the pancreas, &c., the pylorus becomes pressed upon, and a free passage is prevented; in this case, also, we find pain and heart-burn, and sometimes the obstruction is sufficient to produce vomiting, as in Case CXIV.

In other instances, the movements of the stomach are prevented by the presence of fluid in the peritoneal cavity; in ascites and in ovarian dropsy the stomach may be so much pressed upon, that expansion cannot take place, and after food its contents may be rejected or severe pain be produced.

It is probable that in some cases of over-distension from flatus, the muscular coat of the stomach is unable to contract, or becomes paralysed. Dr. W. Philip gives such as his opinion; and cases are not very rare in which, after death, we find the stomach occupying nearly the whole of the abdomen, reaching to the pubes, and apparently causing death, by interfering with the action of the diaphragm and of the heart. Lesser conditions doubtless arise, and are attended with much discomfort, as a sense of distension, flatus, and sometimes intense pain. The symptoms are relieved by ether, by antispasmodics, by the gum resins, as *galbanum*, *assafœtida*, &c.

It must be borne in mind, however, that this tympanitic state sometimes arises from inflammation coming on insidiously, and involving the muscular as well as the peritoneal coats, as in some cases of strumous peritonitis. I have seen several such instances, in which fatal results followed without any pain from the commencement to the close. A short time ago, a policeman complained of fulness of the abdomen, which gradually became tympanitic, but no pain was produced; this state increased for

six weeks, with prostration ; about a fortnight before death, the tympanitis was less, and fluctuation indistinct. He gradually sank, about ten weeks from the commencement of the illness, but he suffered no pain throughout. There was chronic peritonitis, the whole serous membrane being studded over with whitish grains of lymph. There were bands of adhesion, and the peritoneum contained several pints of bloody serum. The serous investment of the spleen was a quarter of an inch in thickness, and contained small opaque cheesy masses. The small intestines were matted together, but not very firmly, and the ileum presented several passive ulcers. In the lungs, at the left apex, was puckering and iron-gray consolidation. We might readily mistake such cases for ordinary dyspepsia with flatulence, since they occur in youth as well as in middle life.

III. Dyspepsia from changes in the vascular supply has already been noticed ; the acute and chronic forms of catarrh, the erythematous condition produced by intemperance and hepatic congestion, and the converse condition of anæmic dyspepsia, have each been described, and we must refer to our remarks upon them.

IV. An appreciation of the condition of the *nervous system* is important in studying disease of the stomach ; diseases in other organs lead to disturbance of this viscus by their nervous and sympathetic relations with it. We find vomiting induced by diseases of the brain, as in concussion, in disease of the membranes, in hydrocephalus, in some cases of abscess of the brain ; by uterine irritation, as in pregnancy and ovarian disease ; by gall-stone ; by disease of the liver, of the kidneys, and suprarenal capsules ; by renal calculus, by intestinal obstruction, by changes in the blood, by some diseases of the lungs ; and by the onset of acute disease, &c. Each of these conditions produces peculiar and characteristic symptoms, but all may be accompanied by violent and most distressing vomiting, and unless care be taken in the investigation, it may obscure the primary malady.

Disease of the stomach, however, leads to sympathetic disturbance of all these viscera, as of the head, causing pain, disturbed vision, *muscæ volitantes*, throbbing in the head and ears, *tinnitus aurium*, and also of the hepatic and renal secre-

tions; and, as it has been justly observed by Dr. Philip, these secondary conditions may become so severe as to be more persistent and trying than the disease of the stomach itself; thus intense neuralgic pain in the face and head may have the stomach primarily for its source.

An equally marked connexion of disease, arising from the state of the nervous system, is the sympathy of the stomach with the lungs; thus dyspepsia gives rise to dyspnœa, and to cough, from the irritation of the gastric branches of the pneumogastric, producing reflex irritation; so also with the heart, by means of the cardiac branches of the same nerve; for palpitation or irregular pulsation may be due to gastric disturbance, and may simulate severe organic disease of the heart.

In phthisis, it has been long noticed, that indigestion may precede the physical signs of disease in the lungs; nausea, loss of appetite, impaired digestion, furred tongue, pain at the scrobiculus cordis, and sometimes severe vomiting being followed by cough, and, after a time, by hæmoptysis and the general signs of tubercular disease.

The observations of Dr. Theophilus Thompson, in reference to the state of the gums in phthisis—a red injected line being produced along the margin of the teeth—is a further confirmation of the irritated condition of the mucous membrane. This early state of phthisis is that in which the greatest benefit is derived from prophylactic treatment; by change to salubrious or sea air, by attention to warmth and clothing, the avoidance of night exposure, by taking cod-liver oil, and sometimes vegetable tonics, a fatal disease may in many cases be warded off.

The pain which is associated with several of the forms of dyspepsia already mentioned, is at times very severe, and appears frequently to be of a neuralgic character, rather than a sign of inflammation or acute disease; a state of excessive irritability of the stomach is often induced, and the contact of any substance is followed by instant rejection. Uterine disease, as leucorrhœa and dysmenorrhœa, may be the cause of this excitable condition of the sympathetic nerve; and these symptoms may exist without producing any emaciation in the patient.

It is to this condition that Sir Henry Marsh has given the name of "regurgitative disease," in which the food or the

greater part of it is regurgitated rather than vomited; and this takes place without previous nausea, or progressive emaciation; and pain may be entirely absent.

We may here mention, that intense pain often results from ulceration of the stomach, and we have found large branches of the pneumogastric nerve involved in the thickened edges of the ulcer, where this symptom, accompanied by vomiting, had existed for many weeks before death. The pain from gastric disease may be situated in the side, or even at the left shoulder.

In the treatment of these forms of gastralgia, much relief is afforded by hydrocyanic acid, by creasote, by calcined magnesia with opium, by chloroform or chloric ether, or by nitrate of bismuth with conium; but opium does not always act favorably, in some instances it appears to increase the disease, even when a few drops of the solution of morphia act beneficially. Small blisters applied to the scrobiculus cordis or to the spine, sometimes alleviate the symptoms.

These remedies may be used in several of the forms of sympathetic irritation of the stomach, although more applicable to cases of true gastralgia. When the symptoms result from pregnancy, the mineral acids will sometimes afford relief. In this condition of excessive irritability it is often advisable to omit all medicine, and allow the stomach to rest. A case of this kind, admitted into Guy's Hospital, was cured by administering, for about a fortnight, nutrient enemata, and only a teaspoonful of water was occasionally allowed to relieve thirst. Nourishment was afterwards given gradually in the usual form, but without producing any vomiting. Another plan may be followed, recommended by Dr. Hunter, of giving a small quantity of milk and water every ten minutes or half hour.

Many of these cases are associated with hysteria, or with dysmenorrhœa, or leucorrhœa; but a much more troublesome form of irritability of stomach is found in chronic gastritis, combined sometimes with ulcer, sometimes with strumous or chronic peritonitis, or with phthisis; an inflammatory condition exists with one of extreme irritability, and the best means of affording relief is by the use of demulcents and anodynes previously mentioned. This state must be distinguished from the irritability of the stomach before referred to, in connection

with phthisis, for we have in these latter cases the symptoms of gastric disease in the red and deeply injected tongue, with pain at the scrobiculus cordis, &c.

Calomel is used by some as a sedative to the mucous membrane of the stomach; but since this condition of irritability is so frequently found associated with an anæmic and chlorotic or hysterical state, the administration of mercurials, except as occasional aperients, is better avoided.

Another form of dyspepsia, which primarily arises from the condition of the nervous system, has been noticed in reference to deficient secretion of gastric juice; namely, the dyspepsia in hypochondriasis, and in an overworked or imperfectly developed brain; this condition is exceedingly distressing to the patient, and equally trying to the physician; it is sometimes the precursor of epilepsy or of mania. In these instances of dyspepsia the whole attention is occupied by the diet, the mind is depressed, and its energies enfeebled; one change after another is tried, but pain and discomfort equally follow; the stomach is sometimes exceedingly irritable; the bowels are over-anxiously watched, the sleep is unrefreshing, and life rendered miserable. To tell the patient nothing is the matter, would be to drive him to some one who would give an opinion more in unison with his feelings. By carefully regulating the diet and the bowels, by cold sponging, by taking frequent exercise, either walking or on horseback, or a pedestrian tour when it is possible, by keeping the mind free from anxiety, and by cheerful society and occupation, all the symptoms may be greatly relieved.

In some men we observe a state closely resembling hysteria; as shown by flatulence, loss of appetite, sensibility of the surface of the abdomen, sensations almost amounting to globus hystericus, disturbed cerebral function, depression, anæsthesia, incapacity for exertion, &c.; in this condition, which is often combined with distended colon, I have found marked benefit result from the use of aloes combined with steel; fresh air and vigorous exercise are important remedial agents when they can be attained.

In other cases much resembling those just mentioned, the head is badly formed, and the forehead is narrow; the body is well

nourished, but the patient complains of pain at the scrobiculus cordis and in the back, or in various parts of the body; the mind is depressed, and the appetite irregular. Although muscular, a man may be quite incapacitated for exertion; the tongue may be clean, the bowels regular, the evacuations normal, or pale, the pulse tolerably full, or depressed and irregular. It would seem that dyspepsia has arisen from ordinary causes, but the sympathetic nerve reacts upon the cerebro-spinal centres, and they being easily disturbed from their healthy balance, again react upon the sympathetic nerve, perpetuating and aggravating the original and slight malady. In this we find the close connection between dyspepsia or disordered chylopoietic viscera, and mental disease, mania, and melancholia.

By acting freely on the bowels so as thoroughly to unload the colon, and by the steady perseverance in the milder preparations of iron, this state may be greatly relieved.

V. The *fermentation* of the contents of the stomach, and the symptoms consequent upon it, are due partly to an abnormal state of the secretions, in part to the muscular movements being impeded, or the pylorus obstructed, and sometimes to the character of the food itself. Dr. Budd has distinguished several varieties of fermentation; so also Dr. Turnbull. 1. The formation of carbonic acid, as in ordinary fermentation. 2. The formation of *sarcina ventriculi*. 3. Lactic or butyric acid fermentation; and, 4. The formation of sulphuretted hydrogen by simple putrefactive decomposition.

When the pylorus is obstructed by cancerous disease, spasmodic contraction, and tumours, the contents of the stomach are prevented from passing onwards; the viscus becomes distended by flatus; pain is produced; and vomiting, which affords partial relief to the patient, generally follows a few hours after food has been taken; the ejected matters are found partially dissolved, and undergoing fermentation; they have a sour smell, and a yeastlike surface; this action is allied to simple fermentation; alcohol is formed, and carbonic acid evolved; some acetic acid is produced; and the *sarcina ventriculi* discovered by Mr. Goodsir are often detected.*

Fermentation of this kind, and the presence of *sarcina*, may

* The *Merismopædia Ventriculi* of Robin, Pl. xii, fig. 1, p. 331.

exist without any pyloric obstruction or organic disease ; and *sarcinæ* have been detected in the urine, in the fæces, in pus, in pulmonary abscess, and on the healthy mucous membrane ; Robin even states, "ce végétal semble être sans action nuisible sur l'animal qui le porte." Fermentation may be produced by the imperfect mastication of food, and by taking exercise immediately after it ; by drinking fermenting or new malt liquors ; by indigestible vegetables, and fruit ; by new bread, salads, &c. Distension is felt almost at once, regurgitation of food into the œsophagus, eructation, palpitation of the heart, &c., take place ; colic is often produced, and sometimes diarrhœa, by the continuance of the fermentation, or by the presence of semi-digested substances in the intestine.

In the more severe cases arising from obstruction the sulphite or hyposulphite of soda, as recommended by Dr. Jenner, is a valuable remedy ; the sulphurous acid is set free, and checks the fermentative action. Charcoal has the same effect, so also creasote. The spasmodic pain from distension is relieved by sulphuric or chloric ether, by chloroform or by opium. In the more easily remediable cases arising from fruits, vegetable, or undigested food, an emetic or purgative may be given, and may be advantageously followed by ipecacuanha and capsicum, or by the nitro-hydrochloric acid with calumba, cascarilla, or gentian. These medicines apparently increase the secretion of the gastric juice or improve the tone of the mucous surface ; but after the immediate relief of the urgent symptoms the most likely plan to afford permanent benefit is by changing the diet, and by taking such substances as the stomach can easily digest.

Another form of chemical change described is that which takes place from the fermentation of starchy elements, milk, &c., and which leads to the formation of *lactic or butyric acid* ; severe heartburn is produced with pain at the stomach and between the shoulders, sometimes with vomiting, but without distension ; the pain is occasionally very severe and persistent, even after vomiting ; there is often a sour, nauseous taste in the mouth, and there may be spasmodic attacks, or even alarming collapse. The state is much relieved by creasote, opium, bismuth, or by magnesia and hydrocyanic acid.

In infants the most severe collapse ensues from the coagulation

of milk in the stomach, and the patient may be utterly prostrate, as if suffering from perforation of the intestine or from cholera ; if recovery take place, small masses of casein and fatty matter are sometimes passed from the intestine.

CASE XCII.—*Infantile Dyspepsia, Sudden Collapse*.—An infant about a year old was seized with sudden collapse shortly after being fed, deathly prostration followed, and it was believed by the parents that the child was poisoned ; the flour, milk, water, &c., of which the food had consisted, were carefully analysed by my friend Dr. Odling, and pronounced normal. The infant became cold, and was apparently in severe pain ; its eyes were sunken, and, after a few hours, several masses of cheesy substance, about half an inch in length, were passed ; these I carefully analysed, and they were found to consist of oily matter and casein ; and the symptoms arose from milk coagulated in the stomach having passed into the duodenum in a solid form. Such at least was my diagnosis of the case ; and the rapid recovery of the little patient showed the correctness of the opinion.

In some persons affected with dyspepsia the breath becomes exceedingly offensive, almost of the odour of sulphuretted hydrogen, being similar to that caused by carious teeth, diseased tonsils, or ulcerated nares. This state is due to the *putrefactive decomposition* of food retained and undigested in the stomach ; it is associated generally with vitiated secretions ; there is headache, mental depression, the tongue is furred, a sense of uneasiness at the stomach comes on, or pain in the bowels ; the evacuations are sometimes dark and unusually offensive, or there is slight diarrhoea. It would appear that, to some extent, effects similar to those observed when sulphuretted hydrogen is respired are the result of this state, and that the blood itself is contaminated by the absorption of gas from the alimentary canal. Putrefactive decomposition may also arise in obstructive disease at the pylorus.

In cases where no obstruction exists, it is well to prescribe a warm saline aperient, as sulphate of soda, and the potash tartrate with aromatic spirit of ammonia ; again, rhubarb, soda, and calumba, or the compound gentian mixture, may be advan-

tageously given. Creasote tends to check the decomposition, but its employment is less suitable in these than in previously mentioned instances.

HÆMATEMESIS.

There are several symptoms of disease of the stomach which demand separate notice, and the first of these to which we shall allude is hæmatemesis, or vomiting of blood.

Great alarm is naturally excited by the rejection of blood from the stomach, whether in small or large quantities; but the import is very different, for whilst in some cases it is a symptom free from danger, in others it is the indication of serious, if not of fatal disease.

The causes of hæmatemesis are:—

1st. Ulceration of the stomach.

2nd. A congested or obstructed state of the portal circulation.

3rd. Vicarious menstruation.

4th. Cancerous disease.

5th. A vitiated state of the blood, as in purpura, yellow fever, typhus, &c.

6th. Aneurism.

The hæmorrhage may, however, have its origin in parts connected with the mouth, the throat, and the œsophagus (as from ulceration, cancerous disease, and aneurism, and from varicose conditions of the œsophageal veins*), and the rejection of blood from these sources may be erroneously regarded as hæmatemesis; or, it may proceed from the nose, the larynx, and the lungs, and in some cases considerable difficulty arises in distinguishing the source of the discharge, for the blood may be swallowed and afterwards vomited.

As to the quantity of blood exuded, there may be the greatest diversity; sometimes it is only recognised by the most careful, or even microscopical examination; at other times several pints or even quarts are rejected at once; and if a large vessel have been divided, the first hæmorrhage may cause fatal syncope.

* 'Schmidt's Jahrbuch,' 1859, Le Diberder und Fauvel.

Blood which is thus discharged into the stomach is generally coagulated, and is often deepened in colour by the action of the gastric juice; it is devoid of the bright, frothy appearance presented by blood from the lungs, which is consequent on the admixture of air. A portion of the blood in the stomach becomes still further acted upon by the gastric juice, and passes into the duodenum. As it extends along the small and large intestine, the depth of the colour is increased, and at last it is discharged as a pitchy, liquid stool, constituting *melæna*. Sometimes this black evacuation or *melæna* is the only symptom of hæmorrhage into the stomach, for no blood may be rejected by the mouth; and when the blood is effused into the small or large intestine, and discharged, the depth of the colour is proportionate to the length of the tract through which the blood has passed, but it never assumes the black colour to which we have referred.

The green fluid which is sometimes vomited in states of great irritation of the stomach has been regarded by Dr. Fraser as altered blood; and the coffee-ground substance so often rejected towards the close of organic disease of the stomach, consists also of blood which has slowly exuded, the hæmatine being acted upon by the gastric juice. In some cases of purpura, a similar appearance is presented from a like cause. Much discussion has arisen as to the possibility of the transudation of blood through *unruptured* capillaries; but the examination of a portion of intestine distended with blood, and presenting points of *ecchymosis*, as found after disease of the mitral valve, will suggest the probable explanation of instances in which blood has been vomited or discharged, and in which no apparent perforation of vessels has subsequently been found. In such a portion of intestine as is present with mitral valve disease, some of the capillaries are found to be beautifully injected, whilst others are collapsed, and blood is extravasated around them, but limited by the basement membrane, thus constituting a point of *ecchymosis*; if the basement membrane had given way, the blood previously extravasated would have escaped, and no ruptured vessel would have been detected. A similar action takes place in the stomach; *ecchymosis* is produced, but the action of the gastric juice prevents our observing the changes with the same facility as in the intestine. There is little doubt that the

capillaries thus become over-distended, and then ruptured in the ordinary form of hæmatemesis, when no ulceration has taken place.

The symptoms which precede hæmatemesis are a sense of faintness, followed by weight at the scrobiculus cordis; the countenance becomes pallid, the pulse compressible and failing, the extremities cold, and sometimes actual syncope takes place; vomiting is then produced, and several pints, or even quarts, of half-coagulated blood are rejected; the patient becomes faint, blanched, and the bleeding is checked. After a few days or hours, there may be return of hæmorrhage, till at last, in some cases, the patient appears almost drained of blood. The subsequent symptoms are especially due to this loss, as found in other instances of anæmia; and severe headache, noise in the ears, disturbed vision, dilated pupil, palpitation or irregular action of the heart, with a sharp but compressible pulse, are present. If a large vessel have been divided, the first attack may, as we have before remarked, lead to fatal syncope. This sudden termination is, however, unusual; the patients generally slowly rally, and after a few days, the black, pitchy discharge of altered blood takes place from the bowels.

The character of the disease which has led to the hæmorrhage must necessarily modify the preceding as well the general symptoms and their termination; thus, in ulceration of the stomach and in cancerous disease, the peculiar symptoms of those maladies are present; in aneurism a pulsating tumour may sometimes be felt, and severe local pain, or pain in the course of the spinal nerves may be experienced. In congested portal system the signs are those of engorged liver, as shown by pain in the right side, dyspepsia, a sallow or semi-jaundiced complexion, furred tongue, occasional nausea or vomiting, impaired appetite, spasmodic pain at the stomach or in the region of the colon, constipation of the bowels, disturbed sleep, and pain in the head; enlargement of the liver and hæmorrhoids are also frequently present.

In vicarious menstruation, local congestion of the mucous membrane, or of the edges of a pre-exciting ulcer—as we sometimes find in an ulcer on the leg—leads to the effusion of blood into the stomach. In these cases we may have very slight

symptoms—absence of the proper menstrual discharge, slight pain in the side, and periodical vomiting of blood, without constitutional disturbance, and without the blanched countenance that we find in hæmorrhage from other causes. With this vicarious discharge we not unfrequently find hysteria, neuralgic pains, and leucorrhœa, &c.

In purpura hæmorrhagica there is a blanched countenance, faintness, &c., but we have indication of the cause in the changed character of the blood, and its effusion from other mucous membranes and into the skin. The hæmatine is probably acted upon, and the corpuscles broken down, so that actual exosmosis of coloured serum takes place.

During the course of fever, hæmorrhage from the bowels, apparently of a critical character, occasionally takes place; the patient who may be in a state of great prostration, with a dry and brown tongue, rapidly amends, and hence the discharge of blood has been regarded by some as indicating a “crisis” in the disease. In the few cases of profuse hæmorrhage which have come within the sphere of our own observation, the effusion of blood has probably taken place from ulcerated surfaces; in one, presently to be detailed, minute ulcers were found in the stomach, from which a profuse and fatal hæmorrhage occurred; in another instance, a young woman, whilst prostrate from typhoid fever, suffered from hæmorrhage to a great extent from the bowels; the patient became blanched, the pulse for many hours could scarcely be felt, but very slowly she completely recovered. These cases resemble those in which very minute quantities of blood are detected on microscopical examination of the evacuations during typhoid fever, but must be distinguished from the hæmorrhages described by Dr. Kennedy* as occasionally taking place during typhus without ulceration, and followed by rapid recovery.

When blood is poured out from the œsophagus or mouth, it is regurgitated or rejected without effort, rather than vomited, and we generally find either dysphagia or ulceration of the throat, &c.

The blood from the lungs is sometimes so retained in a vomica or dilated bronchus, that it loses its frothy appearance and florid

* ‘Dublin Quarterly Review.’

colour, and the patient is often scarcely able to tell us whether he vomited or coughed it up; no actual cough may be produced, but the blood may easily be brought up into the throat and then spat out, or it may be swallowed and then vomited, or discharged by the bowels; in these cases we attach much importance to the general signs of disease, and to the physical examination of the lungs and heart.

As to the prognosis in hæmorrhage from the stomach, we must bear in mind that it is rare for a patient to die from simple hæmatemesis, although such cases occur; patients appear to be almost bloodless, but steadily convalesce. Still the cause of the symptom must be our guide as to its termination; sudden and large bleedings after symptoms of organic disease should always be regarded with alarm, for ulceration often extends into the larger arteries, and the dense fibrous tissue prevents contraction of the adjoining tissues, and thus the hæmorrhage persists unchecked.

As to the treatment when bleeding takes place from ulceration or cancerous disease, the use of styptics is advisable—alum with dilute sulphuric acid, acetate of lead, gallic acid, catechu, tincture of iron, or oil of turpentine, may be used; but in cases where it arises from congestion of the liver, I have generally looked upon the hæmatemesis as to a great extent curative, and prescribed remedies calculated to relieve the congested liver, as a grain or two of blue pill with conium, and magnesia mixture, so as to remove the effused blood from the intestines.

Ice and cold drinks are grateful to the patient, and beneficial in producing contraction of bleeding vessels; but food should be abstained from, because coagula may be removed by it from divided vessels, and hæmorrhage be again produced. After a short time, fluid, demulcent nourishment may be given, but it should be in a nearly cold condition; and when there is the evidence of a cessation of the hæmorrhage, solid substances, easy of digestion, may be given in small quantities. Vegetable tonics with mineral acids, and the milder preparations of steel, will then be found of service; but we shall be often much disappointed by the various astringents, as gallic acid, alum, &c., which afford only partial relief; oil of turpentine, in doses of mxx , has been much recommended, and has been followed by

beneficial results ; its stimulant as well as astringent effects have been well marked.

It is exceedingly important that the patient should avoid those habits or excesses which have led to the disease, but advice on this subject is generally disregarded.

In vicarious menstruation, our efforts should consist in endeavouring to establish the proper and natural discharge, rather than immediately to check that from the stomach, unless it be excessive. Hip baths, steel, aloes and myrrh, change of air, exercise, the avoidance of all tight lacing or unnatural excitement, will probably restore the health, but this hæmatemesis will sometimes continue for a considerable period.

In purpura, the preparations of steel with acids are generally the best remedies that we can use, as the tincture of the sesquichloride or the sulphate of iron, with sulphuric acid ; and I have more confidence in these than in gallic acid or the oil of turpentine. The latter sometimes produces vomiting, and patients complain of its offensive character.

CASE XCIII.—*Hæmatemesis, from Cancer of the Liver.*—The most marked case of hæmatemesis and melæna from this cause that I have ever witnessed, was in a man about 55 years of age ; he was in an emaciated condition, cachectic and semi-jaundiced ; the liver was enlarged, and it was believed that he suffered from cancerous disease of that organ, a diagnosis which was found, after death, to be correct. He was suddenly seized with violent vomiting of blood, and black stools were passed from the bowels ; in about eight hours he died. On inspection, we found cancerous disease of the liver ; there was no ulceration in the stomach, nor evidence of ruptured vessel ; the intestine contained a considerable quantity of blood, but no ulcer. On opening the vena porta, it was found that the cancerous disease had extended into the vessel, and completely occluded it ; and that softened cancerous matter was injected along the branches of the vena porta, so as completely to check the circulation. The cause of the hæmatemesis was at once apparent—the capillaries of the stomach had become suddenly engorged with blood, and had ruptured, leading to the fatal hæmorrhage ; but no openings nor ruptured vessel could be found, for the distension had disappeared, and the minute vessels had collapsed. A similar result is found in many instances of vomiting of blood after intemperance.

CASE XCIV.—*Hæmatemesis from Portal Congestion.*—James P—, æt. 45, residing at Milton Street, was admitted into Guy's Hospital, February 2nd, 1859 ; he was a man of intemperate habits ; and whilst at work some time previously, packing hay and exerting his strength, sickness and flatulence came on, and he vomited about a pint and a half of grumous blood, and

afterwards smaller quantities of clear blood. He was under treatment for fourteen days, and then returned to his work. From that time he had had pain across his chest, which sometimes moved to the epigastrium with much flatulence. Three weeks before admission, immediately after jumping to reach a handle above him, he vomited up about half a gallon of brown-coloured blood in clots; and some blood passed per rectum. When brought to Guy's he had a yellowish, semi-jaundiced complexion, and suffered from pain at the scrobiculus cordis; the lungs were healthy; the pulse was full, soft, 80; the tongue coated; the appetite defective; the bowels open; the urine not albuminous. He was ordered infusion of roses with acid, and milk diet. 4th.—There was slight pain, no return of vomiting, but he had passed blood by the bowels; the tongue was furred. 8th.—He appeared nearly well, and was soon afterwards presented.

This case of hæmatemesis probably arose from hepatic engorgement, due to intemperance; and the hæmorrhage from the over-congested mucous membrane of the stomach was in itself curative.

CASE XCV.—*Hæmatemesis after great Intemperance.*—Alfred W—, æt. 38, admitted into Guy's under my care in May, 1855, was a tall man, perfectly blanched in appearance, and on admission he was almost in a state of syncope. He had been for some time a porter at the Brighton Railway, and had drunk very freely of spirits, although accustomed to eat but little food. During the Epsom races, having harder work than usual, he drank still more intemperately; he had been troubled with occasional pain at the stomach, and with vomiting. The day before admission he felt a sense of weight at his stomach, which he tried to relieve by taking more spirits; a feeling of faintness came over him, and he vomited several pints of dark-coloured blood. He was much excited on admission, and there was considerable tremor of the hands. The skin was moist, the tongue and lips pale, the bowels confined. The liver was much enlarged, and there was slight tenderness at the scrobiculus cordis.

There was evidence in this case of great engorgement of the portal system, and although some additional hæmorrhage took place, I adopted the plan of endeavouring to relieve the distended liver, and constipated bowels, rather than of administering styptics. Blue pill and conium were given, and magnesia mixture. In this way black blood, acted upon by the gastric and intestinal secretions was discharged, and the patient rapidly improved. The hæmorrhage returned slightly on the third day, probably from spirits surreptitiously obtained. He steadily,

however, convalesced; food was given as he could take it, and afterwards steel medicine.

Most of these cases arise from the rupture of over-distended capillaries, rather than from ulceration, and we may generally give a very favorable prognosis. Where ulceration exists, and arteries are perforated by the disease, a fatal result sometimes ensues; several cases of this kind are recorded with ulceration of the stomach, in one of which, although fatal hæmorrhage took place, nearly all the blood passed into the duodenum, and scarcely any was vomited. Death, however, does occasionally follow without any ulceration being detected.

CASE XCVI.—*Hæmatemesis from Ulceration of the Stomach.*—Sarah J—, æt. 35, was admitted into Guy's Hospital, April 21st, 1859. She had been a domestic servant, was of temperate habits, and tolerably well nourished. Eleven years previously she had had an attack of hæmatemesis, and six months before admission a second attack had come on. Menstruation was scanty. On April 21st, she was brought to Guy's in a fainting state; two days previously menstruation had commenced, and on the same evening there was a sense of coldness, followed by vomiting of blood in considerable quantity. The hæmorrhage was repeated the following day. Conium and blue pill were given night and morning, and magnesia mixture three times a day.

On 23rd, she was blanched, and the pulse was compressible; no tumour could be felt at the stomach, nor was there any vomiting, nor pain between the shoulders. There was no evidence of thoracic disease.

30th.—The symptoms were much relieved, and soda and calumba were given.

May 15th.—Ammonia citrate of iron were ordered twice a day. She was convalescent, and regaining colour in the lips and countenance.

CASE XCVII.—*Hæmatemesis from Ulceration of the Stomach.*—Sarah M—, æt. 31, was admitted into Guy's Hospital, December 27th, 1858. She had resided at Lewisham, was married, and had had four children, and was of temperate habits. For three weeks before the attack of hæmorrhage she had complained of a sense of weight at the scrobiculus cordis, and of burning pain at the sternum; she had no pain between the shoulders, and the bowels were regular. On December 27th, after hurrying to the railway station, she vomited a large quantity of blood, and she was at once admitted into the hospital. The abdomen was supple, the tongue clean, pulse compressible. There was no enlargement of the liver, nor of the spleen; the menstruation was regular. She was ordered acetate of lead, gr. ij, opium gr. j, alone, afterwards conium and blue pill and magnesia mixture three times a day.

29th.—No return of hæmorrhage had taken place, but the evacuations were black. Infusion of roses with magnesia was given three times a day.

January 8th.—She left the hospital, convalescent.

The previous symptoms of dyspepsia in this instance led to the opinion that some ulceration of the stomach probably existed, and the excitement of the circulation, from considerable muscular exertion in running, caused the rupture of minute vessels. With quietness, rest, and spare diet, she speedily convalesced.

CASE XCVIII.—*Hæmatemesis, vicarious Menstruation, aggravated Hysteria, simulating Fever.*—Mary H—, æt. 19, was admitted into Guy's under my care in May, 1855. She had enjoyed good health till she was sixteen years of age, when she said that she had a convulsion followed by "brain fever;" and on recovery began to vomit blood three days successively at regular monthly periods; if this did not occur she had pain between the shoulders, at the epigastrium, and dyspnoea; this vomiting of blood continued regularly for three years, but she never menstruated properly. For nine months the discharge had ceased altogether, and three months before admission she had a severe hysterical or epileptic fit.

On admission she appeared stout, tolerably nourished, but prostrate; the tongue was dry and brown, and almost black; she lay motionless in bed, without speaking, and altogether refused food, sometimes groaning, and if taken from her bed appeared to faint. She complained of pain at the lower part of the back, and in the inguinal region; the abdomen was tympanitic and distended: she stated that surgeons had twice removed clots of blood from her; but my friend and colleague, Dr. Oldham, could find no enlargement, nor disease of the uterus, and believed that an attempt had been made to divide the os uteri. She refused to swallow food, the pulse was feeble and very quick. There was slightly increased antero-posterior curvature of the spine in the lower part of the dorsal region.

Milk was poured into the mouth, and she was *made* to swallow it; in this way a considerable quantity of food was taken.

Galbanum and zinc with aloes and myrrh were prescribed, and the bowels were thoroughly cleared by blue pill with colocynth and henbane, and by enemata of rue or soap. Local depletion was used from the groins by the application of leeches, and afterwards quinine and steel were given with wine, and sparks of electricity taken from the spine; a shower bath was occasionally used.

The stomach retained food, and the patient soon became able to walk, and left the hospital in a few weeks convalescent.

This was one of the most severe cases of hysteria that is ever witnessed; and the disturbance of the stomach and alimentary

canal were no doubt produced by the functional disease of the uterus, aggravated by treatment which I think few obstetricians would approve of. The vicarious discharge of blood from the stomach was not observed during the period she was in the hospital, but is received on the testimony of the patient and her friends.

CASE XCIX.—*Vicarious Menstruation from the Stomach.*—Ellen H—, *et.* 23, was admitted under my care into Guy's Hospital, August 28th, 1860. She was a needlewoman, who had resided at Kingsland, and she had for several years been in feeble health, complaining of pain at the chest, and palpitation of the heart, &c. Menstruation commenced when she was eighteen years of age, but the function had been irregularly performed, sometimes ceasing for four to eight months. For six months prior to admission, she had vomited blood at her menstrual periods, but occasionally she had menstruated regularly, and no hæmatemesis then took place; before the attacks of hæmorrhage, and for several days before menstruation, she had pain at the stomach and in the right side, loss of appetite, and nausea; the vomiting of blood then came on and continued for several days. She was a spare, and somewhat anæmic woman, with an anxious expression of countenance, and rather dark complexion; her symptoms were those of amenorrhœa with dyspepsia; and during the time that she remained in the hospital she complained of pain at the scrobiculus cordis, and sometimes also at the right side, and the pain was increased by food; there was occasional nausea, but no vomiting; the bowels were regular. Her general health was improved by preparations of steel and quinine, with a carefully regulated diet; but during the time she was in the hospital there was neither return of hæmorrhage from the stomach, nor was menstruation established. She continued under my care as an out-patient, and was afterwards re-admitted, suffering still from dyspepsia; but she had had no return of hæmorrhage. The dyspepsia was again relieved.

In this case there was probably ulceration of the stomach, and we regard a periodical congestion of the mucous surface, and consequent rupture of minute capillary vessels, as the cause of the repeated hæmorrhage. Although during the time the patient was under observation there was no hæmatemesis, we have no reason to doubt her very positive assertion that for several months she vomited blood at the ordinary period of menstruation, and the discharge was in that respect evidently vicarious. Normal menstruation has since been established, and has continued regularly, but she has suffered occasionally from severe dyspepsia.

CASE C.—*Typhus Fever; Hæmatemesis*.—Ann M—, æt. 19, a hawker in the streets, was admitted into Guy's Hospital, February 2nd, 1859, and died February 4th. Fever existed in the house where she lived, and she had been ill for nine days. When brought to the hospital she had the symptoms of fever, with great depression, and with mottling and lividity of the skin; still she was rational. On the evening of the 4th, the eleventh day of fever, vomiting of blood took place, and blood was passed per rectum; in a few hours she died.

On inspection, the lungs were healthy; the heart was firm but empty; the stomach was full of blood; and at the lesser curvature were several minute depressions or erosions affecting only the surface, and not penetrating the entire thickness of the membrane; one, a little deeper than the rest, appeared to have a minute vessel at the base; but this fact could not be satisfactorily established. The duodenum was reddened and congested; and blood was found in both the jejunum and ileum: and the colon was also full of blood. No ulceration nor disease of the intestine was found; the spleen was large and soft; but the mesenteric glands, as well as the liver and kidneys were healthy.

This instance of hæmorrhage could not be regarded as precisely analogous to those which sometimes occur when the character of the blood is changed, as in yellow fever, purpura, &c., for erosions existed in the stomach from which the blood escaped; but, the prostration of fever doubtless rendered the hæmorrhage more persistent, and perhaps had an important influence in determining the minute ulcerations. The patient was about the age at which perforating ulcer sometimes occurs. In her case, the loss of blood led to a rapidly fatal issue.

PAIN AS A SIGN OF DISEASE OF THE STOMACH.

The two symptoms which are regarded as especially indicative of disease of the stomach are, perhaps, more than any other liable to mislead; we refer to *vomiting* and to *pain* in the region of the stomach; and we shall briefly enumerate the causes of these symptoms as the best safeguard against error. The explanation of the uncertain diagnostic value of these symptoms is found in the intimate connexion of the nerves of the sympathetic plexus with all the abdominal viscera and with the spinal nerves; and, secondly, in the extensive

distribution of the pneumogastric nerve, which supplies, in the abdomen, not only the stomach, but the duodenum, the liver, the pancreas, the kidney, and the supra-renal capsule; and, in the chest, the same nerve extends to the lungs and respiratory tubes, and communicates with the cardiac ganglia. And, again, it is frequently found that irritation of one set of branches of nerves manifests itself in the disturbed function of another part supplied by the same nerve, and that disease at the central origin of the nerve is shown at the peripheral branches; thus, pain in the ear is produced by a decayed tooth, the branches of the fifth pair of nerves supplying both the tooth and the ear; disease at the origin of the pneumogastric nerve in the brain, or of the pulmonary branches, is often manifested by a disturbed condition of the filaments supplied to the stomach, and vomiting is the result.

As a sign of disease pain is of doubtful value: oftentimes it is a certain guide to the locality, if not to the character of the morbid action; at other times, on the contrary, its presence misleads, or its absence disposes us to under-estimate changes which may be going on in the system. Generally speaking, we find that the mucous membranes, except where they approach the outlets of their respective canals are free from ordinary sensibility, and may undergo very marked changes in their condition without any painful manifestation. Acute disease may take place in the mucous membrane of the small or large intestine, in the mucous membrane of the kidney or bladder, with complete immunity from suffering. A similar fact is observed in relation to the parenchymatous viscera; thus the substance of the liver and the kidney is often changed in a marked degree; and if disease, as abscess, form in their structure without much distension, the patient may be unconscious of morbid change. On the contrary, in serous membranes an opposite condition is found to exist; almost any change is appreciated, and in sudden or acute disease the pain is often extremely severe in its character. All well know the stabbing pain of pleurisy, the agony of acute peritonitis, and the intense suffering of severe synovitis. In each of these latter diseases rest is a very essential element in the alleviation of the malady, and this rest can be attained to a great extent without the

cessation of life. In pericarditis, on the contrary, we find, as for many years shown by Dr. Addison, that there is an absence of pain, unless there be pleurisy occurring at the same time. In the pericardium, however desirable rest may be, movement must continue as long as life lasts.

In reference to pain as an indication or non-indication of disease we have to remark—

I. That acute inflammation and disease of the stomach may exist, with entire freedom from pain, if the mucous membrane only be affected. Acute gastritis is generally regarded as an exceedingly rare form of disease, excepting when produced by irritant poisons. This may be the case; but we are of opinion that in many instances the absence of pain has led to this belief. In the gastro-enteritis of children, and not very unfrequently of more advanced life, conditions of great irritability with cessation of the right functions of the stomach, and probably with hyperæmia, must be regarded as closely approaching the character of gastritis. However this may be, we have evidence from the action of irritant poisons that, while the mucous membrane is only affected by them, pain may be entirely absent, excepting that consequent on the violent muscular action exerted in the act of repeated vomiting. Thus in a patient who had taken a large dose of oxalic acid, violent vomiting was produced, failing pulse, and a sense of exhaustion, but no pain. In a few days after taking demulcent forms of diet, she completely recovered. Case XL was an instance of poisoning by sulphuric acid, in which a large portion of the mucous membrane of the stomach was destroyed, and although the patient survived eleven days, she did not appear to suffer from any pain at the stomach. The same fact was still more strikingly shown in Case XLI—when chloride of zinc had been taken, life was prolonged for three months. But the absence of suffering was remarkable till eight days before death; and the pain then, we do not doubt, was due to the formation of an abscess in the left hypochondriac region. I have witnessed the same immunity from suffering in poisoning by arsenic, and by corrosive sublimate; and we are, I think warranted in the belief that acute disease may take place in the mucous membrane of the stomach without any pain.

II. Organic disease of the *mucous* membrane, as for instance cancer, may be comparatively free from pain. It frequently happens in cancerous disease of the liver, that after death tubercles or growths of similar character are observed on the mucous membrane, and of which there had been no indication during life. Thus, in Case LXXIX, a patient, aged sixty, died from cirrhosis, and after death a large villous growth was found attached to the anterior surface of the stomach, but the orifices were free. After admission, there was no complaint of any pain at the stomach, neither had she any vomiting; and it is probable that the burning pain she had before admission was of the character often observed in ordinary dyspepsia, for she was of intemperate habits. The freedom from any obstruction at the orifices, and the growth involving only the mucous membrane, was, we think, the cause of the absence of pain. No supposition was entertained of the presence of this growth in the stomach during life.

III. Disease extending to the muscular or peritoneal coats produces generally severe pain, as observed in ordinary ulceration or cancer. This symptom is present as one of the most ordinary signs of the conditions just mentioned, often coming on directly after food has been taken. In several instances, in which the suffering was exceedingly intense, we have found branches of the pneumogastric nerve involved in the thickened, dense, and fibrous edges of the ulcer. In a case of this kind which I watched with much interest, the cause of death was gradually increasing exhaustion, as the consequence of the intense pain and constant vomiting. In another, Case LIII, a young woman, aged twenty-one, suffered from constant and severe pain, with progressive emaciation, continuing for many months, unrelieved by the administration or application of any anodyne that we possess. A month before death symptoms of acute phthisis came on. A large chronic ulcer was found at the lesser curvature, and several branches of the pneumogastric nerve were traced to the edges of the ulcer, and some passing across its base were only covered by fibrous tissue.

IV. Over-distension of the stomach produces severe pain. The formation of the stomach and its peritoneal attachments are such as to allow moderate distension in the performance of

ordinary digestion ; but whenever the distension becomes greatly increased, pain is the result.

V. Disease, especially of an acute kind, affecting the peritoneum, is also, with few exceptions, accompanied by severe pain. In reference, however, to the position of pain in peritonitis, it is not always a certain guide to the precise seat of injury. I well remember a patient under the care of the late Dr. Golding Bird, a young woman, who was seized with sudden severe pain at the scrobiculus cordis and towards the left side, followed by rapid collapse. From the seat of the pain he was led to diagnose perforation of the stomach ; it was, however, found to be perforation of the appendix cæci.

VI. Dr. Osborne has shown, that in some cases of gastric ulcer the position of greatest ease to the patient may serve as a guide to the exact seat of the disease ; that if the ulcer be on the posterior surface, lying upon the face would be the most comfortable position, and *vice versâ*. Food, on its entrance into the stomach, generally passes directly along the lesser curvature ; and if the viscus be contracted, would come in contact with an ulcer, whether placed on the anterior or posterior aspect of the median line of the curvature. If more distended, there might be less direct application to the diseased surface ; in the case of severe suffering from gastric ulcer previously referred to, the patient appeared to be most easy when leaning somewhat forward and towards the left side, which would have the effect of allowing fluids to gravitate from the ulcer, as mentioned by Dr. Osborne. We have seen several cases which tend to confirm this opinion.

VII. In disease of the lesser curvature, near the pyloric orifice, pain is sometimes experienced by the patient as soon as the food enters the stomach, and, in some cases, this conveys the idea of disease at the œsophageal orifice. This fact may lead to the supposition that the œsophagus is the part affected, and the opinion may be strengthened by the rejection of food almost before it has reached the stomach, as in Case LXXXVIII.

VIII. Many conditions of functional disease are entirely free from pain. It is, indeed, well for us that there is such insensibility, otherwise the least deviation from healthy action

might be followed by suffering, and the strict rules of a dyspeptic be essential in ordinary life.

IX. The pain, in many functional diseases of the stomach, is exceedingly severe; but it is often produced by a mal-condition of the nerves or nerve-centres, and it arises from the intimate connection of the spinal and sympathetic nerves. In some states of exhaustion the whole of the nervous system appears to be in a state of great irritability, and the sensibility of structures becomes greatly increased. We often find, in these conditions, the stomach is incapable of bearing the presence of food; it is at once rejected, or produces intense pain, or flatulent distension is immediately set up, or a sense of fainting; and the means best calculated to relieve are those which invigorate and strengthen the whole system. Of this class are the stomach diseases observed in connection with uterine disease, with loss of blood, exhaustion, mental anxiety, &c. The deficient nervous supply also interfering, perhaps, with the right secretion of the gastric juice.

X. The effect of a diseased condition of the pneumogastric nerve at its centre, or at its peripheral branches, in connection with stomach disease, is of great interest, and it is probable that pain is sometimes the result. We have, however, more frequently observed vomiting rather than pain produced by an irritated condition of the pneumogastric nerve.

XI. In some forms of functional disease of the stomach, in which severe pain comes on three or four hours after food, it is probable, as we have elsewhere stated, that extreme irritability of the pyloric orifice exists.

XII. In functional, as in organic disease, pain often arises from distension of the stomach, consequent on chemical decomposition of the alimentary mass.

XIII. The absence of pain sometimes arises from the destruction of the pneumogastric nerve. This fact was remarkably shown in Case XXVIII, the patient suffering from sloughing at the extremity of the œsophagus; and in cancerous disease of the stomach the same thing has been observed.

XIV. Pain at the scrobiculus cordis, simulating disease of the stomach, often arises from spinal disease, the pain being referred to the extremity of the irritated nerve.

XV. Severe pain at the *scrobiculus cordis* is frequently present in chronic bronchitis and in obstructive valvular disease of the heart; in fact, from any state which leads to over-distension of the cavities on the right side of the heart. In these conditions we very generally find that food produces pain and flatulence, and is very imperfectly digested; the vessels of the stomach and of the whole of the chylopoietic viscera are much engorged, and the surface of the stomach is very generally covered with a thick layer of mucus, a state of chronic catarrh of the gastric mucous membrane is produced. Many observers, however, attribute the almost constant pain at the *scrobiculus cordis* in these instances to the over-filled cavities of the right side of the heart, and we are disposed to refer part of the distress to this cause.

XVI. In aneurism of the abdominal aorta we have sometimes observed pain of a most intense kind, and the disease might very readily have been mistaken for cancerous disease of the stomach with glandular infiltration, producing pressure upon the aorta. In one instance, which I watched with much interest, the aneurism existed at the position of the *coeliac axis*; it was rightly diagnosed, and the patient became exhausted and died from the intensity of the pain, the false sac not having given way. I dissected large branches of the sympathetic nerve spread out upon the surface of the tumour; and the intense suffering and fatal exhaustion appeared to arise from the implication of the nerve structures; no other cause of death could be found on very careful inspection.

Enough has been said to show that the most careful investigation of this symptom is necessary to form a correct diagnosis of disease of the stomach.

VOMITING AS A SIGN OF DISEASE OF THE STOMACH.

The causes of vomiting are still more varied than those of pain at the stomach; and the importance of carefully estimating these causes is in proportion to their complexity; and although some of them are not connected with gastric disease, we still make brief reference to them. These causes of vomiting may

be divided into those which originate in the stomach and intestines, and secondly, into those which are sympathetic.

In the first division we must place—

1. Inflammation of the stomach ; gastritis and gastro-enterite.
2. Undigested food, or foreign bodies in the stomach.
3. Irritants and medicines.
4. Great irritability of the mucous membrane.
5. Ulceration of the stomach.
6. Obstructive disease of the pylorus.
7. Cancerous disease.
8. Acute peritonitis.
9. Pressure on the stomach, as in ascites and ovarian dropsy, in abdominal tumours, &c.
10. Diseases of the duodenum.
11. Hernia, intestinal obstruction, intussusception.
12. Pharyngeal and œsophageal regurgitation.

In the second division are—

1. Diseases of the liver and gall-bladder.
2. Diseases of the kidney.
3. Diseases of the supra-renal capsules.
4. Diseases of the uterus and ovaries.
5. Diseased conditions of the blood and general nervous system, as at the onset of the exanthems, fevers, pyæmia, erysipelas, &c.; ague, yellow fever, and cholera may, perhaps, be classed among these as arising from blood change.
6. Diseases of the spine.
7. Diseases of the brain.
8. Diseases of the lungs.

I. There is something remarkable in the presence of vomiting in circumstances where pain is absent ; thus, in acute disease of the stomach, where only the mucous membrane is affected, the patient may be free from all suffering at the region of the stomach, excepting that produced by the violent straining of the muscles during vomiting. We need not do more than refer to the instances of poisoning by oxalic acid, by sulphuric acid, by arsenious acid, and by corrosive sublimate, which have been already given as illustrations of this fact ; and in the symptoms

of gastro-enteritis the same immunity from gastric pain occurs, whilst vomiting greatly distresses the patient.

II. Undigested substances often remain in the stomach for some time without producing pain, unless they pass within the pyloric valve; and we sometimes find that they are retained for many hours or even days before they are rejected by vomiting.

III. In reference to vomiting caused by medicine and by irritants, it is only necessary to mention that in some instances the action appears to be one of primary irritation of the stomach, in others it is secondary, through the medium of the blood; but whether this secondary action and its consequent vomiting arise from the excretion of the medicinal substance from the mucous membrane of the stomach is doubtful; thus tartar emetic produces vomiting when injected into the blood equally as when taken directly into the stomach.

IV. We have referred, in our remarks on functional disease of the stomach, to states of extreme irritability of the mucous membrane in which food of every kind is at once rejected. This form of functional irritability we have found generally associated with uterine and ovarian disease; or it has been produced, apparently, by irritation of the pulmonary branches of the pneumogastric, to which we shall presently have to refer; but in some instances we have not been able positively to trace the complaint to one or other of these causes, and at present we must acknowledge, though unwillingly, as a cause of vomiting, functional irritability of the stomach itself. It is to this form of disease that Sir H. Marsh has given the name of regurgitative disease, in which food is rejected *without any effort*, and often without *corresponding emaciation*. In his valuable paper on this subject, he refers to its connection with pulmonary and with uterine disturbance.

V. In ulceration of the stomach, vomiting often comes on as soon as food enters the stomach, or a period of variable length intervenes, the pain increasing till the rejection takes place.

VI. In obstructive disease at the pylorus, the vomiting is generally deferred till nearly the close of the digestive process; much, however, may be done to diminish this symptom by the administration of fluid diet, so that sometimes several days elapse between the attacks.

VII. *Cancerous disease* affecting the orifices of the stomach constitutes a common cause of persistent vomiting. It must, however, be borne in mind that vomiting is not a constant sign of cancerous disease of the stomach; if the orifices be free, it may be entirely absent, although the disease is very extensive; and again, if sloughing take place, even when the orifices also are diseased, vomiting often subsides, sometimes in consequence of the obstruction being removed by the sloughing; at other times, apparently, from the destruction of the branches of the pneumogastric nerve. Further, the period at which vomiting occurs does not always indicate the seat of the cancerous obstruction; in some instances of obstruction at the pylorus, with disease at the lesser curvature, vomiting takes place immediately after food has entered the stomach, so as to convey the idea of obstruction at the cardiac orifice or in the œsophagus itself, and the symptom has been regarded as dysphagia rather than vomiting. Case LXXXVIII furnishes a good illustration of the truth of this remark.

VIII. *Acute peritonitis*, especially when the gastric peritoneum is involved, is often accompanied with severe vomiting. The statement has been made, that vomiting does not take place in acute peritonitis unless the peritoneum in the neighbourhood of the stomach is implicated; but although this is generally, it is not constantly the case. Chronic peritonitis is also a cause of vomiting; so also local peritonitis and effusion near the stomach. In some of these instances the stomach is affected by its direct implication in the disease; in others vomiting arises from the pressure of effused pus or the constriction of adhesions.

IX. *Pressure* on the stomach is a direct cause of vomiting. In *ascites* and *ovarian dropsy*, the stomach is sometimes so compressed that vomiting comes on soon after food has been taken, apparently from this cause alone; and when paracentesis has been performed, the pressure being removed, the sickness ceases. When glandular tumours in the neighbourhood of the pancreas or disease of the pancreas itself exert pressure on the stomach, the symptoms closely resemble primary disease of the stomach, and the diagnosis is exceedingly difficult; but, since the pancreas receives a branch from the pneumogastric nerve, it is not easy to ascertain how far vomiting in some of these cases is due to nervous irritation, and how far it is due to direct pressure. In

those cases in which the pancreas has been diseased, without great enlargement and without pressure on the stomach or duodenum, I have not observed vomiting as a prominent symptom. Case CXIV was a very remarkable instance of this kind, and presented many of the symptoms of organic disease of the stomach. Sloughing of the pancreas had taken place, and a large abscess had been formed, which exerted pressure upon the duodenum.

In *aneurismal disease* of the abdomen, the remark which we made in reference to disease of the pancreas and its glands holds good; and the same difficulty arises in determining how far the vomiting is due to pressure or to sympathetic irritation.

In some cases we have found direct pressure made by the patient at the *scrobiculus cordis* the cause of vomiting; many persons can thus at once empty the stomach; and in an instance of a boy, some years ago, in Guy's Hospital, it was only after very careful watching that the true character of the complaint and the deceit of the patient were ascertained.

X. As to vomiting not depending on the condition of the stomach itself, we have to refer to morbid states of other abdominal viscera, and first to disease of the *duodenum*, as inflammation of its mucous membrane, ulceration, and especially obstruction.

There is great similarity between the diseased conditions of the first portion of the duodenum and of the stomach. Ulceration of the first portion of the duodenum produces many of the symptoms of like disease in the stomach. Case CXII, mentioned with disease of the duodenum, presented many points of great interest; old peritoneal adhesion had led to obstruction at the central part of the duodenum, which had become greatly distended and slightly ulcerated; extensive strumous disease of the glands in the neighbourhood of the thoracic duct and of the mesenteric glands had taken place, so that symptoms were produced very closely resembling organic disease of the stomach itself, with which it was confounded; at the same time discoloration of the face rendered it doubtful whether the supra-renal capsules were affected, so great was the prostration; the failing pulse and the vomiting also closely resembled the symptoms found in renal capsular disease. A form of dyspepsia in which vomiting, with pain at the seat of the duodenum, comes on at

the close of digestion has been attributed to the duodenum; but whether this class of cases is connected with an abnormal irritability of the pylorus itself, we cannot affirm. Again, in some cases of acute jaundice, febrile symptoms with violent irritability of the stomach, but without pain, arise; and the disease has been attributed to mischief commencing in the duodenum, and extending to the biliary ducts. In some fatal cases of this kind, great congestion in the duodenum has tended to confirm the idea; so also the fact that symptoms of this kind have come on after intemperance. See Case CVII.

CASE CI.—Acute Jaundice; Violent Vomiting; Recovery.—Samuel H., æt. 22, was admitted into Guy's Hospital under Dr. Barlow's care, March 11th, 1861. He was a cellar-man employed in Tower Street; and he stated that he did not exceed two or three pints of ale during the day. Some time previously he had had gonorrhœa, followed by slight sore throat. He had continued well till March 7th, when he complained of languor and headache, and of a sense of weakness with swelling and pain in the calves of the legs. On the 8th and 9th he had vertigo and pain in the head. On the 10th he had pain in the left side of the chest, and down the arm to the elbow. On the 11th, when admitted, the skin was hot, but *not* jaundiced; the pulse was 96, soft and compressible; the tongue was furred and brown; the bowels were confined; and there was no appetite. There was much headache, but no evidence of disease of the chest. The abdomen was distended and painful; but there was no pain in the region of the gall-bladder nor liver. There was much thirst, and he complained of pain in the legs. There was vomiting, and constipation of the bowels. The urine was scanty, coffee-coloured, and albuminous. He was ordered grey powder, four grains, followed by castor oil and julep of acetate of ammonia with nitric ether, three times a day.

On the 12th, the skin became slightly jaundiced, the bowels did not act from the grey powder and castor oil; and five grains of calomel were given, followed by a senna draught; this dose produced a bilious fluid evacuation. Vomiting of green fluid came on, with pain at the stomach, but not of a severe character. The pulse 92. Effervescing saline medicine was given.

On the 13th, the jaundice was marked; the skin was hot; the pulse slow and compressible. There was mental oppression; he had slight pain in the abdomen, but it was not localised; at the same time he had pain in the shoulders, legs, and thighs; the thigh appeared to be swollen. The tongue was furred and brownish in the centre. The symptoms suggested the idea of pyæmia, and four ounces of brandy and effervescing mixture were given.

On the 14th, *the vomiting was severe*; the bowels were confined; the tongue furred. He had had but little sleep. Fifteen grains of colocynth and calomel were given, and the brandy was omitted.

On the 15th, the bowels were acted on freely, and there was bile in the evacuation. The urine was scanty, but *not* albuminous. The pulse was 80, and compressible; the tongue furred and white. There was less headache, and he had had a few hours' sleep. He was ordered magnesia mixture, with compound decoction of aloes, of each half an ounce three times a day, and soda-water with milk.

On the 18th, the symptoms were relieved; there was less oppression, and the vomiting had ceased; the pulse also was less compressible, and the tongue less furred. He was ordered calomel, gr. ij, compound colocynth pill, gr. ij, ipecacuanha, gr. j, every night.

On the 20th, he had still less pain and cerebral oppression; the skin was less jaundiced; the urine abundant, and less highly coloured; but the motions were clayey. The pulse 76. No vomiting.

On the following day he had a rash on the skin, resembling urticaria.

On the 22nd, the jaundice was disappearing, but the evacuations were still pale and clayey. There was no pain in the abdomen nor limbs, and his countenance appeared more intelligent.

On the 27th.—Two or three days previously, erysipelas of the right side of the face and neck had come on, and suppuration had taken place at the angle of the jaw. The abscess was opened. Julep of ammonia and decoction of aloes, of each half an ounce three times a day, were given; wine four ounces.

30th.—The erysipelas was disappearing; the countenance was pale; the pulse very compressible and slow; the urine clear and of normal colour; the motions pale.

April 30th.—Since last report steadily convalescent, and he left the hospital well.

This case was one of obscurity, and of great interest. At first the action of the kidneys appeared also to be checked; but the irritability of the stomach was excessive, and the prostration typhoid, and for several days the illness was very severe. It was doubtful whether the disease extended from the duodenum to the bile ducts, and thus induced jaundice.

XI. *In hernia, obstructive disease of the intestines, intussusception, &c., vomiting* is generally present. If the obstruction be in the small intestine, the vomiting comes on very quickly; but if the colon, sigmoid flexure, or rectum be the seat of disease, vomiting is often postponed for a considerable time, unless irritant medicines and violent purgatives have been administered. As the vomiting continues, the ejected matters present the character of the fluids at the seat of obstruction; and if that obstruction be intestinal, their odour and appearance have more or less of a fæcal character. The following is men-

tioned to show how easily vomiting may be excited in obstruction of the colon :

CASE CII.—*Old Peritoneal Adhesions ; Twist of the Sigmoid Flexure ; Acute Peritonitis ; Death on the Sixth Day.* (Reported by Mr. Soper).—William B—, æt. 56, a dock-labourer, residing at Deptford, was admitted July 19th, 1861. He had always enjoyed good health till six months ago, when he received a blow on the right side, which laid him up for a fortnight, and he suffered from severe pain. Till July 15th he continued well ; but on that day, whilst at work, he was seized with severe pain across the middle of the abdomen. A dose of castor oil was taken ; no action on the bowels was produced, and the pain continued ; he passed water freely. On the next day, Tuesday, 16th, he repeated the castor oil, with some oil of turpentine, which at once produced *violent vomiting*, without any evacuation from the bowels. These symptoms continued till admission. He was a strongly built, muscular man, with a distressed and anxious countenance ; he was then collapsed, and the pulse scarcely perceptible, 140. The respiration was thoracic, 32. The abdomen was very tense, tympanitic, painful, and partially dull in the umbilical region ; the tongue was furred, and there was desire to go to stool ; the urine was scanty and high coloured ; the breath was warm, but the skin was cold and clammy ; the pupils were contracted. He was ordered opium, gr. ij, at once, and gr. j, every four hours ; the pain subsided, and he had some sleep. On the following day, the 21st, he was evidently dying ; at 1.30 p.m., the hands and feet were cold, the abdomen was tense, tympanitic, but there was no complaint of pain. The opium was continued, and a small quantity of stimulant given. He died at 9 p.m.

Inspection was made on the 22nd. Decomposition was advanced. The abdomen was distended ; the peritoneum was acutely inflamed, intensely injected where the coils were in contact, and the serous membrane was covered with lymph. The cæcum was greatly enlarged, and immediately on its left was a coil of the sigmoid flexure, enormously distended, of green colour, and occupying the greater part of the umbilical and left lumbar region ; the transverse colon was in its usual situation, and also of large calibre. The small intestine was also distended, and situated principally in the epigastric region and in the pelvis. On turning aside and drawing up the small intestine from the pelvis, the sigmoid flexure was found to be twisted on itself and constricted. There was adhesion between the sigmoid flexure and the cæcum, and also between the portions of sigmoid flexure itself. These adhesions had apparently determined the twist ; at the extremities of the twisted coil, the green colour and distension suddenly ceased ; below, passing into the collapsed rectum ; above, into the distended colon. The appendix cæci was intensely congested and inflamed. The other viscera were healthy.

The blow on the right side possibly produced the peritoneal adhesions found after death, and led to the contraction which

determined the twist of the sigmoid flexure. No plan of treatment would probably have been effective in this case; but the purgative medicine, the turpentine, and castor oil given before admission, at once set up violent vomiting, and produced the peritonitis at an earlier period than would otherwise have taken place. Death ensued on the sixth day; whereas if no purgative had been administered, but opium and occasional injections used, life might perhaps have been prolonged for three or four weeks, or even for a longer period.

XII. The regurgitation of food, rather than vomiting, which is consequent on disease of the pharynx, larynx, or œsophagus, must be distinguished from actual vomiting. By carefully observing the process of deglutition, the seat of mischief may be accurately ascertained. In paralysis of the muscles of the soft palate and of the pharynx, deglutition cannot be properly completed, and food is rejected through the nares; so also when the epiglottis is ulcerated from strumous, syphilitic, or cancerous disease, the act of deglutition is scarcely performed before the substance swallowed is violently ejected, and severe pain in the throat, and cough, are set up. It is remarkable too, in these cases, how a solid bolus of food may be formed and swallowed, slipping beyond the diseased surface, whilst the smallest quantity of fluid produces most violent pain and distress. In obstruction of the œsophagus, the act of deglutition is completed, and then regurgitation takes place. Very extensive disease may, however, affect the œsophagus without this rejection of food; for ulceration or sloughing may have removed obstruction, or the branches of the pneumogastric nerve and the whole wall of the canal may be destroyed. Other causes of vomiting are manifestations of the general and intimate connection of the stomach with other viscera; they are properly designated sympathetic in their origin, and their study is of essential importance in the diagnosis of disease of the stomach. As with the other causes of vomiting we shall do little more than enumerate them; and the first of this class to which we allude is—

XIII. Disease of the *liver* and of the *gall-bladder*; large branches of the pneumogastric nerve extend to the liver, as well as numerous nerves from the large sympathetic ganglia. In

gall-stone, violent vomiting is generally associated with the intense pain; and in many conditions of hepatic disease the same symptom is constantly present. The following case was one of great interest.

CASE CIII.—*Violent Vomiting; Gall-stone (?) ; Pleuro-pneumonia.*—James R., æt. 42, was admitted into Guy's Hospital about 1857, under Dr. Barlow's care, with severe vomiting; there was pain in the region of the stomach as well as of the duodenum, and slight jaundice came on. In several weeks he recovered, but the symptoms returned, and he was again admitted; a third time they came on, and a third time subsided. The fourth time on which he applied, September 12th, 1859, the same condition of stomach existed; and the pain in the region of the duodenum had come on as before, but there was no jaundice. A short time before death, symptoms of acute pleuro-pneumonia of the right lung supervened. On inspection the brain was found in a healthy state; the right pleura contained a thin layer of lymph. The lower lobe of the right lung was of a dusky gray colour, soft, very offensive, and gangrenous; the upper lobe was generally more dense than normal, many lobules were dense, and some were consolidated from deposit. The left lung was congested. There was slight enlargement of the bronchial glands; but no evidence existed of pressure or disease of the pneumogastric in the brain, neck, or chest. The heart was healthy. All the abdominal viscera were sound; but along the greater curvature of the stomach some patches of arborescent injection were present; and in the first part of the duodenum there was considerable redness of Brunner's glands, which were unusually prominent. There was no evidence of disease of the liver, gall-bladder, spleen, pancreas, supra-renal capsules, nor kidney.

In this case the cause of the vomiting was obscure; it was supposed to have arisen from gall-stone, and, as far as could be ascertained after death, this was the case. The pain in the region of the duodenum, and the congested state of the mucous membrane at that part, were not inconsistent with this idea. The acute attack of pleuro-pneumonia which came on whilst the patient was in the hospital, and which rapidly passed into gangrene of the lung, led to the supposition that the pneumogastric nerve had been the cause of the symptoms; but no source of irritation in the brain, nor in the course of the nerve, could be detected. The passage of gall-stone is sometimes followed by chronic change, thickening, and the effusion of fibroid tissue, and may in some cases determine the seat of the deposition of cancerous product; and it is in these cases that the diagnosis is

sometimes obscure, and the correct pathological interpretation of the symptoms difficult.

XIV. In disease of the *supra-renal capsule*, vomiting is rarely absent; but sometimes it is a sign of such prominence as to simulate primary disease of the stomach. On post-mortem examination we have found arborescent injection of the mucous membrane of the stomach, and sometimes slight superficial ulceration; but it must also be remembered that the pneumogastric nerve affords a branch to the supra-renal capsule, and that its connection with the semilunar ganglia is a very intimate one. In Case XLIII, the vomiting with prostration was the most prominent symptom. The patient had been on the western coast of Africa, and red dusky blotches on his bronzed face were supposed to be due to syphilis, which he had contracted sometime previously; so that although disease of the supra-renal capsule was mentioned at his bedside, still the case was thought to be one of disease of the stomach. In Case XLVI, also, the irritability of the stomach was a marked symptom, and after death superficial ulceration was detected.

XV. Diseases of the kidneys and renal calculus constitute other causes of vomiting. During the passage of a calculus down the ureter, vomiting is a very distressing symptom. In acute albuminuria vomiting is also associated with nausea; and in chronic albuminuria it is sometimes the precursor of a fatal termination. So severe, indeed, may be this symptom in *ischuria renalis*, as even to suggest the possibility of intestinal obstruction, as shown by Dr. Barlow. The vomiting in albuminuria is not only due to the direct connection of the nerves constituting the renal plexus with those of the stomach, but to the urea excreted from the mucous membrane of the stomach and intestines. It is found to be present in large quantity in the blood, and is separated in all the excretions and secretions; and in the stomach this abnormal excrementitious substance appears to act as a direct irritant.

XVI. Both functional and organic diseases of the uterus are causes of vomiting. In dysmenorrhœa, most distressing irritability of the stomach is occasionally set up; and in pregnancy, vomiting may be so severe as to exhaust and completely to prostrate the patient.

XVII. The remaining causes of vomiting arise from the condition of the nervous system, and are most interesting and important in the correct diagnosis of disease; the first of these is diseased condition of the *spine*. The splanchnic nerves pass from the spinal cord to the large sympathetic ganglion of the abdomen, and constitute an intimate connection between these centres of nerve-force; in those diseases, however, of the spine in which we have observed irritability of the stomach, other sources of disturbance have been present.

XVIII. At the onset of *acute diseases*, especially the exanthems, fevers, pyæmia, erysipelas, &c., vomiting is often present. It is not known how this is produced, whether directly by the altered condition of the nervous system, or secondarily from the state of the blood. Sudden nervous shock, fright, &c., will produce vomiting; and in some more chronic diseases, when the blood is altered in character, as in renal disease and even gout, the same symptom is occasionally very intractable.

Dr. Graves, in his 'Clinical Medicine,' makes the following valuable remarks in reference to this subject:—"Every fever which commences with vomiting and diarrhoea, whether it be scarlatina, or measles, or typhus, is a fever of a threatening aspect; and in all such fevers the practitioner should be constantly on the watch, and pay the most unremitting attention to the state of the brain. There is much difference between the vomiting and diarrhoea of gastro-enteritis, and this *cerebral diarrhoea and vomiting*. The latter sets in generally at a very early period of the disease, perhaps on the first or second day, and is seldom accompanied by the red and furred tongue, the bitter taste of the mouth, the burning thirst, and the epigastric tenderness which belong to gastro-enteric inflammation." He also states very truly, that in cerebral disease there is often a large quantity of bile rejected by vomiting, and passed also by stool; and that leeching the abdomen is less efficacious in cerebral inflammation than in gastro-enteritis.

Very little is known as to the proximate cause of vomiting in cholera and in yellow fever; but we sometimes find in the intermittents of our own country that vomiting is a prominent symptom; and we have several times witnessed instances in

which vomiting, excited possibly by uterine or hepatic mischief, assumed regular periodicity in those who had been exposed to miasmatic poison. The following case was of that kind:

CASE CIV.—*Anæmia. Struma. Vomiting and Diarrhœa every other day.*—Ellen M—, æt. 26, living at Kennington, was admitted into Guy's under my care. For two and a half years she had been married, but six years previously resided in a low marshy locality in Cambridgeshire, as lady's maid; while in that capacity loss of power and vomiting obliged her to give up her situation, and she returned to her native place, Lynn, in Norfolk.

Her father and sister died of phthisis; and she appeared to have had delicate health, being subject to syncope.

She had had two children, and the youngest was, at the time of her admission, seven weeks old. In her first pregnancy she was much prostrated by vomiting, and was unable to take animal food; after parturition she partially recovered her strength, but in a month the sickness and faintness returned; on again becoming pregnant she lost these symptoms, and felt well till the seventh month, when they returned with such severity as to oblige her to keep her bed a month before delivery. There was but little blood lost in labour, but on the sixteenth day she was again seized with vomiting, which gradually increased in severity; there was diarrhœa at the same time as the sickness, and these symptoms came on every other day, on the intermediate days the patient feeling cold and chilly.

She was a spare woman, much emaciated. At night she perspired, and had disturbed sleep; the bowels were relaxed, she had the disposition to take food, but the stomach rejected it soon afterwards; pulse sharp, compressible, &c. There was systolic bruit over the region of the aortic valves; no physical sign of pulmonary disease existed, but she had a slight cough; the urine was pale and normal.

In this case, symptoms of irritability of the mucous membrane of the alimentary canal were brought on apparently by sympathetic connexion with the uterus. It came on first during pregnancy, ceased for a short time after labour, and with her second child the same symptoms occurred. Sickness is one of the ordinary symptoms of early pregnancy, but here this irritability had been perpetuated by some unusual susceptibility of the mucous membrane, or of its nervous centres, and aggravated by a strumous diathesis.

This state is sometimes symptomatic of organic disease of the mucous membrane of the uterus, but we had no proof that such was the case here.

There was remarkable periodicity, a kind of tertian irritability

of the stomach and intestines existed ; on every other day vomiting and diarrhœa, alternated with coldness and chilliness. Long residence in a miasmatic district had probably given this periodicity to the symptoms, and aggravated her anæmic condition.

As to the treatment : the diarrhœa was checked by starch enemata with opium ; and steel in the form of the ammonia tartrate of iron was prescribed ; but quinine gave more relief, at first combined with trisnitrate of bismuth and conium in the form of pill, afterwards with steel. The irritability of the stomach subsided so much, that she was shortly able to take animal food ; the countenance became less blanched, the spirits less depressed, the diarrhœa almost disappeared, and the systolic bruit became inaudible ; by a perseverance in the use of these means she continued to improve, and left the hospital convalescent.

XIX. Irritation of the peripheral branches of the pneumogastric nerve in the abdomen has already been referred to as one cause of vomiting in disease affecting the organs to which they are supplied ; but the same nerve may be changed at the peripheral branches in the chest, and at its origin in the brain. *Disease of the brain*, then, is another cause of vomiting, and one which it is important to bear in mind in the diagnosis of disease ; too often the so-called bilious attacks of children are the first indications of acute hydrocephalus. The irritability of the stomach is sometimes so great, that vomiting is at once produced when the patient is raised from the recumbent position. The diagnosis of these cases is sometimes exceedingly difficult when commencing with symptoms of true gastro-enteric disease ; but it would be well if the remark of the great authority in clinical medicine just quoted were borne in mind, that, "*in all feverish complaints, where during the course of the disease the stomach becomes irritable without any obvious cause, and where vomiting occurs without any epigastric tenderness*, you may expect congestion or incipient inflammation of the brain or its membranes." In simple cerebral disease the abdomen is generally collapsed ; in primary abdominal disease there is, on the contrary, distension. This difficulty in diagnosis is not, however, limited to very young subjects. In strumous disease of the brain, the vomiting is sometimes excessive ; the following

case was of this kind: a state of chlorosis and anæmia was associated with disease of the cerebrum and fatty degeneration of the heart; and it was doubtful how far the vomiting was due to uterine inaction.

CASE CV.—Chlorosis. Vomiting. Tubercles in the Brain. Fatty Degeneration of the Heart.—Maria J—, æt. 18, a needlewoman, who had resided at Woolwich and Chatham, was admitted under my care into Guy's Hospital, May 22nd, 1861. Two years previously, after exposure to cold, she ceased to menstruate, and there had been no return. She had had cough, headache, and pain in the loins; the glands in the neck had slightly swollen. She was admitted in a very anæmic state, blanched, and she complained of headache, and palpitation at the heart; the pulse was sharp, the pupils dilated, and there was a slight systolic bruit in the cardiac region. She was ordered quinine mixture, with sulphate of iron. This, however, she was unable to retain on the stomach, and tincture of iron was given. The vomiting became very violent, and green fluid was ejected; at the same time there was soreness and fulness of the abdomen, and increased prostration of strength. Mr. Stocker ordered brandy and carbonate of magnesia, with compound tincture of lavender.

On the 25th the vomiting and prostration continued as before; and chloric ether, with carbonate of soda in mucilage mixture, were given every four hours, and the brandy continued.

On the 29th she was very prostrate and anæmic; the vomiting continued; the abdomen was rather distended, and there was slight tenderness at the *scrobiculus cordis*; the bowels were loose, the pulse compressible; there appeared to be a listlessness of manner, but it was considered to be due to great prostration. She was ordered ice and brandy. To repeat the mixture, with a small quantity of opium, and to have an enema of starch with opium. At a quarter to nine on the same evening she died unexpectedly. The body was tolerably nourished. *Brain.*—The cerebellum contained in its left lobe several large strumous masses, about half an inch in diameter, extending from the gray into the white substance; there was a small quantity of lymph beneath the arachnoid, in the neighbourhood of these tubercles, and there were numerous points resembling purpuric extravasation in the meshes of the arachnoid. In the *chest* there were old pleuritic adhesions; and at the apices of the lungs were several tubercles; the lower lobes were œdematous. The muscular fibre of the *heart* was pale and soft, and in the right ventricle were many grayish-white points of *fatty degeneration*. On the abdomen there was a moderate layer of fat. The peritoneum was quite healthy. The omentum, with a moderate layer of fat, was spread out over the intestine. The stomach was contracted; towards the lesser curvature there was minute arborescent injection, but the mucous membrane was generally pale. The liver was pale, the spleen healthy; several spleniculi were found. The kidneys and supra-renal capsules were healthy.

In this instance the cerebral symptoms were obscure; and

the pain was not more severe than is often observed in simple chlorosis; neither was the vomiting a distressing symptom till two or three days before death. The pupil was dilated, and the patient was quite sensible till within a short period of the close of life; the pulse, also, had the sharp character generally found in states of anæmia. It was not till the vomiting became excessive that the great prostration ensued; and the degenerated condition of the muscular fibre of the heart had probably an important effect in leading to the sudden termination of life.

After concussion of the brain, vomiting comes on; and in some cases, when inflammatory disease has followed, and suppuration has taken place, the vomiting is excessive. One of the most severe cases of secondary vomiting which I ever witnessed was of this kind. A man in middle life had received a blow at the back of the head; cerebral symptoms came on, and suppuration took place at the origin of the pneumogastric nerve; the membranes were adherent at that part for the space of half an inch, and about half a drachm of pus was effused. The vomiting was excessive; anything swallowed was rejected with violence beyond the extremity of his bed.

The following case appeared to be one of *membranous irritation, with diseased ear*. The symptoms have twice subsided under treatment.

CASE CVI.—*Vomiting (cerebral). Diseased Ear*.—George C—, æt. 30, was admitted under my care, May 23rd, 1860. He had resided at Woolwich, as a stoker at the public baths. His mother had suffered from gout; some of the family had died from phthisis, others from disease of the heart. He had been married for ten years; eight years previously he had ague in the Crimea, and for eighteen months was at Scutari with dysentery. After his return to England he became very intemperate in his habits, drinking as much as a pint of brandy daily, and for more than a week before his illness he had taken more than usual. A fortnight before admission, whilst at work, at 9 a.m., he experienced a sudden pain at the back of the head, as if he had received a severe blow from a hammer; at the same time he had much pain on trying to turn his head, and he was carried home. He tried to work the next day, but again was carried home and placed in bed. Sickness came on the same day, the stomach rejecting everything, and this irritability of the stomach continued till admission; the bowels were acted on after the commencement of the attack. Eight days before admission he had a discharge of blood and pus from the left ear, which lasted three or four hours, with constant tinnitus aurium; and he had slight pain over the mastoid process. He was a spare man, of middle height, light complexion; both conjunctivæ

were in an inflamed condition, and had a purulent discharge from them; the pupils were of natural size; the gums were ulcerated (mercury?); the tongue had a thick brownish fur; the skin was cool. There was no evidence of thoracic disease; the pulse was 48, and labouring. The abdomen was much contracted. He complained of severe pain in the head, over the eyebrow, and at the back of the head; there was excessive irritability of the stomach, fluids were instantly rejected, and the vomited matters contained much green fluid. Enema Ol. Ricini was administered; but, as it did not act, was followed by a colocynth one. Warmth was applied to the feet, a blister to the neck, and effervescing mixture with ammonia and carbonate of magnesia was prescribed. The vomiting continued; no urine was passed; he had no sleep, and became restless.

On the 26th he passed a light watery evacuation; the vomiting, however, continued, and he suffered from intense pain in the head. The hands were cold, the feet warm; the pulse 48, labouring; tongue furred; the abdomen still collapsed, and hiccough had come on.

On the 27th he was still restless; the symptoms continued as before; he was unwilling to be disturbed, and had no sleep from the severity of the pain. The pulse was 50, labouring; the respiration 16. The colocynth injection was repeated; but neither food nor medicine was taken.

On the 30th the pulse was not quite so labouring; the irritability of stomach continued, and soda-water, milk, &c., were at once rejected; his eyes were closed, and he was very unwilling to be disturbed; his hands were cold, his feet warm, his breath offensive, his pupils rather small.

On the 31st he seemed rather better; the pulse 60, rather more full. He got out of bed and passed a pint of urine, sp. gr. 1035, loaded with uric acid, but free from albumen and sugar; he appeared soporose. Nutrient injections were administered twice a day.

On June 1st he had no sleep, and the irritability of stomach continued; so also the pain in the head and the hiccough. Pulse 62. He appeared drowsy, but sensible.

On the 2nd there was no manifest improvement. A blister was applied behind the ear, and ʒj of mercurial ointment ordered to be rubbed in the axillæ night and morning.

On the 4th the pain and the vomiting suddenly ceased, and he began to crave for food. On the 5th the countenance was more intelligent; there was no return of pain. Inunction was continued for two or three days. On the 25th he was presented; he had steadily progressed, and there had been no return of the symptoms. For four months after leaving the hospital he continued well; but he returned to his habits of spirit-drinking; and on September 19th he was readmitted under my care. For two weeks he had had a renewal of the symptoms.

On September 22nd he had violent retching, with severe pain in his head; the pulse was slow, labouring, 80; the pupils were unequal, the right dilated; the hands and feet were cold. He was ordered to be cupped at the back of the neck to ʒvj, to take soda-water and milk, and a colocynth injection to be administered. The vomiting and pain ceased after the cupping.

The tongue, on the 25th, was clean ; the pulse 76. He was then allowed fish and vegetables. He continued in the hospital a few days, and left well.

XX. Disease of the lungs, or irritation of the pulmonary branches of the pneumogastric nerve, is the last cause of vomiting to which we refer. The vomiting in whooping cough appears to be of this kind, and equally so the vomiting which is often present at the early stage of phthisis ; the same symptom may occur in acute as well as in chronic disease of the lung. Sir Henry Marsh has mentioned early phthisis as one of the causes of the irritability of stomach to which he has given the name of regurgitative disease ; and too frequently it leads to the unfortunate expression, that the symptoms of early consumption are "all stomach." It seems that as the pulmonary disease advances, and disorganization takes place, this condition of irritability is lessened, although we often find that the paroxysms of cough are productive of violent vomiting.

Many interesting physiological questions in reference to vomiting might have been dwelt upon ; but sufficient has been stated to show that it is impossible satisfactorily to diagnose and to treat disease of the stomach without duly recognising the value of each symptom in its general bearing ; and further, that the most effective manner of alleviating *any* symptom, however distressing, is not by the treatment of that symptom, but by the removal of its cause.

The importance cannot be over-estimated of distinguishing between vomiting of cerebral, spinal, or nervine origin, and that which arises from gastric or other abdominal disease.

CHAPTER V.

DUODENUM.

THE symptoms which have been regarded by some writers as proceeding from disease of the duodenum have by others been referred to states of the liver, of the stomach, or of the pancreas.

My own observations, and the facts which I adduce in the following remarks, show that there are symptoms of disease justly attributable to this portion of the alimentary canal; and that in some cases we may, with care, satisfactorily diagnose that the duodenum is diseased. The peculiarities of its position and structure deserve our careful attention. Extending from the pyloric extremity of the stomach to the jejunum, it is about twelve inches in length, and may be divided into three nearly equal portions; the first is the most moveable, is almost surrounded by peritoneum, and is horizontal in its direction; it may be called the pyloric or stomachic portion of the duodenum, for it is associated with the stomach in its diseases. The second is vertical in direction, closely fixed near to the crura of the diaphragm, and to the vena cava; it receives the common bile and pancreatic ducts generally by a single opening, and is hepatic in its morbid relations. The pancreas is situated on the left side of the second portion; and the vena portæ, the hepatic artery, and the branches of the pancreatico-duodenal artery are also in relation with it. The third is horizontal in direction, and is simply intestinal in its function; the pancreas is situated above it; in front the superior mesenteric vessels enter the mesentery, and behind it is placed the aorta and the vena cava. The three portions of the duodenum are situated on *different planes*, the first portion being near to the anterior abdominal parietes, whilst the third part is immediately upon the spine;

and this arrangement allows the content of the canal mechanically to gravitate quickly into the jejunum, and assists also the discharge of bile from the ducts.

The muscular layers of the duodenum are double; a circular and a longitudinal coat, as in other portions of the small intestine. The mucous coat is covered with villi, which commence at the duodenum, and soon become exceedingly numerous; so also the *valvulae conniventes* are gradually developed, till we find them as large as in the jejunum. The whole of the surface is studded over with Lieberkühn's follicles; not unfrequently, especially in young subjects, there are solitary glands, as in the jejunum and ileum. There are also the glands of Brunner, minute compound glands peculiar to the duodenum, and which are situated beneath the substance of the mucous membrane; these commence a few lines from the pylorus, and extend about as far as the common bile duct; their function is not definitely known, but they are believed to resemble minute salivary or pancreatic glands. It sometimes happens that the solitary glands are so distinct, that they may very easily be mistaken for Brunner's glands; the latter are, however, situated beneath the mucous membrane, and microscopical examination at once manifests their difference.

There is still another point in connection with the duodenum that deserves consideration, and which indicates its close connexion with the stomach and with the liver. The pneumogastric nerves, branches of which supply the stomach, and also the liver, send filaments along the first portion of the duodenum, continued onwards from the lesser curvature of the stomach; this associates that part of the duodenum very intimately with the stomach.

The pancreatico-duodenal artery, which supplies the greater part of the duodenum, is from the hepatic, and the pyloric branch of the coronary extends into the first part of the duodenum, so that in the arterial supply we find the same association.

State of secretion.—The secretion is stated to be alkaline, and such is probably the case; the acid reaction after death arising from the gastric juice, which has gravitated through the pylorus. Whether a patulous, feeble contractile power in the pylorus, allowing the secretions of the stomach to pass at irregular

periods into the duodenum, is the cause of any of the discomforts associated with the forms of dyspepsia, we have no data on which to form an opinion. Corvisart states that the pancreatic fluid discharged into the duodenum has the power of dissolving albuminous substances; this opinion is, however, controverted by Dr. Brinton; the former describes duodenal dyspepsia as arising from an abnormal condition of this secretion.

Congenital malformation.—The duodenum sometimes has a double sigmoid curvature—a peculiar arrangement which I observed in a patient who died from intestinal obstruction. The ascending colon was adherent to the sigmoid flexure, and the cæcum twisted upon itself was situated in the left hypochondriac region. The person had been born at the seventh month, and the cæcum was preternaturally free.

In a cyclopean monster, I found the viscera of a double fœtus in a single peritoneal cavity; a double œsophagus was united in a single stomach, with a large convexity extending across the abdomen; and a single duodenum, placed vertically, received the biliary and pancreatic ducts on either side.

Diverticula are exceedingly rare as compared with those which arise from the lower part of the ileum; but small pouches are more frequently present, and they consist generally of mucous membrane, thus constituting a sort of hernial protrusion. In the museum of Guy's is one situated near the opening of the duct into the duodenum.

Some believe that the duodenum becomes distended with flatus, or with retained chyme, as the result of indigestion; and where there is mechanical obstruction, which we shall afterwards describe, this may be the case. It is possible also that an enormously distended transverse colon may impede the free passage of the contents of the third portion, but such is problematical. The distension which has been supposed to arise from the duodenum, will generally be found to be from the stomach or the transverse colon; for the duodenum passes quickly to a lower level, and I believe its contents at once gravitate into the jejunum.

As to the strictly pathological states, we find congestion sometimes active, more frequently passive; ulceration, cancer, and lastly mechanical obstruction.

To some it may appear altogether futile to speak of congestion or hyperæmia of the duodenum, but observation of the appearances after death convinces me that marked changes occur, and that in some instances a careful investigation might have pointed out their existence during life.

Great congestion of the duodenum is observed in various diseases in which a similar condition extends to the whole tract of the alimentary canal, as in disease of the mitral valve, and in portal obstruction in hepatic disease; but there are other cases in which we find active congestion, or at least the traces of its previous existence. The latter state of acute hyperæmia is illustrated in the following case :

CASE CVII.—*Inflammation of the Bronchi, of the Bile Ducts, or Biliary Hepatitis, &c. Acute Congestion of the Duodenum.*—Thomas H—, æt. 42, was admitted into Guy's Hospital, March, 1852; he had been ill for three weeks. He was a large, stout man, who for fourteen years had been in the police service; his habits of life had been very intemperate. Four years previously he had received a severe blow in his right side from a prize-fighter, and for some time he had been subject to vomiting in the morning, and the bowels had at times been much relaxed; before admission jaundice came on; he had had more anxiety of mind than usual, and gradually became languid and icteric. For four days his legs had swollen, afterwards his abdomen, and his strength became prostrated. The skin was of a dusky yellow colour; the tongue was dry, brown, and furred; respiration 44; the pulse 100, soft and compressible; the abdomen was much distended with flatus, and fluctuation could also be felt; the liver extended several inches below the ribs, and there was tenderness on pressure in that part. In the chest there were general bronchial râles; he was delirious at night, and slept but little; the motions were light in colour, the bowels relaxed, the urine contained lithates and the colouring matter of bile. Three days after admission he was more prostrate, and was delirious; the pulse was very compressible; he had pain in the right hypogastric region, and on the following day he died.

On inspection, severe capillary bronchitis was detected; the larger bronchi were also diseased; they were somewhat congested, and contained yellow-coloured tenacious mucus. The heart was large, and had around it a considerable quantity of fat; the right ventricle was thin; the left ventricle had undergone partial fatty degeneration. The valves were healthy, with the exception of slight thickening of the mitral. *Abdomen.*—There were several pints of yellow serum in the peritoneum; the intestines were considerably distended with flatus, and the liver extended several inches below the ribs. The duodenum contained *bloody mucus, the lining membrane was very much congested, and in some parts ecchymosed.* The lower part of the small intestine contained clayey fæces. There was a considerable quantity of fat in the omentum, and in the abdominal parietes.

The liver weighed 7 lbs.; its surface was smooth, and of a deep greenish-yellow colour, and some veins were seen upon it; the acini were whitish in colour. The section of the liver appeared coarse along the smaller branches of the vena portæ; the capillary vessels in Glisson's capsule were much distended, and some of them were quite turgid with blood. The smaller biliary vessels contained tenacious mucus, and their lining membrane was congested; this state of the bile ducts contrasted remarkably with the pale colour of the veins. The cells of the liver were gorged with fat, some of them were distended with oil globules; other hepatic cells appeared ruptured, and granules with oil globules were dispersed upon the field of the microscope. The deep green spots did not present any cells, but homogeneous matter with granules.

The larger bile ducts were free, but the opening into the duodenum was very much congested; the gall-bladder was empty; the kidneys were large and congested; the spleen was firm, and contained several fibrinous masses.

The health of this man was much impaired by his intemperate habits; his liver had probably been diseased for a considerable period. The affection of the chest came on subsequent to his admission into the hospital, and consequently after the jaundice. There was evidently acute disease of the smaller biliary tubes, as indicated by the congestion of Glisson's capsule, by the congestion of the lining membrane of the biliary tubes, and the tenacious mucus they contained; bile appeared to have been separated from the blood, but to have been retained in the hepatic structure. The bronchitis which subsequently took place was, perhaps, the cause of the fatal termination, and tended doubtless, to increase the congestion of the mucous membrane. The very congested state of the duodenum near the entrance of the bile-ducts indicated an extension of disease from the duodenum to the bile-ducts, or *vice versâ*; it was much more *localized* than is observed in the secondary congestion of the mucous membrane in pulmonary obstruction. This did not appear to be an affection in which much benefit could be obtained from the administration of mercury, but rather from salines with sedatives.

After burns the mucous membrane of the duodenum has been found greatly congested, and in several cases recorded by Mr. Curling in the Medico-Chirurgical Transactions this part of the intestine was ulcerated. This statement has not been confirmed by the observations of Dr. Wilks, recorded in the Guy's Reports for 1856. I witnessed many of the cases to which he

refers; and although in some the first part of the duodenum was hyperæmic, in none did I observe ulceration. A case of ulceration of the duodenum from a burn is, however, placed in the Museum at Guy's, in which Dr. Gull has drawn attention to a small ulcer existing at this part. The child survived twenty-five days, but died comatose; a small cicatrizing ulcer was found in the first part of the duodenum.

Mr. Curling describes diarrhœa, and the discharge of blood, as having arisen from this condition of the duodenum, and sometimes severe hæmatemesis and prostration. In some instances death took place from peritonitis consequent on perforation. After such severe injury to the skin, it is not surprising to find great disturbance of the circulation or of the internal organs, and especially of the mucous membranes, which are known to sympathise so closely with the skin; in some of these cases stimulants appear to have been administered freely, and these have probably conducted to this diseased appearance of the duodenum.

Chronic Congestion produces grey discoloration of the mucous membrane; and in the examination of the discoloured part we find that the deep colour is produced by the deposit of irregular grains of pigment, very thickly placed in the substance of the mucous membrane, near to its upper surface, and probably in the coats of the capillaries; the apparent explanation of this state being, that gastro-enteritis, or long-continued hyperæmia, has been followed by the deposition of hæmatine or pigment in the substance of the membrane.

In several cases of this grey discoloration the appearance, both in children and in adults, has been uniform. A child, aged nine, a thin, poorly nourished, pale boy, who had been subject for some time to looseness of the bowels, whilst running, hurt his thigh; he shortly afterwards complained of pain at that part; he was admitted into Guy's in a typhoid state, and died in two days afterwards. There was suppuration in the brain, and grey discoloration of the mucous membrane of nearly the whole of the small and large intestine.

Chronic congestion is observed, as before stated, in connexion with pulmonary and hepatic congestion, in fact, in any disease which leads to distension of the vena portæ; and we also find a

less general condition of congestion of the first part of the duodenum in disease of the pylorus, whether it be simple fibroid degeneration and hypertrophy, or true cancerous disease. The mucous membrane becomes thickened, its vessels congested, and its glands enlarged; sometimes, indeed, so much so that the glands might easily be mistaken for minute cancerous tubercles. The continued irritation thus leads to hypertrophy of the glands of the mucous membrane, as we find in other similar structures.

The duodenum is sometimes found, after death, to be filled with blood, and a coagulum is occasionally moulded into its exact form. This is due to extravasation of blood from ulceration and perforation of an artery, in the duodenum or in the stomach.

As to the symptoms arising from the conditions just described, they appear to be so continually bound together with those indicative of simple disease of the contiguous viscera, that definiteness and certainty cannot be attained. The vomiting and pain connected with hepatic disease and gall-stone are possibly due partly to the condition of the duodenum. In the latter there is probably spasmodic contraction of the canal; but of this we do not speak with certainty. In the cases described by Mr. Curling, vomiting was a frequent symptom; and the bilious evacuation in violent vomiting indicates that the first and second portions of the duodenum have been involved.

Instances are not unfrequently met with in which, several hours after food, there is pain at the region of the duodenum, perhaps with violent vomiting, faintness, pallor of the countenance; and these symptoms have by some persons been referred to the duodenum, as a form of *duodenal dyspepsia or inflammation*; by others to the pyloric valve; but occasionally jaundice follows, which appears to strengthen the former supposition. After intemperance, also, violent bilious vomiting, furred state of the tongue, loss of appetite and loathing of food, diarrhoea, tenderness of the right hypochondriac region, are followed by jaundice; and we are prone to regard the duodenum as being in, at least, a state of great hyperæmia. Exposure to cold, with great mental anxiety, tends also to promote this state of duodenal disease; and the mischief appears to be propagated to the like ducts, as in Cases CI and CVII. Sir H. Marsh has drawn

attention to the occurrence of jaundice with disease of the duodenum, in the Dublin Hospital reports; so also Dr. Stokes, in the 'Encyclopædia of Practical Medicine.'

Congestion of the duodenum is best relieved by diminishing portal and hepatic engorgement, and by stimulating the abdominal excretory organs to increased action. These objects may be attained by giving saline and mercurial purgatives, by aperient enemata, and by the application of leeches to the anus or to the scrobiculus cordis. A free dose of calomel, blue pill, or grey powder, followed by a saline aperient draught, often acts very effectively as a purgative; but in many instances, especially where the morbid condition arises from chronic pulmonary disease or obstructive disease of the heart, small doses of mercurials may be very advantageously combined with squills and foxglove, so as thoroughly to act on the abdominal excretory glands; but to give mercury so as to produce salivation, or to prescribe it in every instance where bilious fluid is rejected, appears to be an unwise course. The most bland nourishment should be given, and the abstinence from stimulants should be enjoined; ice and cold drinks often afford great relief when vomiting distresses the patient. In acute hyperæmic states, salines, as the solution of potash, the bicarbonates of potash or soda, the carbonates or the citrate of magnesia, may be given with diuretics in effervescence or otherwise, as the individual case may require. But in chronic hyperæmia, where there is profuse secretion of mucus, more advantage will be found from the dilute nitric or nitro-hydrochloric acids, with laxatives, as taraxacum, and the compound gentian mixture (P.L.), or with cinchona.

The most *acute* form of *inflammation* is sometimes observed after the administration of poisons. In a case of poisoning by sulphuric acid, where several square inches of the mucous membrane of the stomach had been destroyed, the duodenum was found intensely congested, and covered throughout by a thin, adherent, diphtheritic membrane. In this case the vomiting and dysphagia disappeared on the third day, and the patient, though extremely prostrate, did not appear to suffer much from pain. Arrowroot, lime-water, and milk, &c., were administered, and for a week it was thought that the patient might rally.

less general condition of congestion of the first part of the duodenum in disease of the pylorus, whether it be simple fibroid degeneration and hypertrophy, or true cancerous disease. The mucous membrane becomes thickened, its vessels congested, and its glands enlarged; sometimes, indeed, so much so that the glands might easily be mistaken for minute cancerous tubercles. The continued irritation thus leads to hypertrophy of the glands of the mucous membrane, as we find in other similar structures.

The duodenum is sometimes found, after death, to be filled with blood, and a coagulum is occasionally moulded into its exact form. This is due to extravasation of blood from ulceration and perforation of an artery, in the duodenum or in the stomach.

As to the symptoms arising from the conditions just described, they appear to be so continually bound together with those indicative of simple disease of the contiguous viscera, that definiteness and certainty cannot be attained. The vomiting and pain connected with hepatic disease and gall-stone are possibly due partly to the condition of the duodenum. In the latter there is probably spasmodic contraction of the canal; but of this we do not speak with certainty. In the cases described by Mr. Curling, vomiting was a frequent symptom; and the bilious evacuation in violent vomiting indicates that the first and second portions of the duodenum have been involved.

Instances are not unfrequently met with in which, several hours after food, there is pain at the region of the duodenum, perhaps with violent vomiting, faintness, pallor of the countenance; and these symptoms have by some persons been referred to the duodenum, as a form of *duodenal dyspepsia or inflammation*; by others to the pyloric valve; but occasionally jaundice follows, which appears to strengthen the former supposition. After intemperance, also, violent bilious vomiting, furred state of the tongue, loss of appetite and loathing of food, diarrhœa, tenderness of the right hypochondriac region, are followed by jaundice; and we are prone to regard the duodenum as being in, at least, a state of great hyperæmia. Exposure to cold, with great mental anxiety, tends also to promote this state of duodenal disease; and the mischief appears to be propagated to the like ducts, as in Cases CI and CVII. Sir H. Marsh has drawn

attention to the occurrence of jaundice with disease of the duodenum, in the Dublin Hospital reports; so also Dr. Stokes, in the 'Encyclopædia of Practical Medicine.'

Congestion of the duodenum is best relieved by diminishing portal and hepatic engorgement, and by stimulating the abdominal excretory organs to increased action. These objects may be attained by giving saline and mercurial purgatives, by aperient enemata, and by the application of leeches to the anus or to the scrobiculus cordis. A free dose of calomel, blue pill, or grey powder, followed by a saline aperient draught, often acts very effectively as a purgative; but in many instances, especially where the morbid condition arises from chronic pulmonary disease or obstructive disease of the heart, small doses of mercurials may be very advantageously combined with squills and foxglove, so as thoroughly to act on the abdominal excretory glands; but to give mercury so as to produce salivation, or to prescribe it in every instance where bilious fluid is rejected, appears to be an unwise course. The most bland nourishment should be given, and the abstinence from stimulants should be enjoined; ice and cold drinks often afford great relief when vomiting distresses the patient. In acute hyperæmic states, salines, as the solution of potash, the bicarbonates of potash or soda, the carbonates or the citrate of magnesia, may be given with diuretics in effervescence or otherwise, as the individual case may require. But in chronic hyperæmia, where there is profuse secretion of mucus, more advantage will be found from the dilute nitric or nitro-hydrochloric acids, with laxatives, as taraxacum, and the compound gentian mixture (P.L.), or with cinchona.

The most *acute* form of *inflammation* is sometimes observed after the administration of poisons. In a case of poisoning by sulphuric acid, where several square inches of the mucous membrane of the stomach had been destroyed, the duodenum was found intensely congested, and covered throughout by a thin, adherent, diphtheritic membrane. In this case the vomiting and dysphagia disappeared on the third day, and the patient, though extremely prostrate, did not appear to suffer much from pain. Arrowroot, lime-water, and milk, &c., were administered, and for a week it was thought that the patient might rally.

(See "Diseases of Stomach.") In ordinary practice, however, we do not meet with this form of disease.

Ulceration of the duodenum varies both in degree and extent; sometimes it is merely superficial, and is associated with other diseases, as in a patient who died from albuminuria with pericarditis, in whom the duodenum presented superficial ulceration, the result of erythematous or acute inflammation; or there may be chronic ulcer, resembling that found in the stomach, and presenting many symptoms in common with that disease.

Some duodenal ulcers have raised and thickened edges, with depressed centres, being evidently of slow formation. They are mostly found in the first portion of the duodenum; and since this part of the intestine is almost surrounded by the peritoneum, we sometimes have fatal peritonitis, produced by perforation, as in the stomach; the muscular and peritoneal coats being also destroyed by the ulcer; or adhesion takes place with the adjoining structures, as the liver and pancreas, &c.; and these oftentimes constitute the floor of the ulcer.

Several cases have come under my own notice the early symptoms of which were exceedingly slight, till sudden and fatal peritonitis had been set up by perforation. In some instances these ulcers have been associated with violent vomiting, and the persistence and aggravation of the vomiting were attributed to this diseased condition, as in a young woman, aged twenty-four, who was admitted into Guy's Hospital, with very urgent vomiting; the pulse was small and frequent; she was pregnant, and died in a short time from peritonitis; a small ulcer was found in the duodenum.* The vomiting was probably referred to sympathetic irritation from the uterine state; and a favorable prognosis would in many such cases have been given till the symptoms of peritonitis came on.

The second portion of the duodenum is, however, also liable to ulceration, as in a case preserved in the Museum of Guy's, where the coats of the whole of the vertical portion on the pancreatic side are destroyed, and the pancreas forms the base of a large chronic ulcer, in the centre of which is seen the opening of the biliary and pancreatic duct. There was a small ulcer in the third portion of the duodenum; and peritonitis had been

* Dr. Hodgkin on 'Pathology of Serous and Mucous Membranes.'

set up; the pancreas was enlarged. The patient, Samuel R—, was forty-four years of age, and had empyema; he became exceedingly emaciated before death, and suffered from vomiting as well as from melæna.

Ulceration is sometimes followed by constriction; and adhesions also frequently form between the first part of the duodenum and the gall-bladder; in some, ulceration extends from the gall-bladder into the duodenum, thus allowing the passage of calculi; and the gall-bladder is, in other cases, entirely obliterated.

Pain several hours after food, sallow complexion, furred tongue, feebleness of circulation, mental depression, nausea, irritable bowels, have been ascribed to ulceration of the duodenum, but the facts do not fully warrant this conclusion. In the several instances we have observed there were no such indications; in some, there had been disease of the gall bladder; in others, chronic disease of the liver; and the predisposing and exciting cause of the hepatic had probably induced the duodenal mischief.

Ulceration of the duodenum must be remembered both as a source of fatal perforation and of intestinal hæmorrhage.

The treatment of these cases is similar in all respects to that recommended for corresponding gastric disease.

CASE CVIII.—*Ulceration of the Duodenum. Perforation.*—George E—, æt. 30, a man of light complexion, and of steady and temperate habits, was admitted into Guy's Hospital, October, 1851. He was by trade a surgical instrument maker, and accustomed, when at work, to exercise pressure against the umbilicus. Four months before admission he had slight expectoration of blood, but it was doubtful whether it proceeded from the lungs or stomach. On October 20th, whilst apparently in good health, he suddenly experienced severe pain in the abdomen; to use his expression, he was "doubled up;" he fell down in a fainting state, and was taken into a druggist's shop, where ammonia and some castor oil were administered. The pain was situated on the right side. On admission, he was in a state of collapse; the pain of which he complained passed in the course of the ureter. On the following morning he was exceedingly depressed, skin hot, the abdomen tender, and there were the symptoms of general peritonitis; vomiting of coffee-ground fluid came on, and pulsation was felt at the scrobiculus cordis, which suggested the idea of aneurism. He survived fifty-six hours. On examination, the peritoneum was found to be intensely inflamed; lymph was effused, and castor oil was found floating in the peritoneal cavity. At

the first part of the duodenum, about one inch from the pylorus, an ulcer was found of the size of a shilling; and at its base there was a circular opening, the third of an inch in diameter. In the stomach several small aphthous ulcers were observed, and two small ones were covered with coagula. The remaining parts of the small intestine were healthy; so also the cæcum, colon, kidneys, spleen, and liver.

In the chest there were old pleuritic adhesions on both sides, especially on the left, where there was also a small vomica, with indurated lung, and thickened tubes.

The patient was only thirty years of age; and, as he believed, in good health, though evidently of feeble constitutional power, as indicated by the condition of the lungs and the previous hæmoptysis; he was doubtless phthisical, but the disease of the duodenum resembled, in its insidious character, the corresponding disease of the stomach, and gave no previous indication of its existence.

The treatment of the patient, before his admission, precluded all chance of recovery; but such, unfortunately, is too frequently the case. Brandy and castor oil, probably both, found their way into the peritoneal sac; and the necessary removal of the man, at first into a druggist's shop, then to his own home, and afterwards a considerable distance to the hospital, tended to induce increased extravasation and peritonitis; the judicious administration of opium prolonged life many hours.

As to the cause, the stooping posture at his work probably assisted to produce the disease; but this is involved in much obscurity.

The position of the pain did not point out the seat of the perforation; but this is only what has frequently been observed in cases of gastric ulcer; the pain was principally in the right iliac fossa, and it was believed that the ileum, or appendix cæci, had given way.

Mr. Travers, in the *Medico-Chirurgical Transactions*, mentions a case of perforation of the duodenum, about a finger's breadth from the pylorus, in a gentleman, aged thirty-five, who was strumous, but considered to be in good general health. There was a large irregular ulcer in the first part of the duodenum, with a small perforation, which had led to fatal peritonitis and death in thirteen hours; the perforation took place a short time

after a meal, the period at which such accidents are found generally to occur.

CASE CIX.—*Chronic Ulcer in the Duodenum. Carcinoma of the Liver. Jaundice. Granular Kidneys. Obliteration of the Bile Duct.*—George C—, æt. 46, admitted into Guy's Hospital December 14th, 1853, and died January 4th. For a fortnight he had had jaundice, vomiting, and typhoid symptoms, and for three months, from exposure to cold, œdema of the lower extremities had come on. In the liver there were from six to ten carcinomatous tubercles; the bile duct was obliterated near its opening into the duodenum, and throughout the liver the ducts were very much distended; the cells of the liver were normal. In the first portion of the duodenum there was a chronic ulcer, about an inch in diameter, with raised thickened edges, but not cancerous in its character; the rest of the intestine was healthy; the kidneys were large, and their surface irregular and granular.

The disease in the duodenum was not known till after death; the cancerous condition of the liver, inducing pressure on, and obliteration of the ducts, and the albuminuria appeared sufficient to explain all the symptoms. The ulcer in the duodenum, however, was in a chronic and passive condition, but nothing was ascertained as to its cause; we suppose that intemperance increased it. We rarely find such a complication of disease as cancer of the liver, acute disease of the kidney, and the condition of the duodenum just mentioned.

CASE CX.—*Strumous Disease of the Abdomen. Perforating Ulcer of the Duodenum and Cæcum.*—Jane B—, æt. 18, was admitted into Guy's Hospital February 19th, 1860, and died October 4th. At first the most prominent symptom was vomiting, which was supposed to be hysterical; but after a time the abdomen began to swell, diarrhœa came on, and emaciation, &c. increased, and these signs indicated the presence of organic disease. On inspection, the body was much emaciated; the legs were œdematous. The pleura was opaque, from the recent effusion of lymph, and the lungs were studded with lymph. The peritoneum was acutely inflamed; the intestines were reddened, and there was lymph upon them; there were tubercular masses upon the peritoneum, covering the liver. On withdrawing the cæcum, a small collection of offensive pus was found at its posterior part, and the abscess communicated with the cæcum by means of an opening about an inch above the ileo-colic valve. At the seat of perforation was a transverse ulcer, the edges of which were injected; the ulcer was one inch in length, and the opening one third of an inch. A few other ulcers were observed in the colon, but none were found at the termination of the ileum. The mesenteric glands were enormously infiltrated with cheesy deposit; so also were the lumbar glands. Behind the first portion of the duodenum, and close to the pancreas, was a

collection of offensive pus in front of the spine. This abscess communicated with the first portion of the duodenum by an opening about a quarter of an inch in diameter; the ulceration of the mucous membrane was more extensive than the external opening; and near to the perforation was a second smaller ulcer involving the mucous membrane. The first portion of the duodenum appeared to be contracted. The stomach was healthy; so also the kidneys. The spleen contained a softening strumous mass. The liver also was fatty.

Although the history of this case is imperfect, I have introduced it as an illustration not only of the obscurity of strumous disease in its earlier stage, but as an instance of irritation of the duodenum and colon, followed by ulceration and perforation, and producing peritonitis, at first of a local, but afterwards of a general character. The perforations in both situations were not directly into the serous membrane; the abscess connected with the duodenum was close to the pancreas upon the spine, and the colic one was placed behind the cæcum.

Cancerous disease of the duodenum.—It is far more frequent to find the duodenum 'secondarily involved than itself the primary seat of this fatal form of disease. In many cases the disease appears to have commenced in the pancreas or in the adjoining lymphatic glands, or in the liver; and although cancer of the stomach and of the pylorus is generally very defined and ceases abruptly at the commencement of the duodenum, such is not constantly the case, for the disease sometimes extends onward into the pyloric portion of the duodenum. Again, it is oftentimes very difficult precisely to state in which part the disease has commenced.

As to the symptoms, the earlier ones are often very insidious; and are more likely to be mistaken for hepatic disease than the early symptoms of cancer of the stomach; still the first indications are those of dyspepsia and malaise, sallowness of complexion, mental depression, followed by nausea, vomiting, and sometimes pain, several hours after food has been taken. The patient emaciates, and a hardness or tumour is felt about the cartilage of the tenth rib; a very difficult question then arises, as to whether it is the pylorus that is affected, or the pancreas, or the lymphatic glands. Pulsation

communicated to the growth may suggest the idea of aneurism. In aneurismal disease the vomiting is a less marked symptom, and the pulsation more uniform; the pain also is often very intense. In primary pancreatic disease the tumour is generally more central; the evacuations have been found sometimes to contain fat, and until pressure take place on the duodenum, or the disease extend to the stomach, and to the lymphatic glands, the symptoms are less pronounced. (See Case CXIV.) Pyloric disease is indicated by more persistent vomiting than we find in simple duodenal disease. Occasionally local ulceration, with chronic thickening, takes place at the union of the transverse and ascending colon, or cancerous disease may be developed at this site, and subsequently perforate the duodenum. (See "Cancer of the Colon.") The formation of adhesions with the duodenum in these latter instances sometimes causes partial mechanical obstruction; vomiting is produced, and thus the diagnosis is rendered unusually difficult; such was an instance—No. CXII—which we shall presently give. In all these maladies there is emaciation, pallor, cachexia. Lastly, we must refer to numerous diseases of the omentum and of the liver as complicating the diagnosis. Here, however the difficulty is less; for in the former the tumour is more central, there is greater mobility, and the gastric symptoms are less marked; in the latter, hepatic cancer, the tumour is more strictly in the hypochondrium, and the enlarged gland may be often felt with tubera projecting from its surface.

The termination of cancer of the duodenum is generally one of progressive emaciation and cachexia. If enlarged glands press upon the bile ducts, jaundice will be added to the symptoms; if perforation or sloughing takes place, local abscess occasionally forms, which, by giving resonance on percussion, adds increased difficulty in forming a correct diagnosis.

The treatment of these cases generally consists in trying to relieve the distress and pain of the patient, and in sustaining his exhausted powers. Anodynes are required—opium, morphia, chloroform or its preparations; bland, but very nutrient diet, and especially of a fluid kind, should be given. Stimulants assist in keeping alive the flickering flame of life. When great sallowness of the complexion comes on, or jaundice, it is very unwise

to give mercurials; they hasten degenerative changes, exhaust the patient, without any mitigation of his sufferings, and tend to hasten the fatal termination.

CASE CXI.—*Cancer of the Duodenum.* (Reported by Mr. C. Longmore.)—James R—, æt. 40, was admitted under my care into Guy's Hospital, June 23rd, 1858, and died July 5th. He was by trade a coach-builder, and he had resided at Newington; his habits of life had been temperate, and with the exception of a slight winter-cough, he had enjoyed good health till Christmas of the preceding year. The first symptom of which he complained was a shooting pain in the back and stomach; the pain at last became very violent, especially at night after he had finished his work; there were also moving pains in both sides, especially on the right, and in the testicles; he had neither cough nor vomiting; about four weeks prior to his admission swelling of the feet came on, and after a few days his abdomen began to swell; his urine was high coloured, scanty, and sometimes passed in drops. He was a man of sallow complexion, with dark hair and eyes; he was much emaciated, but the feet and legs were anasarcaous; there was dulness of percussion at the sides of the abdomen, and fluctuation was indistinctly felt. In the scrotum on the right side was a large hernial protrusion; and in the abdominal cavity a hard tumour could be felt, situated on the level of the umbilicus, and two inches to its left side; the tumour was an inch and a half to two inches in diameter, dull on percussion, but there was resonance around it; on pressure very slight pain was produced. Over the cartilage of the tenth rib there was also a minute pea-like tumour. The thoracic viscera were apparently healthy; the pulse feeble, compressible, 70. The surface of the body was cool. The tongue was coated with a brown fur in the centre, but was red at the tip. The bowels were freely acted upon, and the evacuations were paler than natural. The urine was scanty, sp. gr. 1032, free from albumen, but loaded with lithates. Small doses of acetate of morphia were given, and dilute nitric acid with infusion of cusparia. On June 25th, the abdomen had greatly increased in size, it was very tense and resonant on percussion except in the lumbar regions; there was hardness and tenderness on pressure below the ensiform cartilage; and he complained of aching pain and a sense of weakness in the loins. The urine was scanty, and it contained the colouring matter of bile. A bed sore had formed, and a spring bed afforded him considerable relief; but he was very restless. On the 26th, the report states that, during the previous evening and on this day, he vomited about two quarts of bitter bilious fluid, but became more comfortable after its rejection; although a sensation of intense thirst came on. About three ounces of high coloured urine were passed with difficulty. The pulse was feeble and compressible, 90. He complained of a dull pain at the lower part of the abdomen. Ice was given to him, and a small quantity of magnesia mixture with dilute hydrocyanic acid, and a castor oil injection was administered. On the 28th he had become jaundiced, and complained of great pain across the loins, of an aching, dragging character. The urine was scanty, cloudy, contained lithates, and under the microscope showed

blood-corpuscles with lithates. The castor oil injection did not produce any effect. Soda-water was allowed, and carbonate of ammonia with $\mathfrak{M}\nu$ of tincture of opium given every four hours, and a warm water injection administered. This injection acted freely, and he felt less pain. His principal complaint was of great weakness and depression of spirits. On the evening of the 3rd July vomiting of coffee-ground substance came on, and continued till his death on the 5th, at 11 p.m. The tumour several days previously seemed larger and more distinct. *Inspection* was made sixteen hours after death. There was rigor mortis; the whole body was jaundiced; the tissue of the heart was pale and softened. The liver was much enlarged, and it contained numerous cancer tubercles. The gall-bladder was much distended, and passed below the margin of the liver. A tumour about the size of the fist surrounded the vessels at the fissure of the liver; the duodenum was situated in front of this growth, and was adherent to it. The commencement of the duodenum was quite destroyed by cancerous ulceration, and a large slough occupied the position of the first portion. The interior of the intestine communicated with the cancerous mass beneath it; the cancer tumour was altered in structure, and contained blood. The gall-bladder was distended to about twice its natural size, and contained a few gall-stones. The hepatic duct was slightly obstructed. The vena cava was even acted upon, and in several places the cancer had penetrated through its walls. The whole liver was filled with cancerous tubera, and the greater part of its substance was thus occupied. These tubera were rounded, vascular, and softened. The disease appeared to run more especially in the course of the portal vessels, as if its entry into the liver had been by Glisson's capsule. The cancer growth consisted of large nucleated cells. The pancreas, supra-renal capsules, and kidneys, were healthy.

Instances of this kind are often very difficult of diagnosis, as to the precise seat of the disease; the glands close to the duodenum were probably first affected; but, although really behind the duodenum, the intestine did not cause resonance, probably on account of its becoming early complicated in the disease. The subsequent symptoms arose from pressure on the bile ducts and the vena portæ, and from the degeneration of the cancerous growth. Mr. John Dix, of Hull, has recorded a very interesting case somewhat allied to this; and in which there was a tumour apparently connected with the liver, but resonant on percussion. "The tumour was hepatic and malignant. It was softening down—sloughing, in fact; and in this process it had involved and laid open the duodenum, to which it was attached; and whence air had escaped into a circumscribed cavity formed by the tumour behind, and the abdominal wall in front, to both of which the transverse colon was adherent

below, forming the lower boundary" of the resonant space. The patient, "Mrs. M—, aged fifty-five, was pallid, feeble, and emaciated; she complained chiefly of pain in the right side of the abdomen, with vomiting and other symptoms referable to derangement of the hepatic and digestive functions. She had suffered, before that time, from jaundice and gall-stones." The patient died in about three months after the first medical examination; but the resonance in front of the tumour remained till death.

Instances also occur of primary disease of the pancreas extending to the duodenum, and we have witnessed such cases in which the mucous membrane of the duodenum had become infiltrated with medullary cancer. Organic cachexia is then generally well marked, but till the pylorus or duodenum become involved vomiting is not generally a prominent symptom.

Mechanical obstruction.—Other parts of the intestine are much more liable to obstruction of a mechanical character than the duodenum. In the course of several years we have observed, or have found recorded, isolated cases of this kind of obstruction, arising from the following causes:—

1. Peritoneal adhesions.
2. Gall-stones of large size, which having ulcerated through coats of the gall-bladder, have become impacted in the duodenum, and have led to fatal obstruction.
3. Enlarged glands, infiltrated by cancer, compressing the second or third part of the duodenum.
4. Diseased pancreas.
5. Hydatid disease of the liver, opening into the duodenum.
6. Foreign bodies.

It is exceedingly common to find, after death, that adhesions have taken place between the *first* portion of the duodenum and adjoining viscera, either the inferior surface of the liver and gall-bladder, or the transverse colon; and, in many instances, the impediment to the free passage of the chyme is so slight that no symptoms point to any disturbed function. In the following case adhesions with the colon were followed, however, by great distension of the first part of the duodenum; but there was also some ulceration of the same part of the intestine; there was

chronic ulcer of the colon, and chronic as well as acute peritonitis, with strumous and glandular disease, so that there was considerable difficulty in unravelling the symptoms, which resembled those of organic disease of the stomach. Still we believe that the pain and the vomiting several hours after food had been taken, were the result of this duodenal obstruction.

CASE CXII.—Chronic Peritonitis; Acute Peritonitis; Tubercular Deposit on the serous Membranes and in the Glands; Constriction of the Duodenum, and great Dilatation of its first portion; Small Ulcer in the Duodenum; Large Chronic Ulcer in the Colon.—William C—, æt. 38, was admitted into Guy's Hospital under my care, April 15th, 1861. He was a married man, by trade a cooper, and he had resided at Dockhead. About seven years previously he suffered from severe pain at the epigastric region; and for several years since that time he had had pain at the same part, but less acute in its character. He had never had any hæmorrhage from the stomach, but he had complained of slight pain in the dorsal region, between the sixth and eighth vertebræ. Some years before he had had violent vomiting; but since that time vomiting had been slight, the attacks coming on some time after food had been taken. He had had slight pyrosis, and acid taste after vomiting. The pain at the epigastric region was not constant, but it was worse after food, and was especially aggravated by constipation.

On admission, he was very much emaciated, with a sallow complexion, and on the forehead there was a bronzed condition of his skin; the skin at the elbows was also slightly discoloured. There was moderate tenderness at the scrobiculus cordis; the abdomen was rounded and supple; no tumour could be felt; the bowels were rather confined, the pulse was very compressible, the tongue was red in patches. No disease could be detected in the lungs or heart. The patient stated that the bronzed colour of the forehead had existed for three years, and had been produced by exposure to the sun; the lower part of the abdomen was also found to be slightly discoloured. He was ordered to take fluid diet, and of extract of *nux vomica* gr. ss, with extract of aloe and of henbane, of each gr. ij, three times a day.

On April 20th. The bowels were freely moved, and he had severe pain at the scrobiculus cordis; the pain was neither relieved nor modified by any change of position. The tongue was slightly furred, with red patches upon it. Brandy had previously been given, and was increased to four oz.; and chloric ether, ℥xv, and nitrate of bismuth, gr. x, in mucilage mixture (G.P.), were administered three times a day. He was unable to take meat.

On the 30th he continued in the same prostrate condition; the pulse was very compressible; the heart was regular, but the first sound could scarcely be heard. He had no pain nor vomiting. No tumour could be felt in the abdomen. The medicine was continued, and nourishment in the form of jelly, &c., given.

May 28th. Since last report he had continued the same treatment, and was rather less prostrate; the discoloration also on his face was less marked.

On June 11th, soon after breakfast, violent pain came on in the abdomen, which was partially relieved by a hot poultice. The abdomen for some weeks had been contracted, but now became rounded, hot, and tender. At 2 p.m. he was very prostrate, his countenance more than usually distressed, and his eyes very sunken. The pulse was very compressible; he had not passed urine for nearly twenty-four hours; and, since he had pain in the region of the bladder, the urine was ordered to be drawn off, and he was to take opium, gr. j, every four hours.

12th. He was considerably relieved by the opium, and by the urine being drawn off; no pain existed in the abdomen, but it was distended moderately, although the tenderness was less; the tongue was moist; the pulse rather firmer.

13th. At 9 a.m. vomiting of bright green fluid came on, with great prostration; pain in the abdomen and back was increased; his hands became cold, and his pulse gradually imperceptible. Consciousness remained till nearly the close of life, and he died about five o'clock.

14th. At 2:30 p.m. *inspection*. The body was very much emaciated. The head was not examined. *Chest*.—On the left side the pleura was firmly adherent, and on tearing it away, rounded yellowish tubercles, two to three lines in diameter, were found thickly covering the costal surface. The left lung itself was very small; but there were no tubercles in it. The right pleura was free from adhesions or tubercles, and the lung was also quite healthy. The heart and pericardium were normal. There were several yellowish-white tubercular masses in the glands in the anterior mediastinum. On opening the abdomen, the intestines were seen to be distended; and the enlarged transverse colon, extending from one hypochondriac region to the other, prevented the stomach from being seen. There were numerous peritoneal adhesions, especially at the upper part of the abdomen; the transverse colon, stomach, and duodenum being united firmly to the under surface of the liver. The coil of the small intestines presented considerable injection at their lines of contact; but neither was lymph effused, nor had the serous membrane lost its shining colour. Numerous tubercles were present on the serous membrane; some were exceedingly small, others three or four lines in diameter, and they were situated on the intestine or on the peritoneal surface of the liver. The mesenteric glands were extensively diseased; and all the glands situated in the neighbourhood of the pancreas, and near the origin of the thoracic duct, were enlarged, although it could not be demonstrated that the duct was compressed. The glands contained much cheesy and cretaceous matter, and some more recent semi-transparent deposit. On removing the transverse colon, the stomach was found partially distended, and an elongated sac was produced, partially contracted, about three inches from the right extremity; this sac was at first supposed to be from hour-glass contraction of the stomach, but, on opening it, the first contraction was seen to be pylorus, and the second enlargement was an enormously distended first part of the duodenum. The stomach and duodenum contained grayish-green fluid and mucus. The mucous membrane of the stomach did not present any abrasion, thickening, nor ulceration, nor was the pylorus hypertrophied; there was a little arborescent

injection. The sac formed by the first part of the duodenum was capable of holding eight to ten ounces of fluid ; there was some arborescent injection. Immediately beyond the pylorus was a small ulcer about five lines by three in size, its edges rounded and without any injection ; it did not extend into the muscular coat. Three inches from the pylorus the intestine was narrowed, and there was a constriction resembling a second pylorus ; there was no thickening nor cicatrix, and it appeared probable that the peritoneal adhesions had looped up the intestine. On the gastric side of this constriction there was a small pouch, capable of admitting the tip of the finger. The rest of the duodenum, the jejunum, and the ileum, were healthy, with the exception of one or two small ulcers with tubercular deposit on their peritoneal surface. Peyer's glands were healthy. The cæcum and appendix also were normal. In the ascending colon the solitary glands were very distinct, and at the commencement of the transverse colon were the remains of an old ulcer ; for two to three inches the mucous membrane was irregularly destroyed and puckered, and of a gray colour. The rest of the intestine was normal. The supra-renal capsules, the kidneys, and the liver, were healthy ; two or three strumous tubercles were, however, situated on the peritoneal surface of the liver.

In the *second cause* of mechanical duodenal obstruction, impaction of a gall-stone, the symptoms resemble those produced by internal strangulation of the intestine, or by hernia, but vomiting is set up at a very early period, and is of a severe character. The vomited matters, however, cannot have a stercoraceous odour nor appearance. The diagnosis is generally obscure and difficult ; but where the symptoms of the passage of a gall-stone, namely, intense pain in the hypochondrium, vomiting, and subsequent jaundice, are followed also by the symptoms of insuperable obstruction, the nature of the malady is sufficiently clear ; but in the ulceration of a large gall-stone through the coats of the gall-bladder into the duodenum, the indications of disease may be so slight as to be almost overlooked, and the subsequent obstruction cannot then be distinguished from strangulation taking place high up in the intestine. The impaction of the gall-stone is generally found to happen near the termination of the duodenum, or in the upper part of the jejunum.

In obstruction from diseased lymphatic glands in the neighbourhood of the duodenum, the occlusion sometimes becomes suddenly complete, and the symptoms are those of internal strangulation ; but more frequently the pressure is less, and the

symptoms are those which we shall presently have to refer to in connection with disease of the pancreas; thus in an instance of femoral hernia after the intestine had been returned, the symptoms continued, and the patient quickly died. The third portion of the duodenum was then found to have become firmly impacted between two enlarged glands.

CASE CXIII.—*Obstruction from Biliary Calculus in the upper part of the Jejunum, thirty inches from the Pylorus.*—The calculus is in the museum of Guy's. The case was under the care of Ebenezer Pye Smith, Esq., and is recorded in the Pathological Transactions of 1854.—The patient was a stout woman, æt. 69; she had good health till three months before death, when she suffered slight pain in the right hypochondrium, which continued a fortnight, unaccompanied by sickness or prostration. She recovered, but continued her usual sedentary habits; five days before her death she began to feel sick, and vomited bile in large quantities; the urine was moderately secreted. The vomiting increased in violence, but with only very slight pain in the abdomen; on the fifth day she became comatose. A calculus composed of inspissated bile, and measuring four and a half inches in circumference by two and a half in its lesser circumference, was found impacted about thirty inches from the pylorus. There was much fibrous tissue on the under surface of the liver; and an ulcerated opening extended from the gall-bladder into the duodenum, below the bile ducts.

There are in these cases three symptoms which especially deserve attention, as guiding us to right diagnosis, when viewed in connection with the previous history. The absence of abdominal distension, the early period at which vomiting takes place, with the character of the ejected matters, and the diminution in the quantity of urine which is voided.

The absence of distension of the abdomen is an important sign of occluded intestine in the early part of its course. In obstruction of the large intestine, or even at the lower part of the small, the abdomen becomes enormously distended, and the peristaltic movements can often be observed in spare persons through the parietes; this is especially the case in disease of the sigmoid flexure of the colon. The stoutness of the patient in Case CXIII, rendered this sign less observable; again, where this duodenal obstruction exists with hernia, the diagnosis must necessarily be most obscure. As to the vomiting, it comes on very early, and the matters rejected are bilious. In strangulation of the ileum, and obstruction of the colon, unless irritating

purgatives are given, this distressing symptom may be considerably postponed; and when it does take place and is continued, the matters are of a stercoraceous character. Still, in acute peritonitis, as from perforation, the sudden bilious vomiting may greatly mislead us. Again, very violent bilious vomiting sometimes takes place in disease of the stomach, and in cerebral disease; but the signs of obstruction are then wanting.

Gall-stone produces, with vomiting and constipation, intense pain in the region of the gall-bladder; this severe character of pain we do not find in intestinal obstruction, but on the other hand it must be acknowledged, that when slow ulcerative absorption has taken place between the walls of the gall-bladder and the duodenum, calculus so extruded is followed by less severe suffering than in ordinary cases of biliary calculus.

A very interesting case, under the care of Dr. Lever, is mentioned by Dr. Barlow, *Guy's Reports*, 1844:—The patient, aged fifty-one, a year before her death had the symptoms of gall-stone, and her bowels afterwards became constipated; a short time before her death, excessive pain, vomiting, and constipation came on, with scanty urine and collapsed abdomen. The gall-bladder and duodenum were firmly adherent; the two upper thirds of the duodenum were contracted, thickened, and would only admit a common quill; about the centre of the ileum was a biliary calculus of the size of a walnut, partially sacculated.

With regard to the quantity of urine excreted being a sign of the seat of obstruction, as mentioned in the paper by Dr. Barlow, just referred to, he argues that the quantity of urine must necessarily be small, from the diminished fluid brought within the range of the absorbing surface of the portal veins; and thus there must be diminished supply to the heart and kidneys; but there is often a large quantity of fluid ejected by vomiting, which would diminish the renal secretion. If the obstruction be incomplete, or low down in the intestine, the kidneys pour out a larger quantity, and the vomiting is also less severe.

Dr. Barlow has, in the paper previously cited, dwelt upon the importance of bearing in mind, that in ischuria renalis, violent vomiting, constipation, and scanty urine are sometimes present.

In diseased pancreas the obstruction is less complete, but acts by inducing firm adhesions about the first and second

portions of the duodenum; and pressure is also exerted by the increased size and hardness of the pancreas, and by infiltrated glands. The symptoms resemble those of obstructed pylorus, vomiting several hours after food, gradually increasing emaciation, constipation; and these symptoms are slowly developed during several months. A tumour can generally be felt near the region of the pylorus.

The following very interesting case was regarded as one of cancerous disease of the glands in the neighbourhood of the pancreas, and secondary implication of the stomach; for the vomiting took place three or four hours after a meal, as in obstructive disease of the pylorus; and the general symptoms resembled those of organic gastric change.

CASE CXIV.—Disease of the Pancreas; Suppuration and Gangrene; Pressure on the Duodenum.—James P—, æt. 60, by occupation a publican, and resident at Camberwell, was admitted under my care on July 4th, 1861. He stated that he had always enjoyed good health till four months prior to admission; when he was suddenly seized with severe pain in the region of the stomach, and with vomiting. The vomiting returned at intervals of three or four days, and came on several hours after food. Four years previously he had begun to feel slight pain at the region of the stomach, which came on every three or four months, but was relieved by taking a little cayenne pepper with brandy. He had not received any blow, nor had he suffered from any hæmatemesis. The pain was situated at the epigastric and umbilical regions, and extended to the spine; it was of an acute kind, and had not the gnawing character of pain often described by patients affected with ulcer of the stomach.

On admission, he was very much emaciated, with an anxious countenance, sallow complexion, and sunken eyes; his skin was hot and dry, and he complained greatly of thirst; the tongue was furred, the pulse frequent and sharp, the respiration normal; he had slight cough, but it did not distress him; and there was no evidence of thoracic disease by percussion nor by auscultation. The abdomen was contracted moderately, except at the lower part of the epigastric and at the umbilical region, where there was a rounded tumour, evident on visual examination. The tumour was dull, and tender on percussion; no fluctuation could be felt, and it had slight pulsation anteriorly from contact with the aorta, but no general aneurismal thrill. There was resonance between the tumour and the liver, as well as between the tumour and the spleen; in fact, both the hypochondriac regions were more than usually resonant. Pressure on the tumour produced a feeling of nausea; the bowels were constipated; and the appetite was very poor. His weakness compelled him to remain quietly in bed. The urine was high coloured and scanty, and was free from albumen. Fluid food was ordered, and soda-water with brandy, and chloric ether $\mathfrak{m}\mathfrak{x}$, with nitrate of bismuth gr. \mathfrak{x} in mucilage mixture.

July 5th. He was slightly relieved by the medicine, but the vomiting continued; the ejected matters consisted of deep-green fluid, containing a large quantity of mucus, of squamous epithelium, and some nucleated cells (from gastric glands). These attacks of vomiting distressed him greatly; every kind of food was rejected at once, but the medicine and ice relieved his distress partially; his prostration, however, increased, hiccough distressed him; and he had an offensive taste in the mouth.

July 8th. He was very restless and prostrate, and the vomited matters were of very deep-green colour. At 9 p.m. he was suddenly taken worse, and continued in great pain during the night. At 7 a.m. next morning he expired.

Inspection seven hours after death.—The body was very much emaciated. The thoracic viscera were healthy, excepting old pleuritic adhesions. The peritoneum contained some dirty gray fluid, and had in some parts lost its shining smoothness; the intestines were slightly distended. The sac of the lesser omentum was distended by a large abscess, which had constituted the tumour felt during life. On tracing the duodenum upwards, at its centre was found an œdematous portion bulging out; and containing fluid resembling that in the peritoneum; but there was no perforation. By dividing the peritoneum between the stomach and the colon, an abscess was opened; it had dense fibrous walls, about two lines in thickness, in some parts irregularly sinuous, and having several bands on its walls, the remains of occluded vessels. Above and partly in front of the abscess was the stomach; below was the colon, and at its superior, right, and inferior parts was the duodenum greatly distended, and its coil enlarged. The abscess contained dirty offensive pus, and at its posterior part was a black slough about two and a half inches in length; some concrete yellow matter was also found on its walls. The abscess rested on the spine, the crura of the diaphragm, and on the superior mesenteric and splenic veins as they formed the vena portæ. It extended on the left to the spleen. The pancreas for two to three inches towards the splenic extremity was healthy, but the rest of the gland was in a sloughy state, and constituted the black mass found at the floor of the abscess. The pancreatic duct existed in the centre, and degenerating gland tissue was observed under the microscope. The gland and duct were separated from their duodenal attachment. The common bile-duct was healthy, and its opening into the duodenum was free; but the gall-bladder contained numerous gall-stones about the size of peas. The liver and spleen were healthy. The stomach was very much enlarged and distended; it contained tenacious green mucus, such as was vomited during life; its mucous membrane presented numerous points of arborescent injection, so also that of the duodenum; but no direct communication with the abscess could be found, nor any ulceration of the surface.

The origin of the disease in this remarkable case could not be ascertained, viz., whether a pancreatic calculus had set up the abscess, or whether inflammation had been produced in the cellular tissue about the gland. No direct blow had been

received, and the disease slowly advanced. Acute peritonitis, from the transudation of offensive purulent serum into the general cavity of the peritoneum, was the cause of the fatal termination.

Dr. Bright believed that the fatty motions which he found in some of these cases were indicative of disease of the pancreas, but this has not been confirmed by subsequent observations.

The course taken by *hydatid disease* of the liver is uncertain; sometimes it is towards the surface, and a rounded tumour is then felt on the anterior abdominal parietes; or it extends through the diaphragm into the lungs. In a case under the care of Dr. Rees, in Guy's, the cyst opened into the duodenum. Hydatids were both vomited and passed by stool, and the former symptom was very severe. The patient was exceedingly ill, and a friction sound was audible over the seat of the tumour, evidently from local peritonitis; the patient steadily improved after the evacuation of the hydatids by vomiting; the tumour disappeared, and he left the hospital; but after a few weeks intense peritonitis came on, and he quickly died. The remains of hydatids were found in the liver; and the duodenum, colon, liver and kidney, were firmly united by adhesions. A large abscess existed between these structures, and had led to the fatal peritonitis. No communication existed between the liver and the colon; and although the duodenum at its second part was firmly adherent, no direct opening could be found.

The patient was twenty-nine years of age, and had resided at Twickenham; he was temperate in his habits; for nine years he had suffered from so-called "bilious attacks," and from vomiting, with slight sallowness of the skin; five years previously he had had severe jaundice, which continued for three weeks. Eight months before admission his appetite became ravenous, but he lost strength and became emaciated; for seven weeks he had been confined to his bed from severe pain about the umbilical region; jaundice came on, but disappeared, and was followed by very severe pain in the right hypochondriac region, extending to the loins, and a rounded growth presented itself below the ribs on the right side.

A remarkable instance of mechanical obstruction in the duodenum, from a foreign body, is recorded by Dr. Blakeley Brown, in

the Pathological Transactions of 1851 and 1852:—A delicate young woman, aged eighteen, became gradually emaciated, and at last died from peritonitis. The stomach, duodenum, and upper part of the jejunum contained casts composed of agglutinated and interwoven masses of string and hair.

Gastric Solution of Duodenum.—The mucus of the duodenum is frequently found in an acid condition after death, which is probably due to some of the gastric juice slowly gravitating through the pylorus; but in some instances the pylorus is so patulous, that gastric juice readily passes, and exerts its solvent power after death in the same manner as in the stomach. Such a state was found recently in a child who died under my care in Guy's.

CASE CXV.—Perforation of Duodenum after Death by Solution of Gastric Juice.—William B—, æt. 4 years, was admitted July, 1856, and died on the 23rd. He was an anæmic child, with large head; on admission he was in a semi-comatose state, and the pupils were widely dilated, and he had occasional vomiting, but no convulsions; six weeks previously he had had measles, and one week afterwards hydrocephalus gradually became developed; he was in an almost hopeless condition on admission.

Inspection was made fourteen hours after death. The arachnoid was covered with a slight layer of lymph, so as to give it a greasy appearance, and at the base of the brain there was considerable sub-arachnoid effusion. The ventricles contained two ounces of fluid, of sp. gr. 1001. There were miliary tubercles in the lungs and in the bronchial glands.

In the stomach there was considerable gastric solution, the mucous membrane being destroyed; but in the duodenum the intestine was quite divided, all the coats destroyed, and the end of the first portion terminated in an irregular ragged margin. The contents of the stomach were found in the peritoneal cavity. There were tubercles in the mesenteric glands, and an isolated one in the kidney.

CHAPTER VI.

MUCO-ENTERITIS AND ENTERITIS.

THERE has been considerable confusion in the application of the term enteritis; Broussais considered it to be inflammation of the colon, Abercrombie regarded it as inflammation of the peritoneal and muscular coats of the intestine; others, again, more particularly apply the term to an inflammatory disease of the small intestine, which commences in the mucous membrane, and extends in severe cases, so as to involve all the coats of the intestine, even its peritoneal investment. These latter and more severe instances correspond to the enteritis phlegmonodea of Cullen; the former, when the mucous membrane only is affected, to his enteritis erythematica.

Watson, Barlow, &c., apply the term only to the more severe cases of inflammation of all the coats, but we shall in this chapter also consider those in which little more than the mucous membrane is affected, called muco-enteritis, and closely allied to gastro-enterite and gastric remittent fever. Dr. Copland describes glandular enteritis, and ileo-colitis; the former we consider in the remarks on strumous and typhoid disease of the intestine, the latter with dysentery.

Enteritis, then, manifests itself under two forms: 1. That involving only the mucous membrane, and which has a disposition to extend in the course of the mucous membrane—*muco-enteritis*; and, 2. That in which the disease extends in depth, rather than on the surface, and implicates the muscular, peritoneal coats and the connecting tissues: both commence in the mucous membrane.

In hernia, whether external or internal, acute enteritis is set up; and there may be symptoms in common with enteritis, as

constipation, vomiting, &c., but their pathology and treatment are so diverse, that a separate consideration of them is required.

It would be difficult to draw a defined separation between cases of inflammatory diarrhœa, as described by Dr. West, and the simplest forms of enteritis; they pass the one into the other. Diarrhœa, however, is not an ordinary symptom of enteritis; for the bowels in the latter disease are frequently constipated.

We shall first consider enteritis in the form of *muco-enteritis* or *enteritis erythematica*. This is very frequent among children during dentition or weaning, and after the exanthemata; but in many cases of infantile diarrhœa and colic, a more transient condition is set up; the indications of which are twisting pain in the bowels, and the evacuation of watery or green motions, with general fretfulness, &c.; these symptoms pass away in a very short time, and they arise from irritation of the bowels rather than from inflammation of the mucous membrane.

In muco-enteritis a child is found to be fretful, without its usual playfulness and mirth. The lips are dry, and the little patient has a circumscribed flush on one or other cheek; the skin is dry or roughened; the abdomen is somewhat enlarged, or considerably distended and tympanitic, and varies in degrees of tenderness; but the restlessness of the child causes it to cry when no pain is produced; it is unwilling to be disturbed; the appetite is irregular and capricious, either craving for cooling drinks, as cold water, or for unsuitable food, which is oftentimes the cause of the malady. The bowels are irregular, either constipated for several days, or attacked with diarrhœa, the motions being offensive and pale, or greenish in colour, or the evacuations consist of slimy mucus, or of food scarcely changed, and these conditions may alternate, the one with the other; the tongue has a whitish fur, and its substance or papillæ are often much injected; vomiting may easily be induced, and probably arises from the extension of the mischief to the stomach, when the disease is called gastro-enteritis. In the evening the child becomes still more restless, the skin is hot, and even pungent; the sleep is disturbed, and accompanied with grinding of the teeth or starting, and it often awakes alarmed; in the morning the

febrile disturbance is less, and the child may be cheerful and playful.

This aggregation of symptoms constitutes gastric remittent or infantile remittent fever, and many look upon it in the same light as typhoid fever, considering that the inflammatory condition of the intestine is a concomitant, not the essential part, of the disease. This is, I think, incorrect; the intestinal disturbance is the source and the cause of the continuance and extension of the disease; and not, as in typhoid fever, the manifestation of a previously existing and general condition.

When the symptoms persist severely for several weeks greater prostration ensues, the child wastes sometimes to an extreme degree, it appears haggard and aged, the lips have dry sordes upon them, the tongue is more injected, and often aphthous. There is less remission in the morning; the child will scarcely sleep at all, or, in very young children, be placed out of the arms of its nurse; the diarrhoea increases, watery evacuations or food unchanged are discharged a short time after having been taken; the pulse becomes very rapid, the eyes half closed, and the child dies from exhaustion, almost before the nurse is aware of any change; or the brain becomes oppressed, and a drowsy, torpid condition, or convulsions sometimes precede death. The convulsions and coma, to which we refer as coming on at the close of this intestinal condition, are closely allied to those produced by exhaustion, as in the hydropcephaloid disease of Dr. Marshall Hall.

Muco-enteritis is frequently followed by tympanitis, and by strumous disease of the peritoneum, or of the mesenteric glands. In such cases, although the more prominent symptoms of vomiting and purging subside, the child remains wasted, the abdomen enlarges, the appetite becomes ravenous, and exhaustion steadily progresses to a fatal termination. (See Strumous Disease.)

In young persons we sometimes find a state of muco-enteritis similar to that described, but without phthisis or strumous disease; the eyes are sunken and bright, the lips parched, the tongue exceedingly injected, and beef-like; the cheek is occasionally flushed by a circumscribed patch on one or other side;

the pulse is compressible, but frequent; the skin is at one time dry, at another perspiring; there is thirst, generally with loss of appetite, and sometimes with great irritability of the stomach; the bowels are constipated, or diarrhœa alternates with constipation. The urine is scanty and high coloured. This condition may persist for many weeks, with gradually increasing exhaustion, and in some cases it terminates fatally; in very many instances it yields to judicious treatment, but there is great danger of relapse. In young women this state is sometimes associated with painful or deficient menstruation, or with leucorrhœa; and may be accompanied with severe neuralgic pain in the abdominal parietes, and below the mammæ. This neuralgia occasionally leads to a more unfavorable prognosis than the case warrants.

CASE CXVI.—*Gastro-Enteritis. Recovery.*—Frederick C—, æt. 18, was admitted into Guy's Hospital under my care, October 8th, 1858; he had been employed in the Patent Office, and had resided in Chancery Lane. A month previously he felt hot and unwell, and a week before admission there was great febrile disturbance, with diarrhœa and prostration of strength; the evacuations were watery. His face was flushed on admission; the mind clear, but there was some excitement, although no delirium; the tongue was red, injected, beefy, and cracked; the bowels were still loose, and the pulse was compressible; the skin was dry, but no maculæ were present; no evidence of disease of the heart nor lungs existed; the abdomen was supple, but there was slight pain below the umbilicus. There was no fever in the house from which he came, nor offensive drains; and the appearance of the patient was *not* that of one affected with typhoid fever. Ordered milk diet and two eggs. Infusion of calumba with aromatic spirit of ammonia three times a day, and Dover's powder, gr. v., every night.

On October 13th the bowels continued loose; the motions bilious, but fluid; the abdomen was slightly tender; he was prostrate, and appeared to sink in the bed; the skin was dry, but the tongue was rather less injected; there was no delirium at night, but his countenance had a somewhat excited appearance; the cheeks were flushed, the body spare. Ordered to take compound krameria mixture \mathfrak{zj} every six hours, and the pills to be repeated.

On the 18th the bowels were quiet, the tongue less injected, and the symptoms less severe. To take quinine mixture \mathfrak{zj} twice a day, and fish diet.

23rd. No pain. Appetite returning. Tongue less injected.

Nov. 12th. He was presented convalescent.

The *second* form of enteritis is more severe, and all the coats of the intestine are involved. The symptoms are exceedingly acute, and too frequently advance to a fatal termination with

great rapidity ; or they may be extended over many weeks or months. Severe pain is generally present, and it has more or less of a paroxysmal character, and is accompanied with great tenderness and distension of the abdomen ; there is frequently vomiting of bilious fluid, and the bowels are often constipated ; the pulse is small, wiry, and sometimes compressible, the tongue is partially furred, the patient lies on the back with the legs drawn up, as in acute peritonitis, and prostration may rapidly ensue ; or, whilst the more severe symptoms subside, the tongue becomes dry, red, and glazed, the bowels loose, and the strength gradually undermined ; or, again, the convalescence may be as rapid as the occurrence of the symptoms. In these conditions the mind is generally perfectly clear.

Several instances which have come under my own observation will illustrate the disease.

CASE CXVII.—*Acute Enteritis*.—A child about seven years of age, after eating freely of raw apples, was seized with pain in the abdomen around the umbilicus ; the bowels were constipated ; the abdomen was tender and distended ; the countenance was expressive of much distress ; the pulse was rapid ; the tongue had a slight fur upon it. The constipation continued ; the abdomen became more tender and distended, and the child was found lying on its back in severe pain, with the legs drawn up, and with occasional vomiting. This state continued for several days ; the bowels then were freely acted upon ; but the child became prostrate, and shortly died, four or five days from the commencement of the disease. On opening the abdomen, the intestines were found much distended with flatus ; the peritoneal surface was intensely injected where the coils were in contact, and was covered with lymph. The mucous membrane of the small intestine was congested, and portions of undigested apples were found in the intestine.

The inflammation had been set up by crude undigested food ; it extended rapidly from the mucus to the muscular and connecting tissues, and to the peritoneum. The inflamed intestine was unable to propel its contents, and hence the constipation ; and sometimes the constipation is so marked, that it is the most prominent symptom. The severe pain in this form of enteritis contrasts with the absence of it where the mucous membrane only is affected.

The symptoms in other instances closely resemble those consequent upon mechanical obstruction.

CASE CXVIII.—*Enteritis simulating Mechanical Obstruction*.—Henry V—, æt. 17, was admitted into Guy's Hospital July 24th, 1850. He was a tall, thin lad, who had been employed in a tobacconist's shop; and a week before admission he had had diarrhoea, which had been checked by an opium pill. The day before admission he felt well, and whilst walking out of doors he ate some apples and cherries; a few hours afterwards severe pain in the abdomen came on. Some rhubarb, with compound chalk powder and opium, was prescribed; the bowels were opened twice during the night; but at seven in the morning severe pain in the abdomen returned; his countenance was then expressive of great distress; the eyes were sunken, and the bowels were confined; the tongue was furred and clammy; he was rolling himself from one side of the bed to the other, from the intensity of the pain; the recti muscles were rigid, but pressure could be borne; an emetic was administered, and some undigested apples and cherries were vomited. Calomel gr. v, with opium gr. iss, were given, but at once returned; a turpentine injection was then administered. Vomiting then came on, at first of bilious, afterwards of stercoraceous fluid; the injection brought away some scybalous matter, but without relief to the pain. The calomel and opium were repeated, but they were at once returned. At 5 p.m. he was placed in a warm bath; the tongue was clean, the pulse 130, the abdominal muscles were rigid, and the paroxysms of pain in the abdomen were very severe. Calomel and opium, of each 1 gr., were given every three hours; and to relieve the intense pain, chloroform in *mx* doses was prescribed. His pulse became exceedingly rapid, and he died at eleven the next morning, about thirty-six hours from the commencement of the attack.

On inspection, the intestines were found to be very much distended; the peritoneum was injected, and delicate portions of lymph passed between the coils; on turning aside the small intestines, the cæcum, colon, and about three feet of ileum were found collapsed, pale, and empty; at this point there was a sudden cessation of the intense congestion and distension, giving the appearance of constriction; but no constriction nor twist could be detected; the mesentery, however, attached to this part, and connected with the last lumbar vertebra, contained several hard and calcareous glands, and appeared slightly contracted; on raising the intestine, and placing it in a straight line, air at once passed, and the constriction disappeared. The intestine was full of pale yellow fluid fæces, and contained some undigested matter; no ulceration existed, and the other viscera were healthy.

In this case severe colic came on after taking indigestible food; inflammation of the mucous membrane of the small intestine was produced; this extended to the muscular and peritoneal coats, and was followed by intense pain, by distension, and by vomiting; it appeared that the slight interference with the movement of the ileum opposite to the calcareous mesenteric glands led to the limitation of the disease at that part, and that over-distension following inflammation was the principal cause

of the obstruction. The abdomen for several hours was tolerant of pressure, and the symptoms of peritonitis came on later; could the enteritis have been subdued, the obstruction would probably have disappeared. This case was under the care of one of my colleagues, at Guy's, and is recorded as a good illustration of severe enteritis.

The following case is one in which the most acute enteritis produced scarcely any symptom; the patient was semi-comatose; but it is closely allied to cases in which local enteritis is apparently set up by obstructed vessel:

CASE CXIX.—*Sloughing Ileum. Peritonitis. Large and Fatty Kidneys. Degeneration of the left Lobe of the Liver. Lobular Pneumonia. Small and Fatty Heart.*—Thomas C—, æt. 43, was admitted into Guy's Hospital, December 7th, 1853, and died December 31st. By trade he was a sail-maker, and during the last two years of his life had been very intemperate. He was admitted with anasarca, and coagulable urine; diarrhoea and wasting came on, and before death he passed into a semi-comatose condition. *Inspection* was made forty-seven hours after death. The body was spare and pallid; the right side of the body was œdematous (from position); there were pleuritic adhesions at the apex of the right pleura; the lungs were very œdematous, and some lobules were softened and breaking down. There was slight atheromatous deposit on the mitral and aortic valves. *Abdomen.*—The intestines were distended; there was general peritonitis, but only slight injection of the peritoneum at the edges which were in contact; eight inches from the ilio-cæcal valve, the peritoneal surface of the intestine for several inches was of a dark grey colour, as if on the point of sloughing; but there was no constriction nor strangulation, nor had there been any symptom of it during life. The mucous membrane at the lower part of the ileum was in a sloughing condition, but this diseased portion was defined, and intensely congested at the margin; the slough was thin; but it affected the whole of the mucous membrane, and was not confined to Peyer's glands; the mesenteric veins were filled with clot. The left lobe of the liver was wasted, forming a fibrous mass, and was white in colour; the remaining part of the gland was fatty. The kidneys were large and white.

In this patient the disease of the kidney had led to uræmic poisoning, and to the semi-comatose condition; hence the non-complaint of pain in the severe peritonitis which ensued. There is great disposition in uræmia to serous inflammation of the arachnoid, pleura, pericardium, and peritoneum; but it is rare to find such a state of acute inflammation as that described in this case.

Pathological appearances.—In muco-enteritis we may find

very much less change than had been anticipated. Neither ulceration nor congestion may be observed throughout the whole canal. It is probable that the injected condition has in these instances, like erythema of the skin, entirely passed away; thus also some of the most severe forms of bronchitis present scarcely any morbid appearance of the bronchial tubes themselves, the congestion having disappeared, although the altered mucus remains.

A change in the character of the secretion from the mucous membrane is a sure indication of its deviation from the normal condition; but, unfortunately, we do not possess the same facility for the examination of the secretions from the digestive as from the respiratory mucous membrane. Adhesion of a thin stratum of faecal matter is an indication of an imperfect secretion of mucus; or a lymph-like exudation takes place; this exudation resembles the diphtheritic membrane occasionally found in the throat, and it consists of an immense number of granules with nuclei; it may sometimes be easily scraped off, exposing an injected surface beneath; in other instances a section of the whole membrane shows that it is firmly united.

The mucus which is found in the intestine presents indications of rapid change having taken place, nuclei and elongated cells of incomplete epithelium being found in great abundance; and similar mucus is found in the evacuations during life.

Crystals of triple phosphate are frequently detected on the surface of the mucous membrane. It is probable that in many instances this is a post-mortem change; but in other cases we find such crystals when the inspection has been made a few hours after death; and they probably result from decomposition of the mucus, as in the urinary bladder after chronic inflammatory action.

The solitary glands may be very large and prominent; a state which is due to the age of the patient, and the functional activity of these structures, or to the excitement of morbid action. In other instances a granular and thickened state of the intestines is observed, as if its mucous membrane were sprinkled over with fine sand. Small aphthous ulcers sometimes exist, even perforating the intestine, as in a case recorded among the inspections at Guy's, in which there were minute

ulcers extending throughout the whole of the small and large intestine ; and perforations of the cæcum and transverse colon had led to fatal peritonitis.

Grey discoloration is often observed around the solitary follicles, or it is more general in character, either in the large intestine, in the lower parts of the ileum, or even in the duodenum. This state consists in the deposit of pigment in the membrane, and is the result of continued congestion ; it apparently follows also as an effect of muco-enteritis.

The most intense form of local enteritis exists when a portion of the intestine has become strangulated ; the mucous membrane is then swollen, and it is then intensely injected ; portions of faecal mucus adhere to the valvulae conniventes, or the whole surface of the mucous membrane is covered by a thin adherent layer of granular lymph ; all the coats of the intestine become thickened, and the areolar tissue is œdematous ; the peritoneum is covered by lymph ; it is intensely congested, and of a purple or slate colour, or even gangrenous.

A condition closely resembling that just described is sometimes found without any strangulation, either affecting only the ileum, as in Cases CXVIII and CXIX, or of a dysenteric character, and involving also the cæcum and colon ; thus, a distended and congested state of the ileum may terminate suddenly, as if constricted, and the portion of small or large intestine below be pale and contracted ; on removing the intestine, the apparent constriction ceases, the canal becomes perfectly free, and the congestion is the only thing that marks the obstruction. There has been much discussion whether in these cases there is really obstruction by a twist of the intestine, by a spasmodic condition of the contracted part, or by a paralysed state of the distended one ; the last supposition is now generally regarded as the correct explanation, namely, that the inflamed intestine becomes distended, its peristaltic contraction enfeebled, so that at last it is unable to contract upon, and propel its contents. The abrupt termination may be determined by a cicatrix, by slight peritoneal adhesion, or by old disease of the mesenteric glands, as Case CXVIII. These instances closely resemble true ileus from strangulation or from other mechanical cause.

Ulceration and sloughing, or gangrene, generally follow this

form of enteritis ; but, although in hernia and internal strangulation the gangrenous part is at the seat of constriction, this is not always the case in obstruction from other causes ; in obstructive disease of the sigmoid flexure, ulceration takes place above the seat of the obstruction ; but the most acute inflammation and ulceration will often be found in the ileum, cæcum, and ascending colon. The inflamed mucous membrane in these parts gives way from the enormous distension ; numerous ulcers, arranged in *transverse* lines, are closely set together in the ileum and cæcum, and some of these occasionally extend through the peritoneum.

Obstruction of the mesenteric vessels usually takes place in cases of internal hernia ; but it is probable that obstruction of these vessels is sometimes the cause of intestinal disease, rather than the effect. Intense engorgement of a few inches of intestine may be seen, and the mucous membrane may be almost in a sloughing state, without any symptom having been manifested during life, and without any obstruction being found after death ; the mesenteric veins or arteries extending to the part are found obstructed, and have perhaps led to changes in the nutrition of the intestinal coats.

In inflammation of the mucous membranes there is a great tendency to the extension of the disease by continuity of structure. Sometimes the alimentary canal, in its whole tract, appears to be inflamed ; at other times it commences in one part, and extends from that as from a centre. Inflammation of the colon will pass into the ileum ; that of the ileum into the large intestine, as well as into the jejunum, duodenum, and stomach. In enteritis the small intestine only may be affected, or the cæcum and colon be also implicated. Broussais* speaks of this extension of disease ; but, though we are not disposed to agree with his opinions, we must, I think, acknowledge the truth of the frequent extension of disease to contiguous and continuous structures. It is probably as true of the mucous membrane as it is of the skin, as exemplified in erysipelas.

The correct *diagnosis* of enteritis is very important, because means may be employed which tend to aggravate the disease,

* Broussais, 'Phlegmasies Chroniques.'

valuable time may be lost, and such aid as might have been of essential service neglected.

Hernia, external or internal, intussusception, and mechanical obstruction from any cause, may be confounded with enteritis arising from simple inflammation.

It is well always to examine the ordinary positions of external hernia; many mistakes would have been avoided by this simple means. In internal strangulation the pain generally comes on after sudden muscular movements or after exertion of the strength; the patients often affirm that until the time of sudden exertion they enjoyed comfortable health, then something seemed to give way, or there was a "catch," and fixed pain was felt, from which the subsequent pain radiated; the seat of pain, however, does not necessarily indicate the seat of obstruction, as found after death; because distension and the movement of viscera produce much alteration in the position of the intestine. After the sudden onset of pain, constipation and vomiting with varied degrees of severity come on, till prostration, collapse, and death ensue; and the rapidity of the symptoms may be as great as in external hernia. We do not observe this fixity of pain in enteritis, although it may be at first localized to a comparatively small space.

In internal obstruction without strangulation, we often find previous constipation, and the commencement of the attack is slower, the pain being sometimes very slight till towards the close of the malady.

In intussusception the sudden severe pain is very different from that of enteritis, and is more likely to be confounded with simple colic. When the symptoms of obstruction from intussusception become developed, the discharge of bloody mucus is often observed; and the value of this diagnostic indication has been shown by Mr. Gorham.* In enteritis it is very important carefully to ascertain the symptoms which marked the onset of the disease.

It is difficult to distinguish some cases of *chronic poisoning*, or even of *acute poisoning*, from enteritis arising from other causes. In these instances, inflammation of the mucous mem-

* Guy's Reports, 1838, p. 300.

brane is produced. I may refer to cases of chronic poisoning by arsenic; the vomiting is often very severe, and the irritability of the stomach a very prominent symptom, but never stercoraceous; the abdomen is generally less tender than in the worst cases of enteritis; but in doubtful cases we must be guided by the concomitant symptoms and the analysis of the vomited matters. In the enteritis from crude indigestible food, and irritants, as some forms of mushrooms, the symptoms may be very similar to those consequent on ordinary poisons, so that we may be unable to distinguish the one from the other.

In *simple colic* there is less difficulty; there is absence of tenderness, and the pain may be actually relieved by pressure.

In *peritonitis*, suddenly induced by perforated intestine, the collapse is greater; the abdomen becomes exquisitely tender and tympanitic; vomiting is not generally produced, unless the peritoneal surface of the stomach and its coats become involved. But from whatever cause enteritis is induced, peritonitis is a very common result; the muscular coat is implicated, and the peristaltic action by a wise provision checked.

In *hysteria*, we sometimes find tympanitis with constipation, with irritable stomach, and with pain in the abdomen; and these symptoms might, by carelessness, be mistaken for acute inflammation. The expression of countenance is not that of severe abdominal disease; the vomiting may be induced by anything put into the stomach, but disappears at other times. The pain is cutaneous, and the abdomen is tolerant of continued pressure, unless there be inflammatory disease of the ovaries. There is generally leucorrhœa, with painful or disordered menstruation; but the patient often remains in a tolerably nourished condition.

Ischuria renalis.—Dr. Barlow has pointed out the importance of bearing in mind disease of the kidney in its sympathetic symptoms. In suppression of the urine, vomiting and constipation are often produced; but the cerebral oppression is generally very marked, and the examination of the urine (drawn off by catheter, if none can be passed) would at once decide the character of the complaint, if there be any obscurity.

Cerebral disease.—It is not unfrequent, as we have before noticed, to find vomiting present as a symptom of disease of the

brain, and then also associated with constipation ; but there are some peculiarities in this state which distinguish it from enteritis and mechanical obstruction. There is no pain or distension about the abdomen ; the tongue, the countenance, and the other symptoms of disease are different. In young children it is sometimes difficult to distinguish muco-enteritis from true hydrocephalus ; there is irritability of the stomach in both, with perhaps diarrhœa, heat of skin, startings in the sleep, loss of appetite, unwillingness to be disturbed, &c. ; but in the former, the abdomen is more distended, in the latter it is collapsed ; the tongue is injected, furred in the one case, but clean in the other. In hydrocephalus also there is greater pain in the head, or drowsiness ; there is disturbance of the pupils, which are contracted, or in the later stages widely dilated, with strabismus ; and the fontanelles become distended ; the vomiting in hydrocephalus is often induced by only raising the body from the recumbent posture. In the exhaustion which occasionally follows severe diarrhœa, or muco-enteritis in infants, a series of symptoms, resembling hydrocephalus, or, as they have been called, hydrencephaloid disease, supervenes ; these, however, are very different from true hydrocephalus ; they should be borne in mind, lest the effect of exhausting disease be misinterpreted ; in these cases we have the half-closed eye, the emaciated expression, diarrhœa, collapsed fontanelle ; and the early symptoms are seen to commence in abdominal, not in cerebral disease.

Causes.—The ordinary causes of enteritis are improper or indigestible food ; this is especially the case in infants and children in whom the disease is set up during dentition or weaning, or after exanthems, especially measles. Exposure to cold or wet, sleeping in damp beds, or in the open air ; violent and sudden contortions of the body, excessive muscular exercise, as in walking, are other causes ; so also, mechanical obstruction, however produced, whether by hernia, intussusception, internal strangulation, tumours, &c.

Prognosis.—The unfavorable symptoms of enteritis are the long persistence of the disease, emaciation, the development of peritonitis, distension of the abdomen, hiccough, prostration of strength, irregular pulse, haggard and anxious expression, sunken eye ; or, after constipation of obstinate character, the onset of

severe diarrhoea, consisting of thin offensive or serous mucus; also, partial sweats, inability to take food, persistent beef-like tongue.

In muco-enteritis, the continuance of diarrhoea, thin serous evacuations like the washing of beef, great exhaustion of the patient, exceedingly rapid pulse, convulsions, are the precursors of a fatal termination.

Enteritis is less amenable to treatment where there is strumous diathesis; the mesenteric glands are prone to become congested, swollen, and infiltrated; and the patient gradually becomes exhausted, or strumous disease is developed in other parts; but there is scarcely any condition of simple enteritis and muco-enteritis from which patients, especially infants, may not recover.

A more favorable prognosis may be given when the pain in the abdomen subsides, when the bowels act naturally, and the evacuations are of a healthy character; when the tongue is uninjected, the skin supple and generally perspiring, the pulse quiet, the countenance cheerful, and when there has been refreshing sleep.

Treatment.—If we consider the pathological conditions of the disease, the indications of treatment become evident. We believe, then, in the existence, in these cases, of an inflamed condition of the mucous membrane, which may, or does already extend, to the submucous, muscular, and peritoneal coats.

It is exceedingly unadvisable to try and produce action on the bowels by violent purgative medicine, as by jalap, senna, scammony, calomel, blue pill, croton oil, crude mercury, and the like. The peristaltic action is checked by the inflamed state of the coats of the intestine, and additional irritation retards it still further. Leeches applied freely to the abdomen, according to the age of the patient, or depletion from the arm, has, in many instances, been followed by free evacuation from the bowels, and by the relief of pain. Warm fomentation should be applied to the abdomen.

When irritating ingesta are retained, producing and perpetuating the disease, we may administer, at an early period, a purge of calomel or grey powder, followed by castor oil, or linseed oil with opium, or a free saline purge, as the potassio-tartrate of soda, or sulphate of magnesia.

When, however, there is tenderness, it is more safe to give calomel, or grey powder, combined with opium, several times during the day; but it is well to avoid the continued use of mercurials.

Alkalies are of service, in acting as sedatives to the mucous membrane, in diminishing its engorged state, and in neutralising irritating secreta; as the bicarbonate of potash, in doses of gr. x, or gr. xv, and the solution of potash, in doses of ℥xv to ʒss. Chlorate of potash, in gr. v to gr. x, and carbonate of soda, gr. v to gr. xv, combined with narcotic remedies, as hyoscyamus, conium, are in other instances apparently beneficial. The latter appear to act on the involuntary muscular coat, and on the nerve supply of the intestine. A valuable combination in less severe cases is grey powder with Dover's powder.

Some administer magnesian salines, as sulphate of magnesia and calcined magnesia; but where there is tendency to extension of the disease to the peritoneal coat, I think sulphate of magnesia is injurious, in increasing the peristaltic action of the intestines; although in its direct effect on the inflamed membrane, it may lead to the emptying of the capillaries by watery evacuation.

Rest in bed is important, that the intestines may not be disturbed in their position; since perforation, in many cases, follows ulceration of the intestine; and there may be also extension of peritonitis from inattention to this simple rule. There must also be abstinence from irritating food; in fact, nothing but the most mild and bland ingesta should be taken; demulcent drinks, milk alone, or united with lime-water or soda-water, as the case may be, will be grateful to the patient. Great care is necessary after the subsidence of the more active symptoms, in the return to nourishing and substantial food, as well as in the use of any active exertion.

In children with muco-enteritis, chlorate of potash is a valuable remedy; in some cases, it appears to act with as much benefit as in cases of stomatitis. Citrate or bicarbonate of potash are also of real service. In other cases, when the motions are clayey and white, minute doses of calomel are sometimes administered, with carbonate of soda, or chalk, as the compound soda powder of the Guy's Pharmacopœia; but much injury is

often done by calomel and grey powder in these cases, and in numerous instances we have found their use unnecessary; astringents may be given, as chalk, with catechu, or krameria, or logwood, with small doses of opium; but in very young infants it is better altogether to avoid the use of opium, if possible.

Maunsell and Evanson mention the value of dilute nitric acid with minute dose of opium and simaruba, and I have often used this combination with advantage. Ipecacuanha is a valuable remedy where there is no irritability of the stomach, and may be combined with chalk medicine.

In children, also, it is essential only to administer food that can be easily digested, and although it may appear of a proper kind, if the symptoms continue, a change should be made. The disease often comes on at weaning; and the greatest care is required in seeking for a suitable diet; "tops and bottoms," with water, and with or without a small quantity of milk; dried flour, biscuit powder, &c., may be given, or milk-and-water alone. I have seen cases where the only food that could be borne was water boiled for a considerable time with rice, and after partaking of this fluid the vomiting and purging ceased, and a gradual return to more substantial food was attained. For some infants, it may be necessary to obtain a wet nurse, but this is a measure to be avoided if possible. Asses' milk is the best substitute for the natural supply; and a small quantity of cream, with water, can sometimes be taken when simple milk cannot be borne.

The prostration in children is sometimes so great that stimulants are necessary. The aromatic spirit of ammonia may be given with infusion of cusparia, and with astringents; but in many instances I have seen life apparently saved by the timely use of brandy, or wine, administered very frequently and in small doses; thus to a child aged three, almost in a dying state, the pulse scarcely perceptible, the extremities cold, the eyes half closed, brandy in doses of 15 drops, diluted with water or demulcents, was given every quarter of an hour, and in a few days the child was really convalescent. In other cases, white wine whey produces a similar beneficial result.

CHAPTER VII.

STRUMOUS DISEASE OF THE ALIMENTARY CANAL.

INFLAMMATORY disease of the alimentary canal in strumous subjects can scarcely be separated from the more slow and insidious strumous disease, associated with less active symptoms. Struma should not be looked upon as a disease of isolated organs of the body ; but one in which the power of assimilation is diminished, the nutritive functions imperfectly performed, and an unhealthy and unorganizable plasma readily poured out. Disease set up by the ordinary exciting causes in subjects of this kind leads to the various forms of strumous deposit and its subsequent changes. A blow on an epiphysis leading to strumous disease of the bone ; a slight bronchitis to strumous pneumonia, and the deposit of tubercular substance in the lungs ; over-excitement of the brain to hydrocephalus and strumous meningitis ; slight irritation of the mucous membrane of the intestine, or muco-enteritis, to deposit of unorganizable product in the mesenteric glands or in the mucous membrane. The abnormal condition antecedent to these deposits is, I believe, correctly designated tuberculosis ; damp air, a want of light and proper food, imperfect rest, hereditary disposition, and, perhaps, syphilitic taint, induce the imperfect elaboration of those products necessary for healthy growth and nutrition ; and in this state the blood, the nervous force, the vital activity of every part of the body, is unable to return to the normal type on the slightest derangement, but strumous inflammation and strumous deposit take place ; nay, more than this, tuberculosis is oftentimes so marked, that apparently without any fresh exciting cause, the forces required for ordinary healthy nutrition are insufficient for their proper function ; and an unhealthy plasma is poured out, constituting miliary tubercle.

1

Strumous disease of the alimentary canal is observed under various forms:—

1. As it occurs in children, indicated by severe diarrhœa with or without strumous disease of the mesenteric glands.
2. Primary disease of the mesenteric glands.
3. Tubercle in the peritoneum, and strumous peritonitis in its several forms.
4. Tubercle in the mucous membrane with enteritis, leading to softening, ulceration, and perforation, as is frequently observed in phthisis.
5. Tubercle in the appendix cæci.

1. *Diarrhœa in strumous children.*—The symptoms are very similar to those which we have described as gastro-enterite, but here, being engrafted upon a strumous constitution, they are more easily induced, and are less yielding to medicinal treatment. This disease causes the death of thousands of infants among the poor of London, but it does not spare the rich, when there is hereditary disposition; the diarrhœa is frequently set up by some change in the diet; by the nourishment having been of an improper character; by disordered secretions poured out from the mucous membrane of the stomach, intestines, or liver; and it often follows the exhaustion of measles, scarlet fever, &c. Many of these instances are cured by the removal of the exciting causes, and by the administration of simple, corrective medicines. When, however, these causes cannot be removed: when the infant cannot be taken from offensive exhalations, and from a damp or cold atmosphere, and have proper food administered to it, especially when there is a very feeble and strumous constitution, too frequently does the diarrhœa continue; the little patient becomes wasted, the countenance is expressive of extreme distress, and has an aged, care-worn appearance; the evacuations consist of greenish thin mucus, of food only partially changed, or they resemble the washings of meat, and are exceedingly offensive. The skin is dry, sallow, and wrinkled; the abdomen is full, sometimes hot and tender, and there is pain of paroxysmal character; the mouth is dry, and sometimes aphthous, the tongue is slightly furred, the

breath is offensive, the eyes languid, the sleep is often disturbed with starting moans. The child is exceedingly restless, fretful, and almost incessantly whining; sometimes the stomach is irritable, or the appetite is craving, and the child distressed by thirst. Such are the symptoms of severe gastro-enterite rendered intractable by strumous deposit, and passing into the conditions described as *tabes mesenterica*.

In some cases even of extreme exhaustion, the little patient rallies when proper remedial means can be employed; in others the diarrhoea persists day after day, slightly abating and then returning with renewed violence, till at last the infant dies exhausted, or convulsions come on before the close of life. It rarely happens that with very severe diarrhoea there is much cough, although the lungs may be throughout filled with miliary tubercles.

Post-mortem appearances.—After death we may find no apparent change in the whole tract of the mucous membrane; the liver, spleen, and lungs may be normal; the mesenteric glands may be enlarged and swollen, and in some instances contain evidence of low organized deposit. It might be questioned, whether disordered mucous membrane did not induce this condition of the glands; but whether so produced or primary in its origin, there can be little doubt that it leads to the maintenance of an abnormal state of the mucous canal, and indicates strumous cachexia. When we have a more vigorous constitution, one free from struma or imperfect nutritive power, the patient often rallies, and the fatal symptoms are checked.

Treatment.—In the treatment of these cases, it is most important to remove all exciting cause of disease, and every impediment to the healthy performance of nutrition and growth, to inculcate perfect cleanliness, and the inhalation of pure air, to administer the most mild and unirritating food, and to afford warmth to the body.

The child should be placed in a warm bath, clothed in flannel, and the air of the room maintained at an equable temperature. Milk will not agree with some infants, in whatever form it may be given; in others, asses' milk, or milk with lime-water or soda-water, is retained, or water boiled for a long time with rice; in others, "tops and bottoms," made with water and

without sugar may be tried; or similar substances, such as dried flour, biscuit-powder should be given, and, as a *dernier ressort*, a wet-nurse employed.

In the medicinal treatment, where chalk mixture made with dill or cinnamon water, and with or without a few drops of ipecacuanha, does not avail, I have found great benefit from the administration of the compound logwood mixture of the Guy's Pharmacopœia:

Misturæ Cretæ fluidunc. vj.
 Extracti Hæmatoxyli dr. j.
 Vini Ipecacuanhæ fluidr. j.
 Vini Opii fluidr. ss.

in doses of one or two teaspoonfuls, *or*, the compound infusion of catechu, with a small quantity of opium, and, if need be, a few drops of aromatic spirit of ammonia. The krameria is also a valuable astringent, with or without chalk and opium, as in the following Guy's preparation:—Decoction of krameria ʒxv, (root ʒix, with water Oj, boiled to ʒxv.) Ipecacuanha wine and tincture of catechu, of each ʒvj, and syrup ʒiss. A teaspoonful to a table-spoonful as a dose, according to the age of the child.

Where a strumous condition exists, great benefit is derived from the administration of cod-liver oil, with steel wine, or from the latter medicine alone. If vomiting be absent, cod-liver oil is sometimes exceeding serviceable; when it cannot be taken, dilute nitric acid, with infusion of cusparia, and a few minims of compound tincture of camphor, are of benefit, especially when other means have somewhat moderated the diarrhœa.

I have frequently observed, at the City of London Dispensary, and still more extensively at Guy's Hospital, many instances of this form of strumous disease. Diarrhœa is rendered uncontrollable by strumous cachexia, or by disease of the mesenteric glands; and some of the worse forms of the disease are found to occur, as the sequelæ of exanthems, coming on especially after measles.

In some cases small doses of sulphate of copper, as $\frac{1}{4}$ to $\frac{1}{2}$ of a grain, or of nitrate of silver in similar quantity, or of acetate of lead in $\frac{1}{4}$ or 1 grain doses may be prescribed with one or two

grains of Dover's powder. Mercurials are, I have generally found, detrimental, and continued doses of calomel greatly aggravate the disease. Small enemata of starch may be used with benefit, and where we have a good nurse, other agents may be well applied in this way; a weak solution of nitrate of silver or of borax tends to diminish the irritation of the lower bowel, and may prevent prolapse. When exhaustion is extreme, nourishment must be administered every few minutes, if the stomach can retain it; and small quantities of wine or brandy conjoined, as previously mentioned. In not a few cases the latter has been the means of prolonging life and restoring infants to health who were apparently in the arms of death.

This form of diarrhœa, consequent on strumous disease, is, however, not confined to children. The following case is an instance of that kind, where apparently simple diarrhœa assumed an obstinate type; no form of medicine or diet checked it for many days, and at last the patient sank. There was evidence of some inflammatory action at the lower part of the ileum; intense congestion, slight diphtheritic effusion, and ulceration of the Peyer's glands were found; but these appearances were so local that they were not considered sufficient in themselves to explain the severity of the abdominal symptoms. There were minute tubercles in the peritoneum, and low organized deposit in the mesenteric glands; and these were indications of the strumous character of the patient. In the lungs there was neither vomica nor miliary tubercle; some iron grey deposit at the apex, and slight strumous deposit were observed, and at the lower lobe there was ordinary hepatization, which had evidently come on a short time before death.

CASE CXX.—Slight Strumous Disease of the Mesenteric Glands. Diarrhœa. Pneumonia.—Charles A—, æt. 30, a waiter, was admitted into Guy's Hospital, August 15th, 1855.

Three years previously he had had severe diarrhœa; and five weeks before admission, he had had pain at the stomach, with loss of appetite, and cramp, &c. He lost flesh considerably. His stomach was sometimes irritable, and he had suffered from vomiting. On admission there was febrile excitement; the diarrhœa and irritability of stomach continued, and were accompanied with loss of flesh. The cause of the diarrhœa was not evident. The respiration was coarse at the apices of the lungs, but he had no cough. The abdomen was collapsed, and free from pain; no tumour nor

abnormal condition, could be detected on careful manipulation; his tongue was moist, and not injected, and there was no hoarseness. His urine was non-albuminous; sp. gr. 1014. The diarrhœa and disturbance of the stomach continued; chalk, kino, opium, copper, oxide of silver, were prescribed; the last appeared most effective; but although the diarrhœa ceased for a short time, he did not appear to derive nourishment from food; an attack of diarrhœa came on a few days before his death, and he died October 21st, 1855. *Inspection* twenty-six hours after death:—The body was extremely emaciated, and the eyes were sunken. In the chest there were slight pleuritic adhesions at the right apex; the lungs were collapsed; at the extreme apices there was old iron gray deposit, with some low organized product; but no vomica nor miliary tubercles existed. At the lower lobe of one lung there was red hepatization, affecting a large proportion of the lobe. The bronchi were slightly granular, and contained frothy mucus; but the larynx was healthy. There was considerable injection of the mucous membrane at the root of the tongue. The bronchial glands were normal. The heart was small, without fat, and healthy. The *abdomen* was exceedingly collapsed. The transverse colon was curiously twisted, so as to make a sigmoid curve in the epigastric region; it was the only portion filled with gas; the rest of the intestines were empty. The *stomach* was pale, and its mucous membrane normal. In the *duodenum*, the glands were very distinct; in the jejunum the mucous membrane was gray; and in the *ileum*, the mucous membrane was intensely congested; one of Peyer's patches, about two feet from the cæcum, was ulcerated; and the membrane in several parts had a thin adherent brownish covering, as of epithelium stained by adherent fæces. On examination this was found to consist of columnar epithelium. The cæcum was intensely congested, and its membrane was ecchymosed. The surface presented epithelium, blood, and some mucus; and the capillaries were full of blood. The colon was in a similar state, but rather less intensely congested; no ulcer could be found. The appendix cæci was full of fæces. The mesenteric glands varied exceedingly in size, from that of a pea to a pigeon's egg; some were swollen, red, and oedematous; others contained yellow strumous product; and in some parts beneath the peritoneum of the mesentery were minute tubercles; there was no evidence of pressure on the thoracic duct; and the pancreas appeared small, but healthy. The kidneys and supra-renal capsules were normal; the spleen was healthy; the liver was also normal in size, and healthy; the gall-bladder was much distended. On examining the semilunar ganglia, the cells were seen to contain much pigmental matter, but no disease could be found.

This case was believed to be one of phthisis, in which there was extensive ulceration of the colon; such diseased state was, however, not found after death; disease of the mesenteric glands appeared to have been the original malady; and this was the explanation of the gradual emaciation. The diarrhœa was the result of subacute disease of the mucous membrane, and of

ulceration of the ileum, but it was increased by the great congestion of the mucous membrane of both small and large intestine. This was the most marked symptom, and tended more than any other to exhaust the patient. The ordinary remedies were unavailing; but of these, the oxide of silver, with conium, appeared to be the most effective. An attack of acute pneumonia was the immediate precursor of a fatal termination.

2. In *disease of the mesenteric glands* a low organized product is effused into the glands themselves, probably between the chyliferous ducts, which become entirely obliterated, and the structure of the gland thereby destroyed. Extensive disease of these glands prevents the absorption of chyle into the system. The glands show the disease in various stages and gradations; in some but scanty abnormal product is poured out, in others the whole gland is destroyed and very much enlarged, constituting a whitish mass, the size of a pigeon's or a hen's egg. The effused product consists of granular blastema and of imperfectly developed cells; and the swollen and injected state of less diseased glands appears to indicate that inflammation or hyperæmia is associated with the morbid change. The increase in size takes place by additions at the periphery of that already deposited, and degeneration soon follows in the centre, from the scanty supply of nourishment afforded to that part. The gland is sometimes enveloped by a firm fibrous cyst, consisting of inflammatory and better organized product, which has assumed the character of fibrous tissue; whilst the centre consists of calcareous deposit, the albuminous portion having been absorbed, and the inorganic only left. Degeneration of another character, however, takes place in the effused product; it is converted into a mass of granular molecules and highly refracting particles, constituting small cheesy tubercles of a yellow colour, or a softened and semi-diffuent mass. Thus, different stages of disease and degeneration may be presented by almost contiguous glands. Sometimes the changes are of an active character; in others it appears probable that calcareous deposit has remained passive for many years. Whilst these changes are going on in the glands, the lacteals between them become enlarged and distended with

similar strumous product, or we can trace the distended ducts to the intestine, where they ramify on its surface; and at this part we generally find a cluster of tubercles and ulceration of the mucous membrane; were it not that the glands appear to be in a state of more advanced disease than the intestine, we should suppose that the strumous ulceration of the mucous membrane was followed by the absorption of like product and by glandular disease; the *absorption* of nutriment is thus more or less completely prevented. The peritoneum is sometimes studded with miliary tubercles, or we merely find minute clusters in the peritoneum opposite points of ulcerated intestine. Inflammatory disease is also found in the serous membrane in various degrees, either constituting bands of adhesion, or uniting the intestine in one mass. (See Strumous Disease of the Peritoneum.)

Symptoms.—Diarrhœa, as we have previously mentioned, is sometimes one of the symptoms of mesenteric disease. There is gradual wasting, from the obstruction of the chyle vessels, and from the cessation of the supply naturally poured into the thoracic duct. The patient has an anxious expression of countenance; there is dryness of the skin; injection of the tongue, which is more or less furred, and a craving appetite; the desire for food is insatiable. The bowels are irregular, being either purged or constipated, and there are occasional attacks of severe pain, and the evacuations have sometimes been found to contain a large quantity of fatty matter.

The marasmus gradually becomes extreme, and, whilst the limbs are wasted, the abdomen is considerably enlarged, and protuberant. The abdomen is full and rounded, but it rarely happens that the enlarged glands can be felt on tactile examination; we more easily discover them in the neck and in the axillæ. Where peritonitis, or ulceration of the intestines, has been produced, pain is a more common symptom.

A fatal termination may result from diarrhœa; or other organs become implicated, as the brain and the lungs; causing death by tubercular bronchitis, by convulsion, or hydrocephalus. In other instances disease in the epiphyses of the bones takes place, but the patients in these cases present less advanced disease of the glands of the mesentery.

Death does not always follow this condition of the chylopoietic glands unless the disease be very extensive; we have evidence of this fact in the calcareous condition in which they are found when death has followed from other causes, as from phthisis, or from tubercular arachnitis; but the interference to the elaboration of chyle increases the tendency to the deposition of strumous products in other parts.

At the period when this less severe disease existed, and the subsidence of which had left the calcareous state just mentioned, gastro-enterite had probably occurred; or, without any febrile excitement, the child had been observed to be imperfectly nourished, its growth retarded, and its nutritive power evidently feeble. It is in this early stage of the disease that proper attention to the health of the child may correct commencing degenerative changes, which will, if fully developed, necessarily prove fatal. Too frequently, however, the physician is consulted when the opportunity of checking morbid action has passed by.

CASE CXXI.—*Strumous Peritonitis.*—The following case was under my care, and was stated to be one of *labes mesenterica*. The child, Sarah G—, was nine years of age, of a very strumous appearance, with bright eye and fair complexion; for about a month the abdomen had become gradually enlarged, the child wasted, the bowels were variable, sometimes loose, at other times constipated, but without any severe pain or febrile excitement. Nothing could be felt on tactile examination, nor could fluctuation be perceived; an unfavorable prognosis had been previously given. Although, however, we had evidence of strumous diathesis, and the existence of deposit in the peritoneum and in the mesenteric glands was rendered probable, there was no evidence of extensive mesenteric disease. With nourishing diet, and by cod-liver oil, with steel wine in drachm doses several times a day, the child rapidly improved.

It would be incorrect to consider this either as strumous peritonitis or mesenteric disease cured, but it presented the early stage of strumous disease of the abdomen, in which we may expect to be of service to our patient; and it would be very easy to adduce many instances of this kind.

Diagnosis.—In its earliest condition strumous disease of the mesenteric glands may easily be mistaken for simple diarrhœa, or gastro-enterite; and what is of greater importance, the sympathetic affection of the brain sometimes renders it exceed-

ingly difficult to distinguish between strumous disease of the abdomen and hydrocephalus. In the former there may be cerebral oppression, grinding of the teeth in sleep, starting, occasional vomiting, and convulsion; but in hydrocephalus the mind is generally less active, there is strabismus, or evident abnormal condition of the pupils; the abdomen is collapsed, rather than distended; there is greater unwillingness to exposure of the skin to cold air, the bed-clothes are drawn firmly down when the patient is sensible; and again, the extreme capillary circulation is more checked, and the vessels yield easily to distension, so that frequently on drawing the finger across the skin, a deep line of congestion remains for a short time; this indication, however, is a very imperfect one. In the cerebral disease vomiting is often a marked symptom, especially at an early stage.

In the cachexia produced by enlarged spleen, by miasmatic disease, by lardaceous disease of the liver, or of other glands, by disordered viscera associated with worms, symptoms arise simulating in some respects mesenteric disease. Lardaceous disease, though closely allied to struma, exists without deposit in the mesenteric glands, and is equally unyielding to treatment. The history of the case, and the presence of enlargement of the liver or spleen, assist our diagnosis.

In worms, the symptoms are pallor, irregular bowels, wasting, distended abdomen, and voracious appetite; but there is often present more irritation about the nose and anus than in mesenteric disease; there is less emaciation, and the amenability to treatment is greater.

Strumous peritonitis is frequently associated with mesenteric disease, and is with great difficulty distinguished from it. The abdomen is less supple when the peritoneum is implicated, and when the disease is advanced the intestines move *en masse*; there is tenderness and distension, the pain is more severe; but the emaciation is less manifest. Ulceration of the small or large intestine and diarrhoea may be present in either disease, but very many of the cases usually designated *tabes mesenterica* are really strumous peritonitis.

In strumous subjects, however, after gastro-enterite, or slight peritonitis, the intestines sometimes become much distended

with flatus, and these cases at first sight resemble ascites; a very unfavorable prognosis may be given, whilst with rest, good air, cod-liver oil and steel, and occasional alteratives, the health becomes established, and the distension and pain disappear. The insidious character of strumous peritonitis must be well remembered; pain may be *entirely* absent, and the emaciation steadily progressive.

The *prognosis* in well-marked cases of mesenteric disease must be exceedingly unfavorable; for when there is general affection of these glands the obstruction to the introduction of food into the system is scarcely less complete than in direct pressure on the thoracic duct. Numerous inspections after death, however, show us that there may be degeneration of many of these glands, which become calcareous and pass evidently into a passive state, whilst others of them have been restored to their normal condition, and life may thus be prolonged for many years, till strumous disease in some other form, or another malady, prove fatal.

The age most liable to mesenteric disease is infancy, the period from the first to the completion of the second nutrition; in those who attain to early manhood it is much more frequently found associated with strumous peritonitis and with phthisis.

The *causes* of this disease have been previously mentioned; they are hereditary predisposition, improper food, insufficient rest, the want of cleanliness and light, the exanthems, as measles, scarlet fever, and small-pox, exposure to cold, and to a damp, humid atmosphere, and probably also congenital syphilis. Children brought up by hand are more liable to the disease; each of these causes diminishes the nutritive energy of the system, and a slight exciting cause then becomes sufficient to set up the disease; and when it is developed to accelerate it, so that it becomes quickly manifest in a marked degree.

Simple mesenteric disease is, however, rare, even in strumous subjects; and in many cases of strumous peritonitis, and of phthisis with ulcerated intestine, the glands are unaffected.

Treatment.—It must always be remembered, that in this disease waste advances and increases, whilst the supply of reparative material to the blood is cut off.

Our chief aim, therefore, must be to facilitate and assist nutrition; whatever is given must be easy of absorption and assimilation.

lation, as we have stated in speaking of the diarrhœa of strumous children. Wine may be often taken with advantage, for alcoholic liquors probably prevent waste going on with such great rapidity, and they are readily absorbed by the venous capillaries without the more gradual entrance into the blood by the lacteals of the villi. The elaboration of the chyle cannot be effected, however, by venous absorption, and the blood is imperfectly restored.

If there be febrile excitement, salines, as the bicarbonate of potash, or the citrate of ammonia, in doses of a few grains, may be administered; and when great restlessness exists, gr. j or ij of Dover's powder, or ℥ij or v of the solution of hydrochlorate of morphia, are of service.

The use of mercurials is better avoided; but when the motions are clayey and pale, and the bowels constipated, their use, as the grey powder, or calomel with carbonate of soda, is occasionally beneficial.

A change to sea air is very desirable in the early stages of mesenteric disease.

With cod-liver oil we may with advantage combine iodide of potassium, iodide of iron, &c. The preparations of iron, however, cannot in many cases be taken; pain in the bowels is produced; but this is less likely to follow the use of steel wine, and the saccharine carbonate of iron, than the stronger compounds.

Iodide of potassium with solution of potash, and with very minute doses of opium if necessary, continued for a lengthened period, are sometimes of considerable service.

As to external remedies, the tincture of iodine may be painted over the abdomen, or strips of the ammoniacum plaster with mercury applied; but the maintenance of an equable and warm state of the skin is of greater importance than external medicinal application.

3. *Tubercles in the peritoneum and strumous peritonitis.*—The state of the peritoneum is so closely associated with that of the alimentary canal, that a consideration of strumous disease affecting that part renders some notice of the diseased peritoneum necessary.

It is manifested in several conditions.

1. Miliary tubercles covering the general surface of the peritoneum—visceral, parietal, and mesenteric.

2. Strumous deposit on the peritoneum and in the glands, associated with inflammation leading to matting together of the intestines, sometimes to perforation from without, and to the formation of small faecal abscesses.

3. Peritoneal ascites with miliary tubercles.

4. Strumous peritonitis with tympanitis.

The presence of miliary tubercles on the peritoneum is found in many cases of phthisis, when there is ulceration of the intestine, the opposed surface of the peritoneum being covered with minute tubercles, apparently from the local congestion.

In children who have died from hydrocephalus, with miliary tubercles in the pia mater, or from acute pneumonia with miliary tubercles studding the whole of the lung, the peritoneum is frequently found affected in the manner described, but in other instances the affection of the peritoneum gives rise to the most prominent symptom.

The deposit is generally in semi-transparent grains, and appears to be situated in the substance of the peritoneum; it consists of an almost amorphous blastema with minute granules, and of imperfectly developed cells; but sometimes around the deposit elongated fibre cells and branching cells are observed.

In some cases of peritonitis, thin layers of lymph are deposited on the peritoneum, and the deposit assumes a minutely granular appearance, almost as if sprinkled with sand; these must not be mistaken for true miliary tubercles; they can occasionally be scraped off, and leave the serous membrane smooth beneath, but this cannot always be effected. The movement of one part of the intestine on another, and the gradual deposition of lymph, appear to produce this condition. A more severe form of disease is that in which, with strumous deposit, there is ordinary inflammatory change; the intestines become matted together by lymph, and by low organized product, which rapidly undergoes degeneration, constituting cheesy masses. These masses are deposited between the coils of intestine, in the omentum, and in the adhesions themselves; so that we find the peritoneal tunics of the liver, spleen, &c., considerably thickened—three to five lines, or more—and in the fibrous tissue constituting the firmer

part of the deposit are tubercles or strumous infiltration. The stomach rarely, if ever, presents strumous degeneration on its mucous surface, but it is not uncommon to find tubercles on its peritoneal surface. The mesentery and its glands are also generally found in a similar condition.

The product thus effused leads to union of the intestine, one part with another, so that the peritoneal cavity becomes entirely obliterated; fresh deposition takes place as the disease advances, and the tendency to degenerate increases. The masses soften down, the peritoneal and muscular coats ulcerate, and this continues till the mucous surface gives way, and an opening is formed into the intestinal canal. The perforation takes place from without, beginning at the peritoneal surface. This perforation does not, however, lead to more extensive peritonitis; the firm adhesions which have already taken place prevent effusion, and no extravasation follows; or a very small faecal abscess is the result. The extension of ulceration amongst contiguous coils of intestine sometimes leads to several portions becoming completely truncated and opening into a faecal abscess, in which six or eight communications may exist. I have examined several cases in which there were from twelve to fifteen communications between portions of the intestine in different parts, but without faecal abscess. Should the adhesions be less extensive, perforation will produce more marked symptoms, if a fatal result does not very quickly follow. It is generally the small intestine which is found perforated; but in others the small intestine opens into the colon; in one I observed the jejunum communicating with the transverse colon.

It is more rare in strumous than in cancerous disease of the abdomen and intestines, to find faecal abscesses followed by perforation of the abdominal parietes. In struma the disease is often very general, and several faecal abscesses exist; but the adhesions and secondary perforations allow the contents of the canal to be transmitted; in cancer the ulceration is more localized in character, and gradually extends through all the contiguous structures. In Case CXXII, in a child aged six, strumous disease of the abdomen was followed by perforation of the parietes.

The effusion in some instances is of an ascitic character, and dropsy is the result. The peritoneum is thickened, clear serum

is effused, and with it more or less strumous product. This is not a rare disease among children, but it is of a slow, insidious character, and very intractable. It sometimes exists with a lardaceous condition of the liver or of the spleen; but this is not always the case, nor is it always preceded by exanthems.

In some instances of strumous peritonitis the intestine appears to lose its contractile power, and yields to distension, so that most distressing tympanitis takes place, or there is simple distension *without pain*, the muscular fibre having lost its power to contract.

The coats of the intestine become so much softened that after death they readily separate the one from the other, and may be torn in long shreds. Dr. Hodgkin placed in the museum at Guy's several specimens showing this condition in a remarkable degree.*

The *symptoms* of this form of disease are also sometimes obscured at the commencement; with well marked strumous diathesis we have pain in the abdomen of a severe character, resembling colic; and it is accompanied with considerable tenderness; diarrhoea and febrile excitement come on, with injected, slightly furred tongue and with distress of the countenance. There is a circumscribed flush on one cheek. Under suitable treatment and precaution the active symptoms subside, and the patient feels relieved; in a few days or weeks, however, the pain returns, and there is renewed aggravation of the symptoms and of the febrile state; it may be that a defined mass is felt in the abdomen, in the umbilical, hypogastric or iliac regions; the tumour is tender on pressure, and imperfectly resonant on percussion.

These attacks are repeated from time to time, and the diarrhoea becomes severe, and occasionally is accompanied by vomiting. The body wastes, but the abdomen is large, and in most cases loses its suppleness, and it moves *en masse*. The tongue becomes more injected, oftentimes red and morbidly clean. The strength of the patient is broken, severe hectic is set up, attacks of pain are more frequent, and portions of the abdomen become exquisitely tender; at length the exhaustion becomes extreme, or more general strumous disease is set up; thus the

* Hodgkin, 'On Mucous and Serous Membranes.'

brain becomes affected by the deposition of strumous deposit, and slight coma or convulsions come on before death.

The symptoms somewhat resemble those of mesenteric disease; the abdomen is hot, often distended, and tympanitic; the recti are rigid; the body wastes, the countenance becomes anxious, and the eyes sunken; the patient is fretful and irritable; the bowels often act with irregularity; the pain is sometimes a marked symptom, but it is often absent, or merely resembles occasional colic. The wasting of the body is however less than in severe disease of the mesenteric glands.

The second form is that in which there is less serous effusion, but the strumous product is accompanied by greater inflammation, lymph is effused, and the intestines matted together by bands of adhesion, or there is cough from strumous disease of the lungs; but this terrible aggravation of the suffering of the patient is generally absent, for the pneumonic disease remains latent.

The severe attacks of pain often indicate the formation of faecal abscess or fresh accessions of inflammation.

In caecal disease and in phthisis we have dwelt on the sudden peritonitis, which is sometimes set up in subjects affected with those diseases.

Where there is considerable effusion into the peritoneal cavity, the symptoms may be exceedingly insidious, enlargement of the abdomen taking place without manifest febrile symptoms, pain, or tenderness; fluctuation is readily perceived; and generally without enlargement of the liver or spleen, and without thoracic disease or albuminuria. The patient becomes anæmic and emaciated, pain is occasionally paroxysmal, or less severe but continued. Strumous disease of other organs generally follows, and leads to fatal results.

There is great difficulty in producing absorption of this peritonitic fluid; for the peritoneum is in a passive condition, and medicines which act on the excretory organs, as diuretics, solution of potash, iodide of potassium, often fail in the desired effect; and mercurials tend in many instances to increase the effusion.

Causes.—Children in their first dentition, and at the age of puberty, in whom the rapid developmental changes are perverted by struma, are very prone to this disease, but at early manhood from 15 to 25, or 30, we observe many instances of it.

The predisposing causes are those of strumous disease generally—hereditary predisposition, unwholesome food, the want of cleanliness, a damp humid state of atmosphere, exposure to cold, insufficient light, &c.; light is as essential to healthy growth as cleanliness; but unfortunately the absence of the one often entails the loss of the other. The dark offensive dwellings of poverty present terrible manifestations of the sources of struma; but with the rich, hereditary tendency, exposure to cold, &c., are sufficient, with very slight exciting causes, to induce these affections.

The hyperæmic state of the ovaries at commencing menstruation is sometimes sufficient to determine the deposition of strumous product, and to cause strumous peritonitis extending then from the pelvic viscera; this is especially the case when menstruation in young subjects has been checked by exposure to cold, &c.

Peritoneal disease of this form is sometimes excited by blows or falls on the abdomen, or it follows diarrhœa from injudicious food or excess; we observe it also as a sequence of typhoid fever; the follicular ulceration of the intestine and irritation of the mesenteric glands being followed by strumous disease.

The *diagnosis* has already been spoken of in mesenteric affections; we may further add that when strumous disease of the abdomen is associated with like disease of the brain, the symptoms are often more than usually obscure; cerebral symptoms, such as delirium and coma, may be followed by prostration, distension of the abdomen, vomiting, &c.; and the general aspect of the disease may closely resemble typhus fever.

The *prognosis* in well-marked cases is very unfavorable; but at an early stage much may be done to render the changes which have occurred passive, and to prevent the accession of fresh disease.

Treatment.—The indications of treatment are very similar to those mentioned in mesenteric disease. As far as possible the exciting cause of the disease should be taken away; and what is less practicable we must attempt, namely, the removal of the condition which constitutes the disease—strumous degeneration; this may in part be effected by sea air, by iodine, cod-liver oil, iodide of potassium, and the milder preparations of steel. Nourishment should be freely given, and of a character that can

be easily assimilated. Improper food may induce most severe colic, and defeat all remedial measures.

The inflammatory state is best counteracted by the application of leeches and counter-irritants; as by cantharides, or iodine, applied externally; in children, it is well to place a portion of tissue paper between the cantharides plaster and the skin, and only to apply the plaster for two or three hours; or to use for a short time the acetum cantharides; or still better, the Cantharidine Blistering Tissue.

When fluid exists, diuretics may be tried, but they are not of much service. Great care is required in allowing exercise, because at the same time that fresh air and change are exceedingly desirable to improve the health, rest is most important for the abdominal organs themselves. Slight movements may break down adhesions, and lead to rapid extension of disease, and to a fatal termination. The importance of rest to the viscera of the abdomen can scarcely be urged with sufficient force.

Again, it is most desirable that mercurial medicines and drastic purges should be avoided; the gentlest laxatives and mild enemata are all that is required.

Various symptoms arise which demand almost daily attention, as diarrhoea, occasional vomiting, each of which must be checked by appropriate treatment. In a case of strumous peritonitis recorded by Dr. Hughes in the Guy's Reports of 1856, creasote was prescribed to check severe vomiting; it produced urine almost of the colour of indigo. Pain must be moderated by opium in small doses, or by morphia; by warm fomentations or cataplasms; by chloroform; or by belladonna applied to the parietes.

Gentle pressure by a bandage, and the irritation of the ammoniacum with mercury plaster, are sometimes of service; and in those cases in which fluid is poured out, tapping is sometimes advisable; for in chronic peritonitis the serous membrane often becomes a thickened, and almost a passive sac.

CASE CXXII.—*Strumous Peritonitis. Fæcal Abscess. Artificial Anus.*—A little girl, æt. 6, had been suffering from chronic peritonitis for about a year. The abdomen became much distended, and there was severe vomiting, with great emaciation. Six months before death, a circumscribed

tumour formed near the umbilicus; this afterwards broke, and discharged *faeces*.

On inspection, the lungs were found studded with tubercles; the intestines were adherent, and several portions were perforated; they had formed a *faecal abscess* near the umbilicus, which had discharged externally. The Fallopian tubes were filled with soft strumous product. (Prep. in Guy's Museum, 2446⁹⁰.)

4. *Disease of intestine in phthisis pulmonalis*.—The mucous membrane of the intestine is frequently the seat of tubercle; and although they are rarely, if ever, found in the mucous membrane primarily, they are often associated with strumous disease of the lung, of the serous membranes, of the brain, or of the bones.

A very common position for this deposit to occur is in the substance of the mucous membrane, at the lower part of the ileum, and generally in the aggregate or solitary glands. The deposition takes place, as in the lungs, without any evidence of inflammatory disease, and the tubercles are found as minute grains, one-sixteenth to one-eighth of an inch in diameter, and of an opaque, cheesy appearance. On the examination of these tubercles they will be found to consist of an immense number of granules of fat, with imperfect nuclei; in other instances the centre is semi-fluid, softened down; still more advanced, we find that the slight covering of the mucous membrane has given way, and a small ulcer is formed, with a depression in its centre, and with an irregular and slightly excavated margin. This deposition is probably in most cases preceded by hyperæmia of the mucous membrane, or by inflammatory action; and although it appears nearly established that inflammatory action is not essential to its deposition, this state tends rapidly to accelerate the ulcerative action, and the repeated deposition of strumous product at the margin of the ulcer. When active inflammatory action has taken place, the edge of the ulcer is found to be injected and irregular, and to have extended rapidly; in some cases, also, sloughing has taken place. The extent of this diseased state varies exceedingly; generally only the lower part of the ileum is affected, and next in frequency the ileum and the cæcum.

With these parts, the colon is sometimes diseased in its whole

length; and lastly, also, the other portions of the small intestines, the jejunum, and even the duodenum.

We frequently find, that, at the base of the ulcer, immediately beneath the peritoneum, there are numerous minute tubercles, apparently caused by the congestion of the ulcer tending to this increased deposition of tubercle. In other instances the mucous membrane is raised, and presents a small swelling, about a quarter of an inch in elevation, and a quarter to half an inch in diameter; and on making a section of this minute tumour, a collection of pus is found in it; a sort of small abscess in the mucous membrane. (*See Cases of Disease of the Cæcum.*)

But strumous ulceration of the intestine, when associated with phthisis, sometimes manifests itself differently. There is scarcely any diarrhœa, but sudden intense pain is followed by collapse, and too often by fatal peritonitis. A minute ulcer has increased in depth, so as to extend through the muscular coat, and then through the peritoneum. It may be that this peritonitis is localized, or that a fæcal abscess is formed, and of these we shall have to speak more fully. The affections of the appendix cæci will also require a fuller notice.

In other instances the diseased intestine is found to be in a healing condition, while the affection of the lungs has steadily progressed, or, having become rapidly aggravated, has led to a fatal result. I have several times seen cicatrices in the intestine in phthisis, when there was no evidence to show previous disease of a different kind, as typhoid fever. In one instance, admitted into Guy's Hospital several years ago, there were symptoms of intestinal, and it was feared insuperable, obstruction; but the patient was spared to linger on for many weeks, and died from phthisis pulmonalis. A cicatrix was found in the ileum, leading to very considerable contraction of the intestine, and no doubt it had been the cause of the previous symptoms of obstruction.

In some patients in whom the jejunum and ileum have been ulcerated throughout, with less affection of the colon, the diarrhœa has been exceedingly severe. When the mesenteric glands are also affected, we have several times observed, extending from an ulcer in the jejunum or ileum, distended lacteals, reaching to the infiltrated glands, and filled with strumous

product. Some regard the ulceration of the intestine as having set up the disease in the lacteals and glands; others consider that the gland was primarily diseased, and that the obstructed lacteals and local congestion consequent upon it have set up the ulceration at that part of the intestine. Simple distension of the lacteals is more common in cancerous disease, producing pressure on the thoracic duct; in struma, abnormal product *fills* and enlarges the lacteals.

In many cases of phthisis the intestine appears to have taken part in the general atrophy of the whole body, and we find the coats of the intestine much thinned, of a pale colour, and even semi-transparent. Again, the mucous membrane sometimes does not present any lesion, although the patient may have had severe diarrhœa.

The extent and severity of the affection of the intestine are very diverse. In cases where the phthisis is of a pneumonic character, when there is extensive effusion into the lung tissue, rapid disorganization, considerable fever, and speedy termination, the intestines are sometimes unaffected. It is in more chronic cases that we generally find this condition most marked.

In one hundred cases of phthisis only thirteen had the intestines healthy, and those were of the character just mentioned, namely, pneumonic phthisis. In sixty-nine cases the ileum was diseased, and generally the colon also, in a greater or less degree; in seventeen cases the colon only was diseased. The ileum is the part most frequently affected. In more severe cases, the colon is also diseased, sometimes in its whole length, or merely at the sigmoid flexure; or we find the whole alimentary tract diseased, and the jejunum, ileum, and colon are all ulcerated and inflamed.

Attacks of diarrhœa generally alternate with constipation in phthisical disease of the intestine, and thin bilious evacuations are occasionally mixed with blood. The discharge from the bowels is sometimes composed of mucus passed in long strings or casts, or it presents the character of yeast; in a case of this kind under my care, the evacuation closely resembled the matters discharged from the stomach in obstructed pylorus, but with a fœcal instead of a sour odour. Under microscopical

examination minute cells and grains in state of change were observed, but not the ordinary torula, nor the sarcina ventriculi. In other instances the disease resembles acute dysentery, blood and mucus are passed, with considerable tenesmus; there is slight griping pain, but the discharge from the bowels resists all treatment; it may be checked for a few days, but again returns, and it is remarkable in some cases how completely the thoracic symptoms are in abeyance; no cough, dyspnoea, pain, nor distress about the chest may trouble the patient, although after death considerable vomicae may be detected in the lung. In some of these instances the appearances of the colon are quite those of a dysenteric character, the extent of the ulceration destroying in some cases the mucous and muscular coats, leaving but small islets of injected mucous membrane; in other instances the surface is covered by diphtheritic membrane, and presents isolated patches of superficial ulceration beneath. Many of these appearances have been observed in the numerous cases of phthisis which die at Guy's Hospital, and it is probable that the more damp air of the Borough of Southwark, the ill-ventilated homes in Bermondsey and Rotherhithe, from which some of these patients have come, have induced this dysenteric state.

It would appear that exposure to cold and wet is sometimes the cause of the unusual severity in the affection of the alimentary canal in phthisis. In other cases, the administration of mercurial medicines, of drastic purgatives, and of improper food, induces this condition.

The presence of fistula in ano, as a complication of phthisis, is frequent; and it is a question upon which opinions are varied, whether the division of the sphincter is advisable. Most surgeons at the later stages would dissuade from the operation; but in the earlier condition, before there is any disorganization, the removal of a depressing and exhausting discharge may tend to re-establish health; sometimes, however, the pulmonary symptoms rapidly increase after the operation.

With albuminuria in strumous subjects, disease of the colon leads sometimes to severe diarrhoea and great exhaustion. The association of phthisis with renal disease is not of very frequent occurrence; it may be the ileum and colon, as in ordinary

phthisis, which are ulcerated, or it may be the rectum which is especially diseased. The use of purgatives to relieve anasarca has been followed in some of these cases by serous diarrhœa, which is very intractable; inflammation and ulceration are set up, and, like ulceration on the extremities in dropsy, may be the immediate cause of death.

Treatment.—The 1st object should be as much as possible to remove the exciting causes of diarrhœa; 2nd, to check irritating secretions by correctives and astringents; 3rd, to soothe the inflamed membrane by demulcents and by opiates.

In most cases the avoidance of indigestible food, of uncooked fruit, and of malt liquors, is sufficient to check the purging in ordinary phthisical disease of the intestine; or an injection of starch and opium may be used with benefit. Suet and milk is an unirritating form of nourishment, and is often of much service; other demulcents are used with advantage; and when the powers of the patient are much depressed, port wine or brandy must be prescribed.

Opium alone, or in combination, is of great value, as in Dover's powder, the compound kino powder; and it may be given with acetate of lead, with sulphate of copper, with bismuth, or with oxide of silver. Bismuth alone will oftentimes quiet this irritable condition of the alimentary canal; alkalies, also, and astringents, as the compound krameria mixture and the compound logwood mixture of the Guy's Pharmacopœia, will often be found valuable remedial agents.

Although in some cases cod-liver oil acts on the bowels, in other instances it moderates diarrhœa, and the bowels act with less violence and pain.

I have found the injection of borax with barley water, or of powdered charcoal with the same agent, of more service in some cases where the colon is much affected, than simple starch with opium.*

If there be severe pain the application of hot cataplasms or of mustard affords partial relief.

It is the exception to find phthisis free from abdominal complication, but the following instances present some peculiarities in reference to this affection: in Case CXXIII, the mesenteric

* See Dr. Th. Thompson, 'On Consumption.'

glands were very extensively diseased, and the lacteals distended with strumous product; the diarrhœa was exceedingly obstinate, and hastened the fatal termination. In Case CXXIV, the pulmonary symptoms were entirely masked, but there is no doubt that the dysenteric inflammation was more intractable in character on account of the disorganization of the lungs. If there had been no inspection after death, the latter would probably have been considered by many practitioners, who did not take the trouble carefully to examine the chest, as simple disease of the intestine. Each case of phthisis must be considered in itself; the different degrees of pneumonic inflammation, of the laryngeal disease, and of glandular or abdominal complication, &c., render the secondary symptoms exceedingly modified and varied, whilst the broad general characters bear very close similarity; much relief may be afforded by suiting the treatment to these sources of discomfort and danger.

CASE CXXIII.—Strumous Disease of the Mesenteric Glands. Obstruction of the Lacteals. Ulceration of the Small and Large Intestine. Dysentery. Phthisis.—William S—, æt. 20, was admitted into Guy's Hospital, August 29th, and died November 1st, 1855. He had been a labouring man at Hoxton; and, with the exception of a slight cough, had enjoyed good health till the January previous; he then had severe cold, and his cough increased in severity; he was confined to his room for two months, and expectorated viscid mucus; and he had gradually become more feeble and emaciated till his admission.

His chest was narrow and contracted; there was dulness on percussion below the clavicles; and in the supra- and infra-scapular regions there was also roughness in the respiratory murmur, with bronchial respiration, more distinct on the right side than on the left, and with increased resonance of voice. The pulse was weak and frequent; the tongue had a white fur upon it; the bowels were relaxed, but the appetite was good. Catechu, with extract of poppies and cod-liver oil were given, and morphia at night.

The diarrhœa continued with short intermissions, and his affection of the throat increased; he became extremely emaciated, and died November 1st.

Inspection twenty-one hours after death. The body was much emaciated; and on the lower extremities presented several spots of purpura. The larynx was extensively ulcerated; the inner surface of the epiglottis was covered by follicular ulcers, which were united together. There were no pleuritic adhesions, and the lungs collapsed well; at the apices were several vomicæ, and throughout both lungs were numerous tubercular deposits and miliary tubercles. The bronchial glands were much enlarged; and they were infiltrated with strumous product.

Abdomen.—The intestines were tolerably distended; the peritoneum pre-

sented granular tubercular deposit, and there was considerable injection of it at the parts of small intestine opposite to ulcerated portions of the mucous membrane; the mesenteric glands were very large and prominent, of a yellowish-white colour, and infiltrated with low organized product; some of them were the size of a pigeon's egg, and they occupied the whole of the mesentery; in several parts of the small intestine, lacteals were observed to extend from the enlarged glands to the walls of the intestine; these vessels were white, irregularly distended, and in some places had a moniliform appearance; they extended in several places upon the walls of the intestine, and beneath the mucous membrane to ulcers situated there. On opening the small intestine, numerous ulcers were observed; they commenced in the upper portion of the jejunum, and extended with greater or less intervals to the cæcum; some were one and a half inches in length; their margins were congested, and were irregular and undermined; their surface was granular, as if presenting minute strumous deposit; the ulcers were scattered about six inches apart, and were larger at the jejunum than in the ileum; strumous tubercles were observed in many parts of the ileum in the substance of the mucous membrane, and there were several minute ulcers about the size of peas. The ileo-cæcal valve was much congested, and was swollen and œdematous. The whole of the cæcum and colon had a remarkable appearance; with the exception of a few islets of raised congested membrane, the whole surface, as far as the sigmoid flexure, was destroyed; the surface was of a whitish granular appearance, presenting some congested points, or irregular pits; the section showed that there was low organized product in the superficial layer; some true tubercles, and cellular tissue dipped down into the muscular coat; and on the surface itself there was granular and imperfectly formed cellular deposit, like diphtheritic membrane. The descending colon presented transverse irregular ulcers, with larger intervening spaces; the rectum was still less affected; the appendix was much distended at its superior two-thirds, and was ulcerated; it contained strumous tubercles. The white substance in the lacteals consisted of particles of fat irregularly aggregated into numerous spherical masses; and in the mesenteric glands was ordinary strumous and imperfectly formed cellular growth. The liver was normal and not fatty; the spleen was healthy.

CASE CXXIV.—*Ulcerated Colon. Phthisis. No Cough.*—Mich. M'C—, æt. 53, was admitted with violent purging, which had existed a week; much mucus was passed per rectum, but he had no cough. He sank in a very short time. The whole of the large intestine was intensely inflamed and ulcerated, and the small intestines congested; an old vomica at the apex of the lung was surrounded by iron-gray pneumonia.

The pulmonary symptoms were masked; he had no cough, but the severity of the abdominal symptoms, dysentery of an acute form, rapidly led to a fatal result.

CASE CXXV.—*Phthisis. Ulceration of the Rectum and Sigmoid Flexure. Hemorrhage from the Bowels. Ulceration of the Appendix Cæci.*—A. B—, was admitted into Guy's Hospital, under my care, March 18th, 1857. He was a

married man, of temperate habits, who had considered himself in health till one month before admission; his principal symptom had been discharge of blood from the rectum with diarrhoea; he had cough, had rapidly emaciated, and he had become completely blanched.

On examination of the chest we found flattening, with imperfect mobility of the left apex, and soft mucous râle, and with increased resonance of the voice. The abdomen was contracted, and free from pain. There was but little doubt that he had phthisis; there was no evidence of external hæmorrhoids, and the administration of the compound logwood mixture of Guy's at once checked the diarrhoea and the discharge of blood; and these symptoms did not recur. Emaciation, however, rapidly increased, the cough became more severe, and the evidence of disorganization of the lung was better marked; he died in one month; and for several days he appeared in articulo mortis.

On *inspection* the abdomen was unusually collapsed; old pleuritic adhesions were found on both sides; at the left apex was a small vomica filled with pus, and the whole of the upper lobe presented patches of red hepatization, minute tubercles, and iron-gray consolidation; at the upper part of the lower lobe there was also a vomica, and a similar condition existed as in the upper lobe. The right lung was less affected; the bronchi were filled with purulent mucus; the larynx presented a small ulcer on the inferior vocal cord, and the aryteno-epiglottidean fold was very oedematous; the heart was normal; there was no post-mortem solution of the stomach; the intestines, especially the small, were empty and contracted; and in the lower part of the ileum were a few tubercles, and commencing ulceration. The transverse colon presented a sigmoid twist near the spleen, and the ascending and transverse colon contained some scybala, and presented several ulcers, oval in form, about half an inch in breadth, with injected irregular margins. In the sigmoid flexure and rectum, the whole of the mucous membrane was injected; it was almost covered with patches of ulceration, and in some parts there were portions of adherent diphtheritic membrane. The appendix cæci was twisted in a sigmoid form; and at the right of the cæcum, near its terminal third, it became very much dilated; the mucous membrane at this part was entirely destroyed, and the muscular coat much hypertrophied. The kidneys, liver, and spleen were healthy.

The ulceration of the rectum and sigmoid flexure had led to the hæmorrhage which blanched the patient; in this state of exhaustion the disease of the lung rapidly advanced. It was not the part of the intestine usually affected in phthisis; and he had no pain, distension of the abdomen, nor severe tenesmus; diarrhoea, with discharge of blood, were the most marked symptoms of abdominal disease. The mesenteric glands were more than usually affected. The appendix cæci was so diseased that it would probably have led to extension into the peritoneum or into the cellular tissue, if life had been prolonged. The loss of

blood apparently hastened the diseased action in the intestine rather than diminished it; and although the purging was checked, the patient never appeared to rally to any extent. He was unable to take cod-liver oil, but appeared partially benefited by hydrochloric acid, with small doses of opium and calumba.

CASE CXXVI.—*Phthisis. Inflamed Hæmorrhoids. Sciatica. Extensive Ulceration of the Larynx and Intestine.*—George B—, æt. 57, was admitted into Guy's Hospital, under my care, October 21st, 1857. He had resided at Greenwich, and had been employed as a pilot. His appearance was that of a gray, aged, tall, but spare man; and his habits of life had been intemperate. A cough had troubled him for three years; and he had had *raucedo* for some time; some pain in the limbs also tried him, but his chief complaint was of severe pain in the course of the right sciatic nerve extending down the thigh, and the calf of the leg to the foot. There was no evidence of any tumour in the pelvis. The heart was normal, but the respiration was coarse at the apices of the lungs. During the first month after admission the pain in the leg and foot became intensely severe, and the patient described it as being of a burning character. He had an anxious expression of countenance, and complained very much of the throat, having a distressing cough, and great difficulty in swallowing. Nothing could be seen in ordinary visual examination of the throat, and the larynx was moveable; but much coarse, and tubular respiration could be heard at the right apex; the pulse was rapid, and he much emaciated. On December the 4th, he was more prostrate and anæmic, and the pulse was irritable, the throat and cough were less troublesome; but he complained of great pain at the anus, where there was a red oedematous mass of mucous membrane extending about an inch from the outlet, and in the central part, the membrane was of a deep livid colour, semi-gangrenous. This protruding mass consisted of inflamed hæmorrhoids and mucous membrane. Opium was administered, and wine, and externally decoction of poppies with alum, &c.; the bowels were freely acted upon. On the 12th the hæmorrhoids were less inflamed; there was no oedema, and the external skin was flaccid; but, internally a superficial slough appeared to have separated, leaving exposed a red surface. The patient, however, appeared very ill, his eyes were sunken; and he had great difficulty in swallowing, so that he was quite unable to take solid food, but there was also hoarseness. The resonance of the chest was imperfect at the apices of both lungs, and in the right axilla bronchial breathing was especially distinct; there was, however, considerable laryngeal obstruction, so that the entrance of air into the lungs was greatly impeded; he feared "lest he should be choked," and had no sleep; the pulse was compressible. Cod-liver oil was given three times a day, and Dover's powder in five gr. doses; and as nourishment, eggs, beef-tea, wine, &c., allowed. Wine and brandy produced so much smarting pain in the throat that he was unable to continue them, but the cod-liver oil was more

grateful than anything else. The hæmorrhoids were, with the bowels, constipated, but his prostration increased rapidly; gurgling became distinct at the apex of the right lung, but it was less evident on the left side. He gradually sank, and died on the 23rd. *Inspection* was made fifteen hours after death. The body was exceedingly wasted, and the arcus senilis well marked. The epiglottis was thickened; and the mucous membrane on the whole of its inner surface was destroyed as far as the superior vocal cords, leaving only one strip of entire mucous membrane. The aryteno-epiglottidean folds on both sides were extensively ulcerated; the destruction of the mucous membrane extended to the pharyngeal side; and on the left side on the level of the epiglottis, there was an irregular ragged ulcer on the posterior pillar of the fauces; the soft palate was a little swollen, but was otherwise healthy; the vocal cords and trachea were normal; the bronchi contained much thick mucus of a reddish colour. The pleura was firmly adherent on the left side; at the right apex the adhesions were only slight. The upper and lower lobes of the lungs contained small masses of yellow strumous deposit; at the apices were vomices, and in the lower lobes were portions of red and consolidated or of congested lung. The pericardium was adherent, and the heart atrophied; there were fibrinous clots in both the aorta and pulmonary artery; the valves of the heart were normal, except slight atheroma at the base of the aortic crescents. In the abdomen the intestines were moderately distended, and the chyle vessels enlarged. The stomach was normal; but there was extensive and irregular ulceration at the end of the ileum; other ulcers extended throughout the colon as far as the rectum, some of these were healing, others were irregular in form, and extended into the muscular coat; there was no disease of the appendix cæci. The liver, spleen, and kidneys, were healthy; the mesenteric glands were healthy, but there was no evidence of pressure on the thoracic duct; neither in the pelvis could any injury of the sciatic nerve be found.

In this case there are several symptoms well worthy of notice. The laryngeal affection was very severe, but it was not of the kind usually observed in phthisis, for the vocal cords were healthy, and the inner surface of the epiglottis and the glosso-arytenoidean folds diseased; the pharynx was partially involved; those conditions were the causes of the distressing dysphagia from which he suffered; still it was doubtful whether the laryngeal disease was modified by any earlier syphilitic taint. We have generally found, where the tip of the epiglottis has been extensively ulcerated, that solids are more easily swallowed than fluids, on account of their ready passage from the mouth over the diseased surface, without coming into actual contact with the ulcerated surface; in this patient solid food pro-

duced much pain, probably from the posterior pillars of the fauces being also ulcerated.

The phthisical disease of the lung was preceded by bronchitis of three years' duration, and, although the symptoms on admission were comparatively slight, they rapidly increased, and evidence was soon afforded of consolidation, and then of disorganization. Again, although the colon was extensively ulcerated, diarrhoea was never a distressing nor an intractable symptom. That of which the patient most complained on admission was intense pain in the course of the sciatic nerve; and it is probable, that this agonizing pain was produced by reflex irritation of the branches of nerve supplied to the anus; when the hæmorrhoids became acutely inflamed, this pain was mitigated, or at least less complained of, but afterwards, when they were relieved and cured, the sciatica disappeared. This cause of sciatica, namely, hæmorrhoids, has been mentioned by Sir B. Brodie, and also by Mr. Ashton.

These instances, and many others which might have been adduced, show the general constitutional character of phthisical disease; and that although it may manifest itself with greater severity in one organ than in another, we should closely observe the state of all the viscera, as having a most important influence on the curative condition of the disease; that these simultaneous developments of morbid action go on very insidiously, as in the diseased appendix in Case CXXV; for although on the verge of fatal peritoneal perforation, the disease would not have been known unless by post-mortem inspection; and lastly, that even when the general state of strumous disorganization may be past the stage of reparative action, much may be done in partially relieving distressing urgent complication.

CHAPTER VIII.

ON DISEASES OF THE CÆCUM AND APPENDIX CÆCI.

THE diseases of the cæcum and of its appendix, are of a character so peculiar and important, as to call for special consideration.

To a certain extent, the cæcum is apart from the direct current of the contents of the alimentary canal; for the valvular opening from the ileum enters at two to four inches from the lowest part of the cæcum; and the capacity of the cæcum is several times greater than that of an equal length of the ileum. Hence also the contents of the canal move more slowly; and at the same time they become less fluid in their character. The mucous membrane is here destitute of villi, but it is exceedingly vascular, and is furnished with numerous solitary glands. The appendix cæci is an elongated glandular sac, which opens into the intestine at the termination of the cæcum, and generally towards its iliac side. At this part of the intestine also the longitudinal muscular fibres assume a different arrangement; they here form three bands, which arise from the position at which the appendix is attached, and are continuous with its muscular layer.

The cæcum is situated in the right iliac fossa, and is only covered by peritoneum on its anterior and lateral surfaces; a considerable quantity of loose tissue separates it from the fascia, and from the nerves and vessels in relation with the psoas and iliacus muscles. The mobility of the cæcum is therefore considerably less than that of the jejunum or ileum; but in this respect there is much variation, for it is occasionally

4

found with a long mesenteric attachment, and the right iliac fossa is then completely covered by peritoneum.

This freedom is far from being of rare occurrence; it is probably of congenital origin, and is of great pathological importance; for, it allows the cæcum to pass into hernial sacs, and it modifies the position which the cæcum assumes in great distension and in obstruction.

In reference to the rotatory movements of the intestine, Rokitsansky describes three varieties: first, rotation of the intestine upon its own axis; secondly, upon the mesentery as an axis; and, thirdly, upon another coil of intestine. The cæcum may be regarded as moving either on its long or short axis. In the former, the colon will assume a direction more or less to the right, or even a horizontal course. (Fig. 1, Pl. IV.) This condition we have observed without any apparent impediment to the passage of the contents; but when there is habitual constipation, it favours distension of the cæcum, and disease of the appendix. By rotation, on the contrary upon its short axis, the appendix may be placed towards the anterior abdominal parietes, or be situated at the posterior aspect of the intestine; or again, the cæcum is found to be sometimes so twisted, that the ileum opens on the right side. This abnormal condition was noticed in a case where the appendix was directed towards the liver, and the transverse colon was adherent to the cæcum, forming a sigmoid curve in the centre of the abdomen. There had not been any evidence of obstruction, but if such had occurred, the diagnosis would have been greatly obscured.

When these movements are combined, not only is the ileum found opening on the right, or on the posterior aspect, but the head of the cæcum is completely inverted, so as to be directed towards the diaphragm. In this state constriction may readily be produced, as in Case CXXX, where an inverted cæcum became distended to the utmost, and led to a fatal result. (Fig. 2, Pl. IV.)

In the second form of rotation of the cæcum, namely, upon the mesentery as an axis, there seems to be a still greater probability of obstruction. This rotation can only take place when the cæcum is very free, and it may then pass into the pelvis, or be completely rotated. Case CXXXI presents an instance of

this kind. The cæcum was in the pelvis, enormously distended and ruptured; it was twisted upon its mesenteric axis, and the passage into the colon obstructed. These twisted positions of the cæcum, and its free mesenteric attachment, are generally found in strumous subjects; they predispose to disease of this part of the intestine, and to lodgments in and ulceration of the appendix. We have not, however, observed that they have any connection with any abnormal state of the rest of the intestine. In Case CXXXII—one of fatal obstruction—the cæcum was apparently twisted on the mesentery as well as on its own axis.

The appendix presents characters still more diverse; but some of these appearances are the result of pathological changes, which we shall presently consider. It is generally three inches in length, but is sometimes only one and a half inch, and at other times it equals five inches in length. It is attached on the inner aspect of the cæcum by folds of peritoneum, constituting a mesentery; but whilst it is generally free, it is occasionally hidden behind the cæcum, or curved in a sigmoid form to the right side.

The mesentery of the appendix is generally short, and is sometimes fixed to the brim of the pelvis, and the appendix is then pendant in the pelvis. Not unfrequently the mesentery of the appendix is attached to the lower part of the true or iliac mesentery; in such case the appendix runs parallel with the ileum for an inch or an inch and a half, and is then, beyond its mesentery, free. The direction thus assumed is that of the brim of the pelvis towards the left, and the appendix nearly reaches the attachment of the sigmoid flexure. When this condition is present, we frequently have a more or less dependent pouch formed between the appendix and the ileum, consisting of the folds of peritoneum. This pouch is worthy of notice, because from irregular pressure it becomes atrophied; then, probably, perforated, so as to form an opening through which other coils of small intestine may pass. The opening thus formed oftentimes becomes the cause of internal hernia; the traction on the borders of this opening being especially manifested in the direction of the attachment of the mesentery of the appendix, that is, along the brim of the pelvis towards the sigmoid flexure, leads to

fibroid thickening in that position, and as it becomes more and more drawn forward, it assumes the appearance of a band of inflammatory adhesion. The same traction may be the predisposing cause of ulceration of the appendix, and we have seen perforation and internal strangulation thus produced, associated with ulceration and perforation of the appendix. Immediately above the ileum, at its angle of union with the cæcum, we not unfrequently have a pouch, which, in atrophic changes of the peritoneum, becomes thinned, and perhaps perforated, thus forming another source of internal strangulation.

When the appendix cæci is fixed by abnormal adhesions, the rotatory movements of the cæcum to which we have referred must necessarily be modified. Again, as to the position of the cæcum, we may further remark, that another congenital peculiarity is sometimes found, namely, its presence in the left instead of the right iliac fossa; this we observed in a patient who died from chronic bronchitis and emphysema, and in whom the whole of the abdominal viscera were laterally transposed, the stomach, spleen, and descending colon occupying the right side, whilst the cæcum, the liver, and ascending colon were found on the left. The thoracic viscera were also transposed.

In a normal state of parts, the ileo-colic valve prevents the regurgitation of fluid from the large to the small intestine; the greater the distension of the cæcum, the more closely are the component parts of the valve compressed, and after death the colon may be fully distended, without escape of fluid into the ileum. Dr. Brinton and Mr. Roper have shown that if the ileum be also over-distended, the valve ceases to act; there is equal pressure on both sides, and the contents of the cavities may intermingle, or pass from the cæcum into the ileum.

The secretion of the cæcum is alkaline in its character. Tiedemann and Gmelin considered it acid, but in many cases that I have examined, it has been found alkaline. Chemical action probably takes place on particles of alimentary matter left unacted upon by the gastric juice, and by the secretions poured into the small intestine.

This action is very much less than that which takes place in the small intestine, and there does not appear to be sufficient warrant for the statement that the cæcum constitutes a second

stomach, and that true digestion here takes place. It is more probable that the watery parts of the chyme, if the semi-fæcal contents of the ileum may be so called, become absorbed by a very extensive capillary circulation; and that the glands remove from the blood excrementitious material no longer of any service to the system.

The appendix is an elongated gland of a very simple character, resembling the pancreatic cæca of the intestine of the fish, and, as far as is at present known, its secretion is of the character of ordinary mucus. Since the fæces here become more solid, were it not for such a secretion, assisted by that of the ordinary mucous follicles, adhesion would be more likely to take place with the parietes, and distension be the result. The secretion is poured out at that part which is most likely to effect this separation, namely, at the origin of the triple muscular band.

Pathology.—The unusual mobility of the cæcum, which has previously been referred to, is of a congenital character; but may induce serious pathological conditions, as before mentioned.

The villi cease at the ileo-colic valve, but we sometimes find in the cæcum and ascending colon elongated processes, resembling enormously hypertrophied villi scattered over the mucous membrane. In a case which recently occurred at Guy's, they were nearly half an inch in length, about one line in breadth, and covered the cæcum and ascending colon, but were not known to have produced any symptoms, nor to have had any influence on the cause of death.

Atrophy.—Since only a part of the cæcum is covered by the longitudinal bands, we frequently find that in atrophic states of the intestine the right side bulges out in a globular and almost hernial form. This condition is more common in advanced life, and in struma and phthisis; and we have observed the contraction of a cicatrix almost shut off this sacculated portion, so as to render the passage from the ileum to the ascending colon about the size of the ileum itself.

Distension.—Abnormal distension of the cæcum is sometimes the consequence of obstruction in the colon, or its own muscular parietes contract with less than their wonted vigour, and then it easily becomes distended by the accumulation of fæces or of flatus. It is probable that diminished secretion from the

appendix cæci may favour this accumulation of fæces, which is often amongst the exciting causes of serious disease, and requires attention. Considerable fæcal distension in the cæcum and ascending colon produces pain in the iliac region, and by pressure on the last dorsal and genito-crural nerve, induces pain also over the hip, as far as the great trochanter, the groin, and the testicle, &c.

The pain thus produced is sometimes of an acute character, resembling colic, and it may excite considerable alarm. Dr. Copland mentions œdema of the right leg as a result of distended cæcum; this I have not observed, except with very feeble power, or a varicose condition of the veins. Pressure of this kind would, doubtless, perpetuate and aggravate varicose condition of the veins of the lower extremity.

Many instances of pain in the region of the cæcum arise from distension, and the symptoms entirely disappear when the colon is gently but freely acted upon, and emptied.

Edema of the mucous membrane is often observed with renal anasarca, and with long-continued congestion of the vena portæ.

Congestion.—The depending position of the vessels often produces a passive fulness of the capillaries of this part of the alimentary canal; but we also find an active congestion, as shown by arborescent injection of the minute vessels. This is sometimes produced, probably, by medicine administered a short time before death; as the elaterium powder in renal, hepatic, or pulmonary ascites, &c., or it is the result of the transmission of irritating substances and secretions from the small intestine, as an excess of bile or excreta of an acrid character, undigested food, &c.

Inflammation.—The distension of the cæcum, to which we have previously referred, induces local enteritis, namely, inflammation of the mucous membrane of the cæcum, and of the peritoneum which invests the part. These constitute a numerous class of cases, which are happily more tractable than those in which peritonitis is set up by a concretion in the appendix cæci. The mucous membrane is congested, its secretion altered, the fæces adhere to it, the muscular coat is unable to propel the contents, which constitute a tumour felt on palpation, and the inflamed peritoneum produces tenderness. In most cases, this

tumour consists of portions of intestine united by inflammatory adhesions, and in still more rare instances it is composed of effusion behind the cæcum, in the iliac fossa.

The direct continuity of mucous membrane from the ileum appears in many cases to be the cause of disease in the cæcum; we find this in typhoid fever and in strumous disease, in which the ileo-colic valve is often acutely inflamed, swollen, injected, and ulcerated.

The cæcum is found acutely inflamed in some cases of dysentery; it becomes injected, the mucus scanty, fæces adherent, or the surface covered with a delicate false membrane. These states are frequently observed in dysentery, the cæcum being affected as a part of the colon from continuity of structure.

Gray Discoloration.—As the result of chronic disease, as of congestion and inflammation, we find gray discoloration, sometimes generally distributed, at other times constituting minute zones around the solitary glands; or there are small circular ulcers, which have originated in disease of the solitary glands, or mucous follicles, around which this gray deposition has taken place.

A *granular condition* of the mucous membrane, as if minutely studded with particles of sand, appears to be the result of long-continued slight inflammation, and is associated with thickening of the mucous and sub-mucous coats.

Ulceration of the cæcum is rare as a disease simply affecting this part, but when associated with other morbid states it is of frequent occurrence; thus in phthisis, tubercle is often observed in the mucous membrane, and, as a consequence, ulceration; so also in typhoid fever it is not unfrequent to find some scattered ulcers in the cæcum. In dysentery, also, the cæcum is not only acutely inflamed, as before mentioned, but it frequently presents extensive ulceration; sometimes a transverse ulcer is found to present ragged and injected margins; at other times the mucous membrane is even in a sloughing state, or suppuration takes place between the layers of the intestines, as was found in a remarkable degree in one of the cases of dysentery subsequently detailed. In the acute inflammation and ulceration of the colon following poisonous substances, as arsenious acid or corrosive sublimate, the rectum and sigmoid flexure are generally more severely affected than the cæcum and ascending

colon; this, however, is not invariably the case. Again, ulceration is often present from over-stretching, as we have frequently found in obstruction of the sigmoid flexure; the mucous membrane yields in transverse lines, as when the skin has been similarly over-stretched and ulcerates.

In severe instances of enteritis affecting this part of the intestine, or typhlo-enteritis as it is sometimes termed, the peritoneum becomes also involved; this may arise from the propagation of the disease from the mucous to the serous layer, or from rupture of the intestinal coats; but perforation into the peritoneal cavity is more frequently found to arise from disease of the appendix than from simple ulceration of the cæcum; extravasation and sudden fatal peritonitis, even when perforation has taken place, may, however, be prevented by adhesions; or if these adhesions be less extensive, faecal abscess takes place. We shall again have to refer more fully to perforation of the appendix cæci from ulceration variously produced. The peritonitis which ensues when no adhesions exist, is almost as sudden in its symptoms, and as fatal in its results, as perforation of the stomach; but in cases much more numerous than is generally supposed, extravasation is prevented by antecedent adhesions, as has been shown by M. Leudet, whose observations I can fully confirm. But perforation from ulceration may take place on the attached surface of the intestine; the disease then extends into the cellular tissue in the iliac fossa; pus burrows beneath the fascia, and forms an opening near the crest of the ilium, or below Poupart's ligament on the thigh. These cases are, however, rare; and ulceration of the appendix much more frequently leads to adhesions of the peritoneum, to faecal abscess, or to general peritonitis.

Cancerous disease not unfrequently attacks the cæcum; sometimes the ileo-cæcal valve presents an appearance similar to that found in like disease at the pylorus, and an extensive cancerous ulcer is found to extend upon the anterior or posterior surface of the cæcum; when the former course is followed, peritoneal adhesion or inflammation take place; and when the latter, the cellular tissue in the iliac fossa is involved, and suppuration sometimes occurs; in either case faecal abscess and artificial anus may be produced.

Constipation is often present in a greater or less degree, and

the obstruction is further shown by the hypertrophy of the muscular coat of the ileum; this is more especially the case when the disease is slow in its progress and of a scirrhus character. Sometimes, indeed, there is so much fibrous tissue deposited that it is difficult to find true cancerous product; and many instances are regarded as of simple inflammatory character. On the contrary, the disease is sometimes soft and fungating, and has the microscopical and general appearance of a medullary growth. Again, epithelioma and colloid cancer are occasionally observed. The disease has a greater tendency to extend along the coats of the cæcum itself than to pass backwards into the ileum; and it may be further remarked that the cæcum is less frequently affected with cancer than the sigmoid flexure and the rectum.

The *Tricocephalus dispar* is described as being frequently present in the cæcum. I have only observed them about three times from many hundreds of inspections, in very many of which the intestines were examined throughout with care.

Appendix—Increase of length.—The appendix is sometimes five or six inches in length, and perfectly free in its movements. It may be free among the coils of the small intestine, or in other cases becomes adherent to the brim of the pelvis, to the parietes of the abdomen, or to the mesentery. In this way loops are formed, which in many cases become the cause of fatal internal strangulation, a portion of small intestine passing beneath the band thus formed. Cases have been recorded of the appendix being found in a hernial sac.

Atrophy.—The orifice of the appendix is occasionally obliterated, and the appendix itself bound down by adhesions; in this way it becomes wasted, and at last almost destroyed.

Dilatation takes place from obstruction at or near the orifice, so that the secretion is unable to make its escape; the canal then dilates, and becomes one fourth to one half of an inch in diameter; the walls are sometimes thickened, and the muscular coat hypertrophied, as if the attempt had been made to overcome the obstruction, or, on the contrary, it becomes exceedingly thinned almost to perforation; when so dilated it is filled with thin mucus, and the follicles have the appearance of minute semi-transparent cysts.

Concretions.—Substances of diverse character are found lodged in the appendix cæci, and whilst sometimes harmless, often produce very serious consequences. Some are extraneous, others are entirely formed within the canal itself; and, lastly, there are those which have a nucleus consisting of some foreign substance, but covered over by layers of concretion, from the irritation produced.

1. Extraneous bodies are found, consisting of nails, pins, stones of fruit, shot, bristles of a tooth-brush,* entozoa, and most frequently, fæces.

2. Concretions, as of albuminous mucus, are formed in the appendix. These are not uncommon, and constitute firm semi-transparent masses, which, when dry, are fragile, and free from earthy matter.

3. Calculi, which generally present a nucleus of fæces, or of some foreign body.

I have frequently found the appendix filled with fæces; sometimes in its whole length, or only forming one or more hard nodules. These minute fæcal masses frequently constitute the nuclei of calculi, and become encrusted with layers, composed of carbonate and phosphate of lime, according to the analyses of Dr. Odling, and of Dr. Golding Bird. A concretion, examined by Dr. Prout, was found to consist of phosphate of lime, with a little carbonate, and contained a small quantity of animal and oleaginous matter.† Thus constituted, layer after layer becomes applied, till the size of a cherry stone is attained; and many of the so-called cherry stones in the appendix are thus constituted. The calcareous matters appear to be derived from the mucous membrane itself; in the same manner as a calculus in the urinary bladder becomes encrusted with phosphate of lime from the abundant mucus thrown out from the irritated surface. In some cases a larger size is attained, and the mass becomes as large as a date stone, or a hen's egg. In the museum of Guy's (No. 1893²⁵) is a large calculus the size of a hen's egg, the surface of which is rounded and fissured; it was removed from a sinus leading from the parietes of the abdomen to the cæcum; no appendix was found, but a large abscess extended

* 'Transactions of Pathological Society,' 1855. Mr. N. Ward's case.

† 'Medical Gazette,' vol. vi.

from the cæcum to the liver. The calculus was composed of phosphate of lime, with alkaline chlorides.

In some it is very difficult to discover a nucleus, a white laminated substance being presented throughout. The nucleus, however, may be exceedingly small, as in a case described by Mr. N. Ward, where the bristle of a tooth-brush formed the centre of a calculus; or it may be a portion of pin, or a hair.

Diminished contractile power of the muscular coat, with distension of the intestine and over-exertion, are the probable causes of the propulsion of fæces into the appendix; or it may be that the peristaltic contraction is rendered irregular by an irritated condition, from acrid and crude materials impelled into the cæcum, and that this irregularity of action causes the fæces, perhaps more fluid than normal, to pass into the appendix. In whatever way produced, any concretion in this part may lead to very serious results.

I. It excites irritation and ulceration of the mucous membrane, followed—

II. in most cases, by extension through the muscular coat; the ulceration may be limited to that part, or it may extend through the peritoneum.

III. The perforation sometimes sets up inflammatory action of a purely local character; effusion of fibrinous material takes place, adhesions form and prevent extension to the general surface of the peritoneum. Coils of the small intestine may be thus firmly united to the cæcum, and constitute a compact mass, felt on manual examination of the abdomen.

IV. The inflammatory action, although local, may produce a less organizable product; and suppuration may take place, constituting an abscess, into which a greater or less quantity of fæces may escape. The subsequent course of this abscess is very different:—1. It may constitute a dried mass of semi-calcareous product. 2. After sudden exertion the adhesions which localize the pus may break down, and extravasation take place into the general cavity of the peritoneum, with a speedily fatal result. 3rdly. It may pass into the intestine by a second opening, and thus be harmlessly discharged; this opening may be into the ascending colon or the ileum; or it burrows down into the pelvis; and in a specimen in the Guy's museum, an

elongated and ulcerated appendix had opened into the rectum. 4thly. The cellular tissue behind the cæcum may become involved, the abscess extending sometimes upward behind the ascending colon, or down towards Poupart's ligament; the latter opening either below that ligament, or near the anterior and superior spinous process of the ilium.

In a case under Dr. Barlow's care, in Guy's, this ulcerative extension of cæcal disease destroyed the parietes of the iliac artery, and led to almost immediate death, from the sudden and uncontrollable hæmorrhage.

Abscesses of this kind sometimes contain fæces; and we have then but little chance of preventing repeated attacks of inflammatory action, the strength at last giving way, or life is cut short by intense and general peritonitis.

V. The perforation sometimes takes place directly into the peritoneum, and sets up peritonitis so severe and general, that a fatal result follows in a few hours, or at most in a few days.

These are terrible cases, with scarcely any premonitory symptoms; the patient is struck down in fatal collapse, resembling the equally fatal cases of perforation of the stomach, where scarcely any indication has been given of such an attack. In fact, sometimes the pain is not situated in the region of the cæcum, but above, nearer the stomach; whilst I have seen a case where the pain preceding fatal collapse was in the region of the cæcum, when the perforation arose in the stomach. It is difficult to explain this occasional event, but generally speaking, the pain is situated in the neighbourhood of the diseased viscus. The position of the concretion, whether fæcal or otherwise, is various; sometimes quite at the termination of the appendix, at other times close to the opening into the cæcum. Fæcal masses are also found adherent beneath the ileo-colic valve, and in sacculated depressions on the surface of the cæcum; but I have never witnessed true concretions in these parts as in the appendix.

In strumous patients, these concretions more readily tend to an unfavorable result, leading to perforation, and to fæcal abscess or peritonitis; but the appendix is itself the seat of ulceration, without the irritation of concretions; and especially so in strumous subjects. In phthisis it is very common to find ulceration in the

appendix cæci, from the degeneration of tubercle and subsequent ulceration; sometimes several small ulcers are present, at other times the appendix is almost amputated. This condition may lead to fatal peritonitis in the earliest stage of phthisis; again, the low organized product of struma is found either as minute tubercles in the substance of the mucous membrane, or as a large cheesy mass, filling up the extremity of the appendix.

Symptoms.—The symptoms of some of these pathological conditions have been already alluded to; but in others no sign indicative of their presence exists during life. Death results from other causes; and conditions are found which would have acted as disturbing causes, predisposing to serious, if not to fatal disease, if life had been prolonged.

Distension of the cæcum is indicated by fulness and pain in the iliac region, especially when the erect posture is assumed, or after walking; it is generally accompanied by fulness in that part, dulness on percussion, and slight febrile excitement with congested portal circulation, and with loaded colon. Hence, we often find other symptoms present, not arising from the cæcum, but from associated disease; thus depression of mental energy, sallow complexion, furred tongue, offensive breath, pain in the head, arise not from the condition of the cæcum, but from the retention in the blood of waste material, which would be thrown off, if the liver and excretory glands of the whole alimentary canal rightly performed their function. The mechanical distension, however, sometimes by its pressure leads to pain in the loins, or in the course of the last dorsal or genito-crural nerve, the pain extending over the dorsum of the ilium, or into the groin or testicle; in women it interferes with the proper function of the ovaries and uterus.

Irritation or inflammation of the mucous membrane of the cæcum may be productive of diarrhœa, and generally there is but slight pain in the region of the cæcum. It is, however, in most cases, only part of a more general disease of the mucous membrane; as in bilious diarrhœa, from acrid excreta poured into it, in dysentery, and in struma; disease of the mucous membrane alone is not productive of pain. If *all* the coats be affected, or ulceration have taken place, a very marked train of symptoms follows. After some irregularity of the bowels, either

diarrhœa or constipation, generally the latter, and perhaps after more than wonted exertion, severe pain comes on, in many cases suddenly, in the right iliac fossa. The pain may be confined to this spot, and be accompanied by excessive tenderness, radiating over the abdomen, and be very quickly followed by collapse, and the signs of general peritonitis, namely, extremely anxious countenance, sunken eye, cold extremities, distended and tympanic abdomen, clammy partial sweats, failing pulse, and death in a few hours; or the tenderness and pain in the neighbourhood of the cæcum are accompanied with fulness and slight dulness on percussion. There are febrile symptoms, the skin is hot, the tongue is slightly furred, the pulse is often compressible and somewhat excited, local peritonitis is set up in connexion with ulceration or inflammation of the coats of the cæcum. These are the symptoms of what has been called tumphlo-enteritis. There is often a gradual subsidence of these symptoms, the pain and distress cease, the fulness disappears, the bowels return to healthy action, and the patient is restored to health. Or the fulness, tenderness, and pain continue, and a more defined tumour is perceptible, repeated attacks of severe pain come on, and gradual loss of strength, or sudden accession of fatal and general peritonitis. The local peritonitis has given rise to suppuration or to fæcal abscess; and perforation of this abscess is the cause of the sudden collapse and speedy death. Or, we may have the same result as before described, but retarded for a time by local adhesions. Or, instead of peritonitis, diarrhœa may be set up, irritability of stomach, injected and brown tongue, failing pulse, and the ordinary symptoms of hectic fever. But even from this condition recovery sometimes takes place, by the discharge of pus from the peritoneal abscess into the intestine itself, or through the abdominal parietes; or by the absorption of the fluid parts of the pus, when a semi-cretaceous mass is left; if, however, fæcal abscess have formed, recurrent attacks of peritonitis, with increasing prostration, generally lead to a fatal result.

In *cancerous* disease of the cæcum, the symptoms are very similar to those already described, namely, pain and fulness in the region of the cæcum, tenderness on pressure, and a more or less distinct tumour; there is diarrhœa or constipation, but

generally the latter; the cæcal pain is often greatly aggravated by food, especially of a fluid kind; and the accession of pain is sometimes found to arise directly after the nourishment has been taken, as in Case CLVIII.

There is a greater tendency to the local form of disease in early manhood than in later life. Many cases occur under 20, but the disease is not rare at later periods of life, 30, 40, or 50 years of age.

Diagnosis.—In the diagnosis of cæcal disease it must be borne in mind, that simple excessive distension of the cæcum is sometimes accompanied with severe pain.

2. That after blows on the abdominal parietes, or from other causes, suppuration sometimes takes place in the cellular tissue, or even in the muscles of the iliac fossa, and may be accompanied by local pain or peritonitis without cæcal disease.

3. That suppuration connected with the right kidney, or its envelopes, sometimes extends into the iliac fossa.

4. That we may have disease of the vertebra, or iliac bones, leading to suppuration in the iliac fossa.

5. Pain arises in the course of the last dorsal nerve from diseased spine, or in the course of the genito-crural nerve from renal calculus, and might be confounded with cæcal inflammation, &c.

6. Inflammatory disease in connexion with the ovaries, leading to local peritonitis and severe pain, is frequently mistaken for cæcal disease.

7. Cancerous disease of the cæcum; and

8. Disease of the ileum in struma or after typhoid fever, as well as,

9. Strumous peritonitis must each be remembered in forming a correct diagnosis.

10. I have observed the peritonitis from cæcal perforation regarded at first as gall-stone, the sudden pain on the right side, with violent vomiting, closely simulating the symptoms of that disease.

Dr. Battersby, in a very interesting paper in the 'Dublin Quarterly' of 1857, refers to other fallacies, as hernia, disease of the hip, and of the genital organs.

The pain in simple distension of the cæcum is less severe than

in acute inflammation. Disease in the parietes in a very short time manifests its local character; but at first the diagnosis is obscure, and mere faecal odour of pus does not necessarily imply communication with the intestine. The pain and swelling connected with suppuration of the spine or kidney differ in position; that of the kidney is more in the loins, or, if it extend anteriorly, is nearer to the median line. Spinal suppuration extends beneath the iliac fascia, and would be distinguished from cæcal disease burrowing beneath Poupart's ligament, by the faecal character of the discharge in the latter.

The neuralgic pains connected with urino-genital disease, are not accompanied with the tenderness or the other symptoms of intestinal affection. It is, however, sometimes difficult to distinguish inflammatory disease about the right ovary from cæcal disease. There may be in both excessive tenderness, febrile excitement, constipation, severe pain in the lower part of the iliac fossa. The symptoms which will serve to guide us are, that the ovarian disease comes on with irregular menstruation or with sudden cessation of that flux, and that the pain is situated lower down in the hypogastric region; in some cases, observers have believed that they have felt the swollen ovary. Dr. Barlow records a case in which peritonitis of so severe a character was set up around an inflamed ovary, that the patient succumbed. In cancerous disease of the cæcum, which sometimes occurs in young subjects, it is almost impossible, unless there be indication of cancerous disease in other parts, rightly to diagnose its character. These are, however, rare cases. In strumous peritonitis the disease is not confined to one part of the abdomen, but in severe cases the intestines are so completely united by peritoneal adhesions as to move *en masse*. It is impossible to distinguish perforation of the ileum in struma or phthisis from perforation of the appendix cæci; this is, however, of little moment, since the only remedial agents which are likely to be of service in these almost universally fatal cases are precisely similar in both.

Prognosis.—In cases of cæcal distension, when the mucous membrane only is affected without ulceration, our prognosis is generally a favorable one, unless we find the patient of a strumous habit, in whom there is greater tendency to ulceration and perforation. When, again, there are the symptoms of local

peritonitis, many patients do well; the reverse, however, is the case when the onset of the disease is marked by severe collapse, or by urgent vomiting and by general abdominal pain.

Causes.—The *predisposing causes* are strumous diathesis, sedentary habits, habitual constipation, typhoid fever, &c.

The *exciting causes* are, over exercise, much standing, violent athletic exercises; in many cases cæcal disease has come on after very long pedestrian excursions, after indigestible food had been taken, after blows upon the abdomen, after constipation, or an irregular condition of the bowels.

Treatment.—I cannot urge in too strong language the importance of avoiding in cæcal disease powerful drastic purgatives. They tend to increase the disease by inducing violent peristaltic action; by increasing the irritation of an already inflamed membrane they hasten ulceration; and if ulceration have taken place, or if peritonitis have resulted, the only hope of the patient is taken away.

If there be simple distension, with only very slight pain in the erect posture, we should enjoin rest, and administer grey powder with Dover's powder, followed by a dose of castor-oil, or by a castor-oil enema; afterwards mild aperient tonics, as the compound gentian mixture, are useful.

If tenderness exist, or there be the symptoms of local peritonitis, rest is still more positively required; the patient should not move from the bed on any consideration. Local depletion is exceedingly valuable; ten or fifteen leeches applied to the region of the cæcum, and warm fomentations, are often followed by most marked benefit. At the same time mild mercurials may be administered, with opium, or equal parts of grey powder and Dover's powder, or calomel with opium; but I prefer opium or Dover's powder without the mercury. These remedies may be combined with saline medicines, with the acetate of ammonia and bicarbonate of potash or nitric ether, according to circumstances; but little food should be taken, and only of bland unstimulating kind. It is a great temptation, when the pain has subsided, and the febrile excitement has disappeared, for the patient to try and get out of bed and use slight muscular effort; but this is exceedingly injudicious, and is sometimes followed by fatal result.

The remedies just mentioned often induce action on the bowels; but if not, although the pain have subsided, it is better to wait, or to administer a gruel or castor-oil injection, than to give more powerful purgatives, as aloes, jalap, senna, colocynth, scammony, &c.

If there be persistence of slight pain, with fulness and dullness, it is well to continue the opium, and either to repeat the leeches, or to apply a blister to the iliac region.

Afterwards iodide of potassium, with mild vegetable tonics, are of great service, still maintaining rest. Irritability of stomach sometimes arises, which may be alleviated by saline effervescing medicine, by hydrocyanic acid, by soda-water with milk, or with brandy, &c.

If there be evidence of suppuration or of fæcal abscess, whilst we endeavour to limit the action by slight counter-irritants, or by occasional local depletion, we must sustain the power of the patient by a generous diet, by quinine, and by tonic treatment. Opium is often of great value in its anodyne and narcotic action, in checking peristaltic action, in relieving pain, in soothing an over-excited nervous system, in diminishing the excitement of exhaustion, and often in procuring refreshing sleep.

When there is collapse and tympanitis, evincing perforation of the appendix or intestine, nothing should induce us to administer any aperient, or to urge an action from the bowels. We desire to limit the mischief produced by checking the movement of the intestines, and to diminish inflammatory action by soothing the nervous system; opium must be given very freely, and only a very small quantity of food administered.

Cases CXXVII to CXXXI are instances of abnormal position of the cæcum, and several were connected with fatal obstruction; a similar instance of fatal obstruction is recorded by Mr. Avery, in the 'Pathological Transactions' for 1850, where the operation for artificial anus was performed.

CASE CXXVII.—*Unusually Free Cæcum*.—John J—, æt. 5, on November 1st, 1856, whilst running down stairs, fell upon an earthen vessel that he was carrying; the internal jugular vein was divided, and he died about an hour afterwards from the loss of blood.

The viscera were found to be healthy, but in the abdomen the cæcum was situated among the small intestines, quite surrounded by peritoneum, and as free as a portion of the ileum.

CASE CXXVIII.—*Unusually Free Cæcum. Fatty Degeneration of Organs.*—A young woman who had lost a considerable quantity of blood after miscarriage, gradually sank in six weeks. On inspection, the heart, liver, kidneys, &c., were found to have undergone fatty degeneration. The uterus contained a sloughy lining membrane, and apparently some decidua.

The *cæcum* was attached by a long mesentery to the right side of the spinal column, so that the whole iliac fossa was perfectly free, and covered by peritoneum.

These conditions in the last two cases were congenital, and had any subsequent disease been set up in the part, the symptoms might have become considerably modified.

CASE CXXIX.—*Cæcum inverted. Phthisis. Local Empyema. Large White Kidneys.*—Alfred A—, æt. 42, was admitted into Guy's Hospital November, 1856, and died December 14th.

He suffered from phthisis. At the apex of the left lung was a small vomica with tubercles and iron-gray consolidation around it. In the lower lobe was an irregular vomica, and a sloughing mass of lung tissue, which communicated with local empyema at the base of the lung.

The heart and liver were healthy; the spleen was lardaceous; the kidneys were large, white, and mottled.

On examining the intestine, the appendix cæci was found to be long, and extending over the brim of the pelvis, where it was fixed. The rounded termination of the *cæcum* was directed towards the diaphragm as if inverted. The ascending colon was contracted, and attached deeply at the side of the right iliac fossa, directly opposite the ileo-colic valve, and at an acute angle with the *cæcum* (see Plate IV, fig. 1). Very great distension of the *cæcum* in this twisted state might lead to obstruction, for the ascending colon appeared, even in this case, constricted by the sudden twist and by the acute angle which was formed.

No symptom had apparently been produced by this condition of the *cæcum*; but in a state of constipation, when the *cæcum* is distended with *fæces*, considerable impediment to the free passage would be the result. It is probable there would be a greater tendency to ulceration, and to the passage of *fæces* into the appendix cæci.

CASE CXXX.—*Intestinal Obstruction of the Ascending Colon. The Cæcum twisted to the left side into the Left Iliac and Hypochondriac Regions. Death on the 20th day.* (Reported by Mr. Galton).—Eliza S—, æt. 40, a cook, was admitted into Guy's Hospital under the care of Dr. Addison, November 9th, 1856. She had lived regularly and temperately. There were marks of distension upon the abdomen, but she stated that she had never been pregnant,

20th.—Flatus was again passed.

22nd.—There was thick stercoraceous vomiting; the tongue was dry, the pulse 86. The seat of pain was between the umbilicus and the scrobiculus cordis.

23rd.—An injection of beef tea and eggs was retained for a short time, and was returned with small lumps of fæces.

The fæcal vomiting continued, the pain became more severe and general, with gradually increasing prostration, and she died on the 27th, about twenty days after the commencement of the symptoms.

On opening the abdomen the small intestine was found enormously distended, and the cæcum was situated in the left hypochondriac region and iliac fossa, forming a large, enormously distended, almost spherical sac; the appendix was situated on the left side. The whole of the visceral and of the parietal peritoneum was intensely injected, and was covered with lymph. The right iliac fossa was filled by coils of small intestine, the peritoneum being perfectly smooth. By attempting to unravel the intestine, and tracing the large intestine upward from the sigmoid flexure which was normally situated and perfectly collapsed, a stricture was found about the middle of the ascending colon; the stricture, however, was situated near the brim of the pelvis, on the *left* side, and adhesion of the omentum was found at this part, between the ascending colon, sigmoid flexure, and a coil of ileum.

The line of obstruction was perfectly defined; all the intestine below being quite empty, collapsed, and non-injected. The obstruction was four feet four inches from the anus, and appeared to have been produced by the cæcum revolving on the termination of the ileum, which was fixed by its adhesion to the sigmoid flexure. No transverse colon could be found, because it was hidden behind the cæcum near the left iliac fossa. (See Plate IV. fig. 2.) On removing the intestine the stricture disappeared.

The mucous membrane of the cæcum was intensely injected, and a patch on the interior surface was of a leaden colour; at the centre of this part was a minute slough, and perforation extended into the peritoneal cavity; but no fæcal extravasation had taken place.

The mucous membrane of the ileum was healthy, but congested, and contained both solid and fluid fæces. The stomach contained fluid fæcal matter, such as was found in the cæcum. The duodenum was healthy, and the liver, kidneys, spleen, &c. were normal.

The previous attack of inflammation of the bowels had probably led to the adhesion between the termination of the ileum and the sigmoid flexure; and this was one of the causes of the fatal obstruction. The cæcum was apparently unnaturally free, and its distension associated with this adhesion had led to the twisted and inverted position which was found after death. The pain had commenced at the seat of the disease, near the right iliac fossa, extending to the umbilicus. She had had severe

but that when a child her abdomen had been much enlarged. Her father died from phthisis. Her mother from dropsy. She was a seven months' child, but enjoyed good health until she was fourteen years of age, when she fell against the kerbstone at the head of a well, whilst she was drawing water; pain in the loins, with difficulty in micturition and hæmaturia came on. Catamenia afterwards appeared, but were scanty. Many years before admission she had jaundice, with great pain in the stomach, and was told she had inflammation of the bowels; she, however, had good health until 1846, when, during frosty weather, she fell down in a yard, striking her left side against the corner of a stool; she suffered from pain and tenderness at the part, with cold chills; the urine was scanty, but no blood was passed; after remaining in bed for three or four days she felt no further inconvenience. In 1850 she had vertigo, pains in the head and in the bowels, with diarrhoea. The bowels had been frequently confined for three or four days together, but without known inconvenience. On admission into Guy's she was anæmic, but her complexion was rather dark; three days previously, without apparent cause, pains came on in the right side, extending to the umbilicus. No improper food had been taken, nor was there any stomach derangement. She felt chilly; the bowels were opened, and again very slightly, at the time of admission. There had been no vomiting till a short time previously, after this symptom had supervened everything was rejected. There was no tenderness of the abdomen, but it was distended and flatulent. The skin was cool and moist, the urine abundant, pulse 80, the tongue slightly furred. A soap injection was administered, and ʒss of castor-oil given.

November 11th.—A small motion was passed, but no flatus, and she was unable to take nourishment. The abdomen was more distended, and there was tenderness in the right hypochondriac region; the pain was paroxysmal, with a sense of twisting. Magnesia mixture with vinum opii was given every six hours.

12th.—A portion of distended intestine could be seen at the umbilicus; in other respects she remained as before. Calomel gr. v, opium gr. i, were administered; and Oij of warm water were injected into the rectum, and a hot poultice applied to the abdomen.

13th.—All the symptoms were aggravated; the vomiting was less, but the paroxysms of pain were very severe. Since the bowels had acted slightly after admission, it was thought advisable to give purgatives a trial, and colocynth and calomel, gr. v, were given every hour for four times, and the enema repeated.

14th.—The injection was returned, but without fæces; the vomiting became stercoraceous, and the pain very severe. Opium, gr. i, was then given every six hours.

15th.—There was less pain and distension of the abdomen, the pulse full, 92, the skin dry, the urine moderate in quantity.

17th.—There was extreme tenderness of the abdomen, and the stercoraceous vomiting continued as before.

18th.—An ivory tube was introduced into the rectum and retained for a short time; its withdrawal was followed by flatus and small fæcal discharge, and afforded considerable relief.

20th.—Flatus was again passed.

22nd.—There was thick stercoraceous vomiting; the tongue was dry, the pulse 86. The seat of pain was between the umbilicus and the scrobiculus cordis.

23rd.—An injection of beef tea and eggs was retained for a short time, and was returned with small lumps of fæces.

The fæcal vomiting continued, the pain became more severe and general, with gradually increasing prostration, and she died on the 27th, about twenty days after the commencement of the symptoms.

On opening the abdomen the small intestine was found enormously distended, and the cæcum was situated in the left hypochondriac region and iliac fossa, forming a large, enormously distended, almost spherical sac; the appendix was situated on the left side. The whole of the visceral and of the parietal peritoneum was intensely injected, and was covered with lymph. The right iliac fossa was filled by coils of small intestine, the peritoneum being perfectly smooth. By attempting to unravel the intestine, and tracing the large intestine upward from the sigmoid flexure which was normally situated and perfectly collapsed, a stricture was found about the middle of the ascending colon; the stricture, however, was situated near the brim of the pelvis, on the *left* side, and adhesion of the omentum was found at this part, between the ascending colon, sigmoid flexure, and a coil of ileum.

The line of obstruction was perfectly defined; all the intestine below being quite empty, collapsed, and non-injected. The obstruction was four feet four inches from the anus, and appeared to have been produced by the cæcum revolving on the termination of the ileum, which was fixed by its adhesion to the sigmoid flexure. No transverse colon could be found, because it was hidden behind the cæcum near the left iliac fossa. (See Plate IV. fig. 2.) On removing the intestine the stricture disappeared.

The mucous membrane of the cæcum was intensely injected, and a patch on the interior surface was of a leaden colour; at the centre of this part was a minute slough, and perforation extended into the peritoneal cavity; but no fæcal extravasation had taken place.

The mucous membrane of the ileum was healthy, but congested, and contained both solid and fluid fæces. The stomach contained fluid fæcal matter, such as was found in the cæcum. The duodenum was healthy, and the liver, kidneys, spleen, &c. were normal.

The previous attack of inflammation of the bowels had probably led to the adhesion between the termination of the ileum and the sigmoid flexure; and this was one of the causes of the fatal obstruction. The cæcum was apparently unnaturally free, and its distension associated with this adhesion had led to the twisted and inverted position which was found after death. The pain had commenced at the seat of the disease, near the right iliac fossa, extending to the umbilicus. She had had severe

falls and blows upon the abdomen, one in particular, in which she struck the right side, and which, perhaps, tended to produce displacement or inflammatory mischief. The bowels had generally been confined, but she had occasionally suffered from diarrhœa. The first symptom was pain in the right iliac fossa, and then constipation; the severe colic, distension, tenderness, and vomiting were later symptoms. The mode of commencement appeared to indicate that it did not arise from simple impacted fæces. For four days there had been no vomiting, which showed the absence of internal hernia, of sudden strangulation, and of intussusception. The symptoms were neither those of enteritis, nor of acute peritonitis.

It was evident that there had been some chronic changes in the intestines or peritoneum, and it was difficult to decide the character of those changes.

No tumour could be felt; but there were three causes of obstruction left, between which it was exceedingly difficult to decide. 1. A slow growth connected with the intestine itself, as chronic contraction or cancer. 2. Old bands of adhesion; and, 3. Twisted intestine. An approximate opinion was formed as to the seat of the obstruction; either that it was at the colon, or at the termination of the ileum. The vomited matters were so fæcal in their character that it was even suggested that the tranverse colon might have formed a communication with the stomach.

The following case is recorded by Dr. Gull in the Guy's Reports of 1858, p. 179:

"CASE CXXXI.—*Displaced Cæcum. Perforation. Peritonitis. Paraplegia.*
—William J., æt. 28, of middle stature; fair hair; emaciated; anxious expression; large head; broad and prominent forehead. Though he had never been robust, he had good health until the beginning of the year 1857. He was first seized with vomiting, which came on without any discoverable cause, and lasted for several days. As he recovered from the attack the legs became weak. After three months he had a second attack of vomiting, followed by an increase of weakness in the legs. He was admitted into Guy's Hospital under the care of Dr. Gull, November 11th, 1857. He was then unable to stand without support. In a recumbent position he could flex and extend the legs with some freedom, but the movements were sudden and vague, from want of control over the action of the muscles; the spinal centres, when stimulated by the will, seeming to shoot off their influence at

once, making the feeble muscles contract to their full extent with a jerk. In other words, there was no power to regulate the muscular contraction. The movements of the fingers were also wanting in precision. He was awkward in handling small objects, or in applying the hand to grasp larger ones. The muscles were thin and flaccid, corresponding to the general emaciation of the body. The muscular irritability was excessive. Weak currents of electricity, and sufficient to affect healthy muscles, excited well-marked contraction; whilst a little stronger, but yet very moderate dose of electro-magnetism, produced cramps lasting for several seconds after the stimulus was discontinued. The arms were weak, with an obvious want of control over the voluntary movements. There was numbness of the feet and hands, and a burning formication in the fingers and toes. The sensation of the other parts of the body was normal. No involuntary contraction of the legs. Sphincters good. Urine acid. The lower ribs depressed, and but little moved in inspiration. Headache; vertigo; cerebral confusion; tinnitus. Pupils largely dilated, the left the most so; sight dim; occasionally transient amaurosis. Sleep disturbed by dreams. Frequent nausea and vomiting, with pain from the epigastrium to the spine. Abdomen not distended; soft. Skin hot and perspiring. Pulse permanently quick, 126, small, and feeble; respiration, 32; spine straight. No tenderness on pressure or percussion. He could give no account of any accident or injury to the spine, except such as might have resulted from a fall, flat on the back, from the height of a few feet, eight years before his symptoms began. His habits had been temperate; no syphilitic taint. After admission into the hospital he continued to have repeated attacks of vomiting, lasting for many days, uninfluenced by any remedies. The vomited matters were copious, greenish, and mucous. The bowels continued to act freely, but without relief to the sickness. The irritability of the stomach was attributed to the state of the cord. The attacks of vomiting increased his anxious aspect. The paraplegic symptoms continued unchanged. There were, as before noted, headache and vertigo, and sometimes transient amaurosis. The pupils remained permanently dilated, and with the same inequality. The pulse quick (120 to 130), and feeble. On only one occasion was the urine noticed to be alkaline, when first passed. It never contained mucus. There was no band-like sensation around the abdomen. He often complained of pain from the epigastrium through to the spine. About the middle of February, 1858, he first had a sense of bearing down about the rectum, and complained of great distress after an action of the bowels, and of startings in the legs. March 8th, he had an attack of vomiting, apparently such as had often occurred before. This continued on the 9th. On the 10th he was collapsed and pulseless, with cold sweats, and other symptoms of ruptured intestine. There was no cerebral oppression. He died on the 11th."

Inspection, March 13th.—There was no evidence of disease in the lungs, heart, and brain. The spinal membranes were healthy. Dr. Gull states, that "after hardening the cord and making fine sections, it was seen that the posterior columns were atrophied throughout their whole length, and amongst the tissue were numerous exudation cells in a state of fatty degeneration (granule cells)." *Abdomen.*—There was general peritoneal

inflammation, and extravasation of *faeces*. The cæcum was in the pelvis, enormously distended, and ruptured, presenting a large opening, two inches long, and the mucous membrane intensely injected and sloughing. The ascending colon passed up to the spleen on the left side, then after a very sharp angle the colon descended on the same side, to the sigmoid flexure. The ascending and descending colon were adherent, and empty. The cæcum appeared to have been twisted on itself.

During life it was difficult to account for the vomiting from which this patient suffered; Dr. Gull has referred "the early attacks of vomiting to the lesion of the cord itself as the primary disturbance, especially since the degeneration of the columns extended up to the neighbourhood of the medulla oblongata." Such may have been the case; but, in very many instances of spinal disease, no such irritability of stomach has existed; and we have evidence of intestinal obstruction in the position of the cæcum and ascending colon, and this obstruction was increased by the condition of the spinal nerves. Constipation is a common sign of spinal paralysis, and in this case so far increased the obstruction as rapidly to hasten the fatal termination.

CASE CXXXII.—*Twisted Cæcum. Obstruction. Peritonitis.*—David O'C—, *æt.* 30, was admitted into Guy's Hospital October 11th, 1859. He was a strong muscular man, and for three weeks his bowels had acted irregularly, but their precise mode of action could not be ascertained. On October 9th he partook freely of pork, ale, &c., and was soon afterwards seized with intense pain in the abdomen, which "doubled him up;" vomiting soon afterwards came on; these symptoms continued, and excessive prostration followed. The bowels had acted on Friday the 7th, but not afterwards till the day of his death. At the time of his admission into Guy's, on the evening of the 11th, he was cold, almost pulseless, perfectly sensible, but collapsed; he suffered intense pain in the abdomen, and distension; the abdomen was tympanitic, and presented distension in the epigastric region. Opium was given; an enema tube was introduced into the rectum, but only passed about six inches, and returned smeared with blood; a catheter drew off no urine from the bladder. Mr. Stocker thought, that the introduction of an acupuncture needle into the distended part might relieve the enormous distension; several punctures were made, and not affording relief, a minute trochar was introduced, much flatus passed, and a small quantity of thin *faeces*; but this discharge afforded the patient much relief.

Tincture of opium $\text{m} \times \text{l}$, chloroform $\text{m} \times$, were given in camphor mixture. On the following morning he still lived, cold, almost pulseless, legs drawn up; abdomen less distended, but very painful; he had passed water during the night; he stated that he had passed flatus several times from the bowels since the

commencement of the severe symptoms. About 11 o'clock there was an evacuation from the bowels, but the state of collapse continued, and he died about 3 p.m.

Inspection on the 13th, nearly twenty-four hours after death.—Decomposition was commencing in the abdomen, which was still much distended. The thoracic viscera were quite healthy. On opening the peritoneal sac the appendix cæci was observed about the centre of the abdomen, and a very large, distended cæcum occupied the greater part of the epigastric, umbilical, and *left* hypochondriac regions; a distended coil of ileum and a portion of ascending colon were attached to its inferior part; these portions were all of a green colour, and evidently strangulated. The rest of the serous membrane was much injected, reddish, covered with a thin layer of adhesive lymph, and smeared with red-coloured, opaque serum. The small intestines were distended. On passing the hand beneath the strangulated mass the constriction was found to be situated near the *right* iliac fossa, and was produced by the cæcum having rotated on a free mesentery, and then twisted on its axis. By turning the cæcum downwards all constriction disappeared. The line of strangulation was very distinctly marked across the colon, about four inches above the cæcum, and rather less distinctly across the ileum, about fifteen inches from the valve. The mucous membrane of the whole of this part was intensely injected, in some parts presenting ecchymosis; it very readily separated, and was covered with a thin, gray, diphtheritic layer; there was effusion of blood also into the constricted mesentery. The intestine in this part and above it contained fluid fæces, but some half-digested masses of food were also found in the cæcum. The stomach was healthy, as also the rest of the small intestine; the stomach contained green fluid almost of the character of that in the small intestine. The descending colon was contracted, and in the rectum was found a small ulcer immediately above the sphincter. The course of the intestine was very peculiar. The sigmoid flexure was situated in the *right* iliac fossa, the left being perfectly smooth and free from intestine; the descending colon crossed the spine in the centre of the lumbar region, from the inferior part of the spleen; the transverse colon was in its ordinary position, but had the enlarged and twisted cæcum concealing it; the descending colon had a curved position in the right loin, passing from the cæcum, which was situated immediately in front of the spine; and at the lower part of its curve old adhesions existed between the ascending colon and the sigmoid flexure, situated in the right iliac fossa. The left kidney was situated at the brim of the pelvis, and received its arterial supply from the aorta and the common iliac arteries; the ureter left the kidney anteriorly. The left supra-renal capsule was in its normal position. The liver, spleen, kidneys, &c., were healthy.

The cæcum in this case was preternaturally free, and the sigmoid plexure in the right instead of the left iliac fossa. It is probable that several weeks before death the cæcum had a twisted position, and that after an indigestible meal flatulent distension increased the twist, and rendered the obstruction complete. The

cæcum had apparently turned on its mesentery, as an axis, and then on its own axis. The adhesion between the ascending colon and sigmoid plexure was old, and possibly formed during foetal life. The position of the kidney, partly in the pelvis, and its unusual arterial supply, were evidence of congenital malformation. The introduction of the needle afforded no relief, but the trochar gave immediate cessation of pain, and the patient begged to have the operation repeated. The instrument had perforated the distended cæcum, and had diminished the distension of the strangulated part; it was left in the wound, so that no extravasation took place, and it is probable that this temporary relief to the distended bowel prolonged the patient's life several hours; not that we are prepared to recommend the repetition of such an operation.

The following are instances of a form of cæcal disease very frequently met with; they arise from distension of the cæcum, which induces local enteritis, with partial peritonitis; the latter varies greatly in intensity, being sometimes severe, at other times scarcely observable. They are related as briefly as possible. With proper care and judicious treatment, most of these instances recover. The symptoms are less severe than those in which the appendix is ulcerated, or contains a concretion; they come on more gradually, the pain is less intense, the dulness and tenderness are entirely removed as the inflammation subsides, and the bowel is freed from its contents. As in cases of more general enteritis, purgatives do considerable harm; they fail to empty the distended bowel, increase the enteritis, lead to ulceration, and in some to perforation and fatal peritonitis. The benefit arising from the action of the opium is very marked—the bowels act, the pain subsides, and the dulness lessens; mild mercurials with the opium are sometimes used, but we now prefer opium alone; abstinence from solid food and absolute rest are very important, and should be continued for several days after the subsidence of the pain. If the pain be severe, local depletion by leeches afford considerable relief, and should be followed by hot fomentations.

CASE CXXXIII.—*Cæcal Distension and Inflammation*.—Crota W—, a strumous-looking boy, an apprentice to a cook at a large tavern, after harder work than usual, was seized with severe pain in the abdomen on the

right side; after a few hours this partially subsided, but again returned on his making exertion, so that he was obliged altogether to discontinue his work. The bowels were occasionally constipated.

He was of fair complexion, with long eyelashes, and his countenance had an anxious expression; the abdomen was hot, tender and full, especially in the region of the cæcum; the tongue was red, the pulse soft, the thoracic viscera normal; he had no vomiting.

Calomel gr. j, opium gr. ss, were given every six hours, and a hot poultice applied to the abdomen. Eight leeches were afterwards applied, and spare diet allowed.

He rapidly improved, and in a few days was convalescent. He was kept in bed, however, for a longer period, although all the symptoms had subsided.

The marked strumous character of this boy rendered the disease less likely to assume a favorable character; there is greater disposition to disease of the appendix in such diathesis, and few things are more painful than a fatal relapse after partial convalescence.

CASE CXXXIV.—*Cecitis*.—Benjamin B—, æt. 15, a pale, thin lad, who had been employed on the river, was admitted into Guy's Hospital, January 14th, 1852.

About three days before admission he had experienced griping pain in the abdomen, which had increased in severity. The bowels were constipated, but there was no vomiting, nor could it be ascertained that he had partaken of improper diet. There was fulness in the right iliac region, with dulness and considerable tenderness. Eight leeches were applied, and of calomel and opium gr. j each every four hours. On the 19th the pain had considerably diminished, but still much fulness and hardness remained; there was no febrile disturbance; the tongue was clean and the pulse natural. The calomel and opium were omitted.

February 3rd.—He felt much relieved, but had a haggard look; the eyes were sunken, and occasional pain came on across the abdomen. There was no marked indication of progressive disease; and the fulness in the iliac region gradually disappeared.

23rd.—There was again very perceptible fulness and some tumefaction in the right iliac region, and gurgling on pressure; slight pain had returned, the symptoms were, however, very much less severe than before, and he was allowed to move about the ward. He afterwards left the hospital convalescent.

The symptoms in this case were at first very severe, and warranted a very cautious prognosis. They were probably associated with strumous diathesis, and more than usual disturbance

of the other abdominal viscera. There is much fear that slow strumous disorganization would extend in this case, and ultimately lead to a fatal result.

The permission to sit up led probably to the increase of the symptoms, but happily the relapse was not of a character to prevent his convalescence. This was an instance in which great care, nourishing diet, change of air, might be followed by complete restoration to health.

CASE CXXXV.—*Cæcal Distension and Inflammation.*—Charles W—, æt. 19, was admitted into Guy's Hospital March 15th, 1848. He was a confectioner, and three months previously had had a similar attack to that for which he was admitted. The bowels were generally constipated.

Five days before admission sudden pain came on in the umbilical region, which continued for one day and then passed to the right side; the pain was severe, but sometimes increased in severity; he took senna and salts twice, which acted freely, but with only partial relief.

On admission the countenance was expressive of much distress; the pain was in the right iliac fossa; the tongue was white, the bowels were confined, and the pulse was irritable.

Dover's powder and grey powder in gr. iij doses were given every four hours, and on the following day a small dose of castor oil.

March 17th.—The pain was much less severe, but the skin was hot; the sleep was disturbed, the bowels were open, the tongue was injected at the edges; the pulse was more full and softer, 92.

A blister was applied over the cæcum, and the medicine was continued. The cæcal symptoms were much relieved, but on the 24th he had swelling, pain, and redness of the right wrist-joint, resembling rheumatism; the skin was perspiring, and the bowels constipated.

Dover's powder was given night and morning, and castor oil in small doses.

He slowly improved, and left the hospital convalescent on April 11th.

This case was increased in severity by the purgatives given before admission and by the absence of proper rest. It appeared that he had had a similar attack previously. It was not thought advisable to repeat leeches, but the application of a blister was followed by considerable relief. It may be doubted whether the pain in the right wrist was really rheumatism; pyæmia is found in connexion with some cases of cæcal suppuration, but we had no proof of suppuration here, nor did the mild character of the complaint bear resemblance to pyæmia; however, it considerably retarded his convalescence.

CASE CXXXVI.—*Cæcal Distension and Inflammation*.—Jane L—, æt. 24, a single woman, of fair complexion, engaged in domestic service. Three weeks before admission she experienced severe pain in the region of the cæcum, and castor oil was administered in several ʒss doses, but without effect. On admission there was severe tenderness at the right iliac fossa, and other symptoms of cæcal disease. Leeches were applied, grey powder and Dover's powder given, with saline mixture.

The symptoms gradually disappeared, but diarrhœa came on; the pain and fulness subsided, and she steadily convalesced.

In this case constipation preceded the attack, but the purgatives produced no beneficial effect; the castor oil increased the pain, which only subsided on the application of leeches and the administration of opium, &c., with rest. The affection was evidently of an enteritic character.

CASE CXXXVII.—*Cæcal Distension and Inflammation*.—William W—, æt. 43, a tailor, and of intemperate habits, was admitted into Guy's Hospital December 24th, 1856.

For two months he had suffered from pain in the region of the cæcum, and the bowels had been kept open by the action of medicines. Two years before, the bowels had been very irregular. After considerable exertion in shooting, the pain gradually increased; coughing produced a stitch in the side; at last, whilst pulling on his boot, the pain became so severe that he could not rise.

On admission there was fulness and tenderness in the region of the cæcum; the pain was increased by respiration; there was slight dulness on percussion, and the bowels were confined.

Ten leeches were applied to the region of the cæcum; an injection of gruel and salt was administered; and grey powder, with Dover's powder, of each two grains, were given night and morning. Low diet only was allowed, and rest enjoined. The pain and tenderness disappeared after the leeches had been applied. The bowels acted, but remained more or less confined.

On January 1st he was free from pain, and had considerably improved; and on the 12th he was considered well, all the fulness and tenderness in the region having entirely disappeared.

CASE CXXXVIII.—*Cæcal Distension with Inflammation*.—George W—, æt. 31, was admitted into Guy's Hospital August 31st, 1850. He was a man of light complexion, who had worked at a candle manufactory. For three months he had been an out-patient at the London Hospital, complaining of pain in the region of the cæcum when he moved about, but he was relieved by rest.

He gave up his work nine days before admission, suffering as before; the bowels were not constipated; the tongue was clean; the pulse was compressible.

On examination, immediately above the iliac fossa was a small rounded mass, tender on pressure. Rest in bed was enjoined, and calomel with opium prescribed.

The patient left the hospital relieved in one month. The case is interesting, as showing cæcal disease in its mildest form; the pain absent during rest, but preventing the patient from taking exercise. The neglect of such a condition, or its frequent recurrence, would probably cause ulceration of the cæcum or of the appendix, and more serious peritonitis. After the subsidence of the more active symptoms, preparations of steel, and vegetable aperients, as rhubarb, ipecacuanha, with aloes and soap, or the compound gentian mixture, &c., are very likely means of preventing their recurrence. Violent muscular exercise should be carefully avoided.

CASE CXXXIX.—*Cæcal Inflammation simulating Hip-joint Disease*.—James C—, æt. 11, living at Gravesend, was admitted into Guy's Hospital, under my care, February 18th, 1857. He was a strumous child, but he was stated to have had good health till three months before admission. He was roughly used and beaten whilst at work; and he did not feel well afterwards; pain came on in the abdomen; but it did not become severe till a short time before admission, when his foot slipped, and his abdominal muscles were brought into powerful action. Severe pain in the region of the cæcum then came on, and was much aggravated by pressure; the rest of the abdomen was soft; the tongue was normal; pulse 75. The bowels were confined, and the urine was normal. The right leg was flexed at the thigh, and could not be straightened; rotation of the hip, striking the heel, &c., did not produce pain, nor was there any pain in the knee or in the spine. Seven leeches were applied to the right iliac fossa; grey powder gr. ij, Dover's powder gr. iij, three times a day, with rest and low diet, were ordered.

The leeches and hot poultice afforded much relief; he was able then partially to straighten the hip, which had evidently been drawn up to relieve the pain, by relaxing the flexor muscles of the hip. The bowels on the second day acted by soap injection, and on the third day the leg was straight. The pain and fulness gradually ceased; he was, however, kept in bed; the medicine was continued once a day for a short time, and animal food was allowed very sparingly. The bowels acted without trouble. On March 2nd he was convalescent; cod-liver oil was given three times a day, and on the 13th he left the hospital well.

CASE CXL.—*Local Peritonitis. Cæcitis (P).*—Martha P—, æt. 19, admitted into Guy's Hospital October 3rd, 1855, was a tall girl, who had been a servant, and for some time had not enjoyed good health. She had been lately in St. Thomas's Hospital with pain in the side, &c. She had menstruated a fortnight before admission, and a few days afterwards pain in the left side came on again. This, in three or four days, gave place to severe pain in the right iliac fossa. The bowels had been sometimes purged, at other times confined; but latterly were in the former state.

On admission she was very ill. The iliac pain was very severe, with great tenderness on pressure; the abdomen was collapsed; the knees were drawn up; the countenance was expressive of great distress; the skin was hot; there was no vomiting; the tongue was slightly furred; the pulse was small and somewhat hard, 128; the sounds of the heart were sharp, and the act of respiration produced pain. Sixteen leeches were applied to the region of the cæcum, and gr. j of calomel and of opium given three times a day.

The pain gradually subsided, the tenderness and fulness ceased, and in about ten days the patient was convalescent. It was questionable whether this severe pain was not peritonitis, consequent on ovarian irritation. The pain was rather lower than usual in cæcal disease; less dulness in the iliac fossa existed, and the onset of the attack was directly after the cessation of menstruation. Partial peritonitis evidently existed, but whether set up by ovarian or cæcal disease could not be satisfactorily ascertained.

CASE CXLI.—*Cæcal Disease. Recovery. (Reported by Mr. Bretcke).*—James B—, æt. 23, a draper's assistant, residing in the Borough, was admitted under my care into Guy's Hospital, November 2nd, 1861. Whilst sitting at breakfast, on the 31st of October, he was suddenly seized with severe pain on the right side below the ribs; the severity of the pain bent him double, and rendered him almost insensible; medical advice was at once sought, and medicine given every two or three hours, which, however, produced vomiting whenever it was taken. On the 1st November he was partially relieved, but was worse again in the evening. On the 2nd he was brought to the hospital; the pain, which had subsided, again returned severely, and was accompanied with an urgent but ineffectual desire to pass water. The bowels were acted upon once daily till the 31st October, but no evacuation was again passed till after admission. He had a pale and anxious expression of countenance; he had lost his appetite, and, since the attack, had been deprived of sleep. In the right iliac region there was defined hardness, imperfect resonance, and great tenderness on pressure; but the abdomen generally was neither distended nor tender. There was pain on passing water, and only a small quantity was discharged. The tongue was furred, and the mouth and throat were dry. The pulse was 64,

and compressible. There was no pain produced by respiration; the urine contained a small quantity of sugar. He was ordered, of soap and opium pill gr. v night and morning, and to have an injection of castor oil, and a diet of beef tea and arrow-root.

This treatment was continued, and the bowels acted, no vomiting was produced, and the tenderness in the cæcal region subsided, and on the 7th both pain and swelling had disappeared. The bowels on that day acted regularly, pulse 60, tongue clean, and there was no difficulty in passing water. Infusion of cusparia ʒj was given twice a day. On the 11th he left the hospital quite well.

In this case the symptoms of cæcal disease were well marked; there was severe pain, with hardness and swelling in the region of the cæcum, and with obstinate constipation; vomiting had apparently been produced by attempts to obtain action from the bowels; but by the continued use of opium, with bland, demulcent diet and a castor-oil injection, the severe pain subsided, the tenderness ceased, and the bowels acted naturally. No mercurial was given, nor did it seem necessary, for the bowels would not have acted more readily if it had been conjoined with the opium, and the convalescence would probably have been less rapid.

CASE CXLII.—The following is of great interest, as showing a state of *strumous inflammation of the cæcum*, in itself probably remedial, and allied, if not identical, with those previously detailed, but rendered fatal by its association with phthisis.

Ann C—, æt. 46, was admitted into Guy's Hospital under my care February 28th, 1855, and died March 30th. She was a married woman, who had resided in Southwark, and in her employment as a milkwoman had been much exposed to the weather. Some of her family had died from phthisis. For several years she had been subject to cough, but it had become more severe during the last eighteen months; menstruation had ceased, and her health had been much worse a few months before admission, and once she had suffered from hæmoptysis. She was thin and haggard, the face was slightly congested, and the physical signs were those of general bronchitis with phthisis. The pulse was irregular and intermittent, and there was a systolic bruit below the nipple. The bronchitis was slightly relieved, and then the signs of disorganization of the lung became more marked. Three days before death severe pain came on in the right side, accompanied with increased dyspnoea. Some irritation of the bowels supervened, but not to a great extent, and she gradually sank.

On inspection the abdomen was distended and tympanitic. The larynx

was healthy, the bronchi were much dilated; this bronchial dilatation was very marked on the right side, and on section the dilated tubes constituted a considerable portion of the surface. The mucous membrane in them was much congested, and covered with tenacious mucus; they were surrounded by crepitant lung. The bronchi on the left side were much less dilated. The left pleura was universally adherent, but the right was only adherent at its apex; at the right base the pleura was covered with pus-like lymph, and about a pint of pus was effused; a small vomica situated immediately beneath the pleura had given way into the serous membrane. There was a large, irregular vomica at the left apex, bounded by a smooth membrane with several intersecting bands. In the lower lobe were other smaller vomices and numerous miliary tubercles. The heart was healthy. There was a firm clot in the right ventricle, extending into the pulmonary artery. *Abdomen.*—There were old adhesions generally in the peritoneum, and several adherent cretaceous deposits. The *cæcum* was inflamed, and presented raised patches about the size of peas, soft, situated in the mucous membrane and containing pus; some of these collections of tuberculo-inflammatory product had given way, and slight ulceration was the result. The ascending colon was in a similar condition. The appendix cæci and the other portions of the intestine were healthy. The liver was fatty, the kidneys healthy, so also were the mesenteric glands.

This case might be considered as one of chronic bronchitis, and afterwards of phthisical disorganization. In the *cæcum* it is probable that the solitary glands became diseased, and low-organized product effused, which led to the production of minute abscesses and ulceration.

In many of the cases which we have detailed the patients were of strumous diathesis; but the disease, perhaps, had not advanced to disorganization nor to ulceration. There appears to be a greater tendency to this local enteritis of the *cæcum* in strumous than in other subjects. The condition just described is an earlier condition of what we so frequently find in both the *cæcum* and ileum in phthisis; here, however, the *cæcum* only was diseased.

CASE CXLIII.—Perforation of the Cæcum. Abscess extending to the Groin. Phthisis.—Michael R—, æt. 34, was admitted into Guy's Hospital under Mr. Key's care, in September 1835; he was a temperate man, but of strumous habits, and by trade a compositor. For a year and a half he had been subject to flatulence, indigestion, and occasional purging. Four days before admission, after four days of diarrhœa, he had experienced sudden pain in the right iliac fossa, where was a firm swelling, and pain had continued; the bowels were variable; the constitutional disturbance was slight; the pulse was soft and quickened, and the tongue was slightly furred. Leeches

were applied, and antimony was administered; suppuration became more manifest in the tumour, fluctuation became distinct, and hectic supervened. Six weeks after admission into Guy's an opening was made into the tumour, and 3viij of offensive pus evacuated; symptoms of phthisis gradually developed themselves, and he died the following June. Two openings existed above the right groin; they communicated with a contracted space, which was surrounded by dense membrane. The cæcum was found bound down by firm cellular adhesions to the neighbourhood of Poupart's ligament. The appendix was thick, opaque, and filled with a pasty fluid, and communicated with the cæcum. A sinuous canal of one and a half inch in length, narrow and apparently closing, led from the opening on the surface into the cæcum at its posterior part, nearly opposite to the opening of the ileum. The coats of the intestine were thickened, and the mucous membrane did not appear to be changed, except that a few contractions from cicatrices were evident. (Prep. in Museum, 1879²⁰.)

In this case the patient survived the immediate effects of the cæcal disease; the perforation, instead of setting up inflammation in the peritoneum, produced suppuration in the cellular tissue of the iliac fossa, and the pus was discharged near the anterior and superior spinous process. If there had not been any cicatrices in the cæcum, we should have questioned whether the disease had not commenced in the iliac fossa, and afterwards extended into the cæcum, as we have found to occur in connexion with the sigmoid flexure. The commencement resembled that of ordinary cæcal disease; but in its progress it might easily have been mistaken for abscess in the parietes of the abdomen.

Disease of the appendix sometimes exists without manifesting any symptom; this is especially the case in phthisis. The appendix often presents strumous deposit in larger or smaller masses; is often filled with fæces; and not unfrequently we find it distended with thin pus, with occlusion of the orifice, or with ulceration, without any pain or tenderness having been complained of, as in Case CXXV, and in the following instance:

CASE CXLIV.—*Bronchitis. Phthisis. Diseased Appendix Cæci*.—Thomas S—, æt. 40, was admitted January 7th, 1857, into Guy's Hospital, and died on the 20th. He was a tall, spare, anæmic man, who had been ill for six months, and on admission suffered from urgent dyspnoea and the symptoms of severe bronchitis. Inspection was made two days after death. The lungs were found studded with tubercles and congested throughout, and in some parts were in a state of early consolidation. At the apices there was iron-gray consolidation, and several dilated bronchi as large as a goose-quill, were

found to extend nearly to the margin of the lung; there were also several vomicæ. The peritoneum was healthy. The cæcum was rather more free than usual.

In the ileum were several healing ulcers, and strumous deposit was present in the substance of the mucous membrane. Close to the junction of the ileum with the cæcum was a flattened gland, infiltrated with strumous matter. In the cæcum was a small ulcer. The appendix cæci appeared full, and this distension arose from thin pus; or tracing the appendix towards the cæcum, the canal was found to be almost impervious close to the cæcum, on account of a sudden bend in the appendix, and from consequent adhesion. By carefully dividing this stricture, an ulcer about a quarter of an inch in diameter was found, which had extended through the coats of the intestine, except, perhaps, the peritoneal, which might have given way on removal. It was doubtful whether the peritoneum had not actually become perforated before death, and extravasation prevented by slight adhesion with the intestine.

In this case the symptoms of disease of the chest obscured every other symptom; but it is probable, that if life had been spared a few days, acute peritonitis would have been set up.

CASE CXLV.—Phthisis. Ulceration of the Larynx and of the Ileum. Concretion in the Appendix.—Thomas E—, æt. 18, a delicate, strumous lad, was admitted with phthisis on February 27th, and died May 4th. *On inspection*, eighteen hours after death, the lungs were found to contain strumo-pneumonic deposit, old gray induration, and a large vomica at the left apex. The larynx was deeply ulcerated at the inferior vocal cords. The ileum contained in its mucous membrane strumous deposit, and a large ulcer existed at the valve; the appendix contained a waxy concretion, white, lamellated, about one inch long, and placed at the extremity of the appendix; the rest was filled with mucus. This concretion appeared to be composed of inspissated mucus.

CASE CXLVI.—Pyæmia. Necrosed Humerus. Cæcal Disease.—William S—, æt. 72, was admitted into the hospital January 30th, and died February 16th, 1856. He had received six months before death a compound fracture of the left humerus, and Mr. Birkett had removed a portion of the necrosed bone; the wound did not heal, and the patient became increasingly prostrate; nine days before death he suffered from pain in the abdomen.

Inspection was made forty hours after death. The body was much decomposed; the lungs, liver, and kidneys were too much changed to decide as to the existence of acute disease. There was considerable development of fat; the peritoneum was greasy, and in the right cæcal region several coats of intestine were adherent; on removing them about a cupful of pus was poured out; this was found to arise from the neighbourhood of the appendix cæci. The appendix contained several small, circular ulcers, and one of these had a pinhole opening into the peritoneal cavity. The whole of its parietes were much thickened, especially at the extremity, which was white

and fibrous; the appendix contained pus. The cæcum itself, the ileum, and the rest of the intestines were healthy. There were no tubercles, nor evidence of phthisical disease in the lungs. The right shoulder-joint, the sternoclavicular articulation, &c., were filled with pus. On the left side was an oblique, ununited fracture of the humerus.

It is very unusual to find a patient at seventy-two years of age the subject of cæcal disease; neither did it appear to be the direct cause of death; the man died from pyæmia, consequent on necrosed bone. Cases, however, may arise of pyæmia produced by cæcal disease alone; the probability is, that in a poisoned condition of the blood, slight irritation at that part had been followed by ulceration, perforation, and subsequent suppuration.

CASE CXLVII.—*Perforation of Appendix Cæci. Laminated Concretion. Fatal Peritonitis.*—John H—, æt. 36, was admitted August 9th, and died on the following day. He was a strong, muscular man, by trade a carpenter, who, six days before death, had diarrhœa; this ceased, and was followed by constipation. Some months before he had had severe pain in the abdomen, but on admission he was too ill to give much account of himself. Inspection was made six hours after death. The head was not examined; the heart and thoracic viscera were healthy; but the diaphragm was much pushed up by the distended abdomen. The *abdomen* was much enlarged by the flatulent distension of the small intestine, resembling large, distended coils of the colon. The whole peritoneum was much injected, and every recess contained a collection of thin pus; there was a considerable quantity in the pelvis. On gradually removing the ileum from the cæcum, a small concretion was found in the peritoneal cavity near the cæcum; and on further inspection a slough was also found at the lower part of the cæcum, which contained a second concretion, about the size of a cherry-stone. The slough was found to be situated about the centre of the appendix cæci, the extremity of which was united to the cæcum immediately beneath the ileo-cæcal valve. The concretion was white at its centre, composed of laminæ; externally it was brown, and it easily crumbled into powder. The intestine itself, cæcum, and ileum appeared healthy; Peyer's glands very distinct from pigmental deposit; and there were some old adhesions at the lower part of the ileum.

Although the fatal attack commenced suddenly, the patient had some months before had pain in the abdomen; diarrhœa was followed by constipation; the latter the result of perforation, and the consequent peritonitis. The concretion was not analysed chemically; no nucleus was found in it; and it appeared to have been composed principally of phosphates.

CASE CXLVIII.—*Phthisis. Ulcerated Appendix Cæci opening into the Ileum.*
 —Y. D—, æt. 34, was admitted into Guy's Hospital February 23rd, suffering from phthisis; he died March 5th, 1859. The right pleura was adherent. A large vomica existed at the upper part of the right lung, and some white, strumous deposit. In the left was a vomica at the apex, and similar deposit as on the right side. The left lower lobe was in a state of red hepatization. There was considerable ulceration in the ileum, and miliary deposit on the peritoneal aspect of the ulcer. There were intestinal adhesions about the cæcum and in the pelvis, and the appendix cæci was concealed. On careful dissection the appendix was found almost beneath the cæcum, and adherent to a coil of the ileum; at the apex of the appendix was an ulcer, which had perforated all its coats, and those also of the ileum, so as to form a communication between these two portions of intestine. There was also ulceration of the ileo-colic valve and of the cæcum. The liver was slightly fatty. The spleen and kidneys were healthy.

This case is recorded to show a comparatively harmless sequence of perforation of the appendix cæci. The disease of the appendix had led to partial peritoneal adhesions, and its extremity had become firmly united to the extremity of the ileum. Perforation took place into the ileum, and established a second channel of communication between the small and the large intestine. The presence of this cæcal disease was not known during the life of the patient, and after death might very easily have been overlooked.

CASE CXLIX.—*Disease of the Cæcum following a Blow. Perforation of the Appendix. Suppuration. General Peritonitis. Almost complete secondary Perforation of the Cæcum.*—Christopher B—, æt. 21, was admitted into Guy's Hospital, in a dying state, June 1st, 1859, and expired a few hours afterwards. Two years previously he had received a severe blow in the region of the cæcum, but it was a week before admission that sudden pain came on in the abdomen. On inspection the abdominal serous membrane was found to be intensely injected and acutely inflamed. On separating the last coil of the small intestine from the cæcum, a small abscess was observed, which communicated with the appendix cæci; nearly the whole side of the appendix was destroyed by ulceration, and the pus had separated the coats of the intestine as far as the cæcum, into which it was on the point of forming a second opening. The mucous membrane of the rest of the intestine, and of the other viscera, were healthy.

The blow had probably in this instance set up cæcal disease, which manifested slight symptoms till perforation took place a week before his death. Suppuration followed, and was localized

by adhesions; but, most unwisely, he was taken from his bed, necessarily shaken in his transit to the hospital, which he reached in a dying state; the adhesions, on which the prolongation of life depended, were broken down, and general peritonitis established. The almost complete secondary perforation of the cæcum is an illustration of the course which suppuration sometimes takes.

CASE CL.—Local Peritonitis. Perforation of Appendix Cæci. Strangulation of the Ileum by the Appendix.—Miss — was a young lady about 23 years of age. About a year before her death she jumped from a gate, and experienced pain at the lower part of the abdomen, with slight vomiting and some uneasiness for several days; she remained, however, in apparently good health till August 13th, 1858, when, after partaking of veal and a glass of port wine at dinner, severe pain at the lower part of the abdomen came on, the pain extending from the hypogastric to the epigastric region. The next day the bowels were moved, but the pain towards the right iliac region still remained, and she felt ill. On the 16th her medical attendant saw her, and found her suffering from pain in the abdomen, with anxiety of expression in the countenance, which appeared to indicate more than ordinary colic. Aperient medicine was administered, and the bowels were moved on the 17th; she stated that the pain was partially relieved, and that she felt better. On August 18th, in the evening, severe vomiting of offensive matter came on; the pain continued at the lower part of the abdomen; the pulse was below 100, and there was very little tympanitis. Calomel and opium were given; ice was allowed; afterwards grey powder was administered, and mercurial inunction used; twenty-five leeches were applied to the abdomen. The vomiting ceased, and the pain partially subsided; the abdomen was still tense at the lower part, and the tympanitis increased; the pulse continued below 100; the urine was abundant; but the patient became more prostrate. A large injection of cold water was administered. The bowels acted at 2 a.m. on August 27th, and I was requested to see her in consultation the same day. At 10 a.m. I found her with a flushed face and with a distressed and anxious expression; the skin was clammy; the abdomen was flattened at its upper part; but below the umbilicus, and especially towards the right iliac region, it was very tense, tympanitic, and tender on pressure. A rounded and dense mass could be felt in the region of the cæcum and of the termination of the ileum. No evidence could be found of distension of the transverse or descending colon, nor could enlarged coils of intestine be observed through the parietes, and there was evidently greater distension of the right than of the left loin. The vomiting had ceased, the tongue was slightly furred, the gums were sore from the action of the mercury; about a pint of urine had passed, and menstruation had come on. The pulse was 120, and irritable; the respiration 40. She had not passed any flatus nor blood from the bowels, but an evacuation had been produced by the injection a few hours previously; she had had sleep during the night. She was recommended to omit the mer-

curial, and to take opium gr. ss every four hours; to have perfect rest and quietness, and to take ʒij—iij of brandy in divided doses, but frequently repeated, with nourishment, as arrow-root, beef tea, coffee, &c. On the 31st the pain had subsided in the abdomen, but had not ceased in the right iliac region; there was also increased borborygmi and restlessness; the tympanitis had increased, the pulse was 122 to 144, and there was also hiccup, with greater prostration. The opium treatment had been continued, and a blister had been applied to the seat of pain. During the day five or six fæcal evacuations were passed, with considerable relief. Till September 3rd she continued in a hopeful state, but was very prostrate, and the bowels acted. In the afternoon of the 3rd she suddenly awoke from sleep with great distress of breathing, and died in a few hours.

Post-mortem examination.—On opening the abdomen the peritoneal serous membrane was found to be dry, but no lymph was effused. At the left side of the cæcum there was considerable peritoneal inflammation, bounded by coils of ileum, and by the cæcum. At this part, deeply situated, the appendix cæci passed over the termination of the ileum, and extended to the mesentery of a portion of adjoining ileum, where it was strongly united, and formed, with its own mesentery, a firm loop. The mesentery of the appendix was adherent to a small gland, which completed the band of adhesion with the ileum; at the same site another coil of ileum was also adherent by a broad band of organised adhesion. The termination of the appendix was softened, sloughy, ragged, and perforated; but the appendix itself was pale. Above these partially constricting bands the ileum was somewhat distended, but the cæcum was also distended with flatus and fæces, showing that the strangulation had not been complete. The coats of the ileum towards its termination were exceedingly softened, and broke parts down in several parts on removal; but there was no evidence of fæcal extravasation before death. The cæcum was also softened. The opening into the appendix was free. The uterus and ovaries, &c., were healthy.

In this case there was evidence during life of local peritonitis in the neighbourhood of the cæcum and at the termination of the ileum; and although there was obstruction of the bowels, this was peculiar in its character; it was more severe than in ordinary cæcal disease—for, during a short time, there was stercoreous vomiting—and, on the contrary, it was less severe than in complete strangulation of the intestine, for the bowels were many times acted upon. In reference to the position of the obstruction of the bowels, the ileum and cæcum were manifestly concerned; there was dulness and tympanitic distension at that part, with tenderness, and there was no distension of the transverse nor of the descending colon; but the pain was situated near to the pubes, and nearer to the median line of the abdomen than in simple disease of the cæcum; and, although it was evi-

dent that the symptoms were principally due to inflammation connected with the cæcum or the appendix, it appeared more than probable that the pain which had been experienced a year before, and which came on after sudden muscular exertion, might have been induced by some abnormal movements of the intestine. These opinions were confirmed by post-mortem examination; the appendix cæci had been adherent for some time across the lower part of the ileum. A slight attack of enteric mischief produced pain, distension, and almost complete occlusion of the intestine; more severe inflammatory changes then became manifested, closely resembling those of ordinary disease of the appendix; softening and perforation of the appendix took place, but the peritonitis was localized by adhesions. The strangulation of the intestine then lessened, and the bowels were acted upon several times. Although prostrate and exhausted, there was hope of recovery; but sudden collapse supervened, and death followed in a few hours. This, we believe, arose from the extension of the disease to the general peritoneum, which was found to be dry, although sufficient time had not elapsed for lymph to be effused. The value of an opiate plan of treatment was well shown in this case, when associated with complete rest; the pain subsided, the bowels acted, and the vomiting ceased. The mercurials had rendered the contents of the canal more fluid, but had probably induced less plasticity and firmness of those adhesions, on the stability of which the prolongation of life depended, for by them the peritonitis was localized.

CASE CLI.—*Ulceration of the Appendix Cæci. Perforation. Fæcal Abscess. Secondary Perforation of the Ileum.* (From Museum Records.)—Thomas T—, æt. 24, a florid man, who stated that till his last illness he had enjoyed good health, was a ropemaker by trade, and in destitute circumstances. One month before admission he had diarrhœa, which several times recurred, and for which he took brandy. The diarrhœa was accompanied by severe pain. One week before admission he walked fourteen miles, and then slept in a stable, and four days afterwards was seized with vomiting, then with tenderness of the abdomen, and with constipation.

In that state he was first seen; the pain was principally in the right hypochondriac and lumbar regions; the vomiting was frequent, and the pain severe. In three days the pain subsided, and the bowels acted freely. He afterwards became prostrate; the abdomen continued tender, and before death slight delirium supervened. There was general injection of the peritoneum, which was covered with layers of fibrin and bathed with fæces. The

small intestines were distended; in the ileum was an oval patch of ulceration, at which the peritoneum was perforated at one or two places. Several other ulcers were found in the lower part of the ileum. In the right iliac fossa was a circumscribed fæcal abscess, formed by the cæcum, by the omentum, and by the ileum. The abscess contained the appendix cæci, nearly divided by an ulcer at its middle portion; the whole of the appendix was thickened and granular. (Prep. in the Museum 1832⁷⁶.)

The destitute circumstances in which this patient was placed increased the symptoms, and rendered the disease fatal. It is probable that after the commencement of the cæcitis, as indicated by the pain, &c., the patient would have recovered with rest and judicious treatment. He was obliged, however, to walk many miles, slept in a stable, and suffered great privation; in this state signs of peritonitis came on; this was probably caused by the ulceration and perforation of the appendix; fæcal abscess followed, and ended fatally in several weeks. The perforation in the ileum was, perhaps, secondary; it formed part of the walls of the abscess; and the ulceration had extended into it from the external to the internal coats. The peritoneal was more extensively destroyed than the mucous coats, so that several valvulæ conniventes were left.

CASE CLII.—*Fæcal Concretion. Perforated Appendix.* (From the Museum Records.)—Mrs. L—, æt. 45, was a very stout woman, and the mother of three children. For several weeks she had appeared unwell, and the bowels had been generally constipated. She was observed to complain of pain in the right iliac region, induced by coughing. On Saturday, 25th August, whilst engaged in washing, she suffered considerable accession of pain. The pain became more general and constant, and was greatly increased by pressure; leeches were applied, and she was bled, without relief; purgatives acted freely, but the stomach became very irritable; the tongue was red, clean, and parched. The abdomen then became exceedingly distended and tympanitic, and the tenderness general. The principal pain was in the right iliac region. The pulse was weak, but rather quick and sharp. The extremities were cold and clammy. The patient was restless, but sensible; and she died on the third day.

There was general injection of the peritoneum, which contained offensive pus, and some adhesions in the right iliac region. The appendix was short, and fixed by adhesions; it was distended by an elongated fæcal concretion about the size of the little finger, and two thirds of an inch from the mouth of the appendix was a minute ulcerated perforation. Other viscera were healthy, but the whole body was much loaded with fat. (Prep. 1881²⁰.)

At the onset of the pain, it was possible that the perforation had not taken place, and there can be no doubt that the injudicious administration of purgatives in this case hastened the fatal result. The cæcum had probably been distended for several weeks, and the pressure of the abdominal muscles during coughing had induced the impulsion of fæces into the appendix.

CASE CLIII.—*Concretion in the Appendix Cæci. Perforation.*—(From the Museum Records.)—George N—, æt. 20, a young man who had been accustomed to live intemperately, had had for several years occasional severe pain across the abdomen. Two days before admission into Guy's Hospital, in October, 1828, he was supposed to have enteritis, and was bled very freely. The pain, which was severe, was situated in the right hypochondriac region, extending to the back and to the pelvis; it was paroxysmal in character, and was increased by pressure and by the respiratory movements. The febrile excitement was slight; calomel and opium were administered, and purgatives were given. He died, apparently from syncope, after being removed from a warm bath.

On inspection, the general peritoneal cavity was found to contain opaque whitish puriform fluid, which escaped from the right side, where was a circumscribed cavity, bounded by the liver and gall-bladder, by the parietes on the outer and posterior aspects, and on the inner side by the ileum and colon; these parts were feebly glued together, and contained about twelve ounces of pus.

About two inches from the mouth of the appendix cæci was a soft brown concretion of the size of a large pea; and immediately beyond this was an ulcerated opening about three quarters of an inch in length, which very nearly detached the lower third of the appendix. The retained part of the appendix was much thickened; and there was also injection of the mucous membrane of the intestine. (Prep. 1881¹⁰.)

This case well illustrates the course of cæcal disease after perforation; and shows the importance of rest, and an opiate rather than purgative mode of treatment. At the time of his admission a fæcal abscess existed, and the peritonitis had become limited; but this attempt at reparation was entirely defeated by the use of purgatives, and by the mechanical disturbance of the abdominal viscera.

CASE CLIV.—*Fæcal Concretion in the Appendix. Perforation. Peritonitis.*—(From the Museum Records.)—Mr. C—, æt. 22, a well-developed man, whose general health had been good, after excess at table, on May 5th, 1829, suffered much disturbance of the stomach and bowels, but recovered partially in a few days. On the 9th, severe pain in the abdomen came on, with vomiting and prostration; he was depleted, and active purgatives were

administered, without relief; afterwards calomel and opium were given; there were delusive symptoms of improvement a short time before death, which took place on the 14th.

On the right side there was a circumscribed cavity bounded by small and large intestine and omentum, and it contained about 3ij of feculent pus. The appendix cæci was found bound down and distended to at least three times its ordinary size; and it contained an indurated concretion of about the size of a chocolate nut, which was composed of indurated fæces; a small perforation had taken place. Internally the mucous membrane was thickened, congested, and partially ulcerated; and there appeared to be inflammation of the adjoining parts of the ileum and colon. The valvula conniventes were abraded.

The concretion in the appendix had probably existed for some time, but did not produce any symptom till the period when vomiting and diarrhoea were set up. At that time, ulceration appears to have taken place, and on the fourth day perforation and peritonitis occurred, and a fæcal abscess was formed. With treatment which we now consider to be adverse, he survived for five days.

CASE CLV.—*Ulceration of the Appendix, with Disease of the Kidney.*—(From the Museum Records.)—William J—, æt. 36, was a thickly-set, corpulent man, with dark complexion. Sixteen days before admission, he was seized with severe pain in the right iliac region, which, on the following day, extended to the loins. From that time he passed less water than natural, and occasionally vomited; he was bled, warm bath was used and opiates given; he had rigors, and before admission a fit; at that time the intellect was disturbed, no urine was discharged, but a very small quantity of highly albuminous urine was drawn off. Occasional hiccup, with twitching of the face and grinding of the teeth took place. The bowels were freely purged. The mind became clear, but was again oppressed, and he had occasional rigors; more urine became secreted, but he had a sense of coldness of the skin. After free purgation he sank.

Adhesions were found existing between the colon, ileum, and the abdominal parietes; and the appendix cæci, with part of the cæcum, were destroyed by ulceration. The ileo-colic valve was also ulcerated. The kidneys were large and flabby.

The adhesions which had taken place in this case had entirely prevented fæcal extravasation, and it is probable that the cæcal disease was in an almost passive condition. The principal symptoms doubtless arose from ischuria renalis, which was also the cause of death; but the case is one of much interest, as

exemplifying one of the difficulties which might have arisen in the diagnosis. The severe pain in the right iliac fossa, from cæcal disease, might have been attributed to a renal calculus impacted in the ureter.

Many of these cases of perforation occur in early life. In my notes I have cases at the age of 12, 14, 19, &c., death generally taking place on the third to the seventh day; but although the leading symptoms are very similar, and well marked, each one has its own minor peculiarities. The detection of foreign substances in the appendix, without any severe irritation having been produced, is by no means uncommon; thus a pin was found with its head downwards, and its point extending into the coats, half surrounded by fibrous membrane. Again, I have observed an iron nail in the appendix, without injury having resulted from its presence; shot, and various substances are sometimes thus lodged. The presence of fæces in the appendix is often the precursor of ulceration and fatal perforation. In the volume of the Guy's Reports for 1856 is a case of much interest, recorded by Dr. Hughes, of strumous peritonitis and perforation of the cæcum coming on in a boy aged fourteen, after typhoid fever. Seven months after fever, while at work, sudden and severe pain came on in the abdomen, which subsided in a few days, but again returned, continuing for several hours in each attack. When brought to Guy's, the pain in the abdomen was general, with tenderness, and there was much febrile excitement. After several weeks the general distension subsided, but a hard, tolerably defined mass was felt in the region of the cæcum. This hardness continued, and he had occasional attacks of severe pain, sometimes with diarrhœa; hectic supervened, the skin became hot, the stomach irritable, and he exceedingly restless, fretful, and distressed; the abdomen moved *en masse*; he sank about ten weeks after admission.

There was slight strumous deposit in the lungs; but the abdomen presented the usual appearance of strumous peritonitis; the disease, however, was most marked in the region of the cæcum, the anterior surface of which was destroyed, and a fæcal abscess had resulted; the termination of the ileum was also perforated. Other parts of the small and large intestine were ulcerated.

The ulceration consequent on the typhoid fever in this child appears to have predisposed to slow organic changes of a strumous character in the abdomen.

CASE CLVI.—*Cancer of the Cæcum. Abscess in the Groin.*—William J—, æt. 56, by trade a coach trimmer, of very temperate habits, had enjoyed excellent health, till he ruptured himself in carrying a heavy weight; he afterwards had an abscess in the right groin, for which he kept his bed.

In October, 1855, he experienced pain and sense of heat at the lower part of the abdomen, and then found a swelling about the size of a walnut, which gave him great pain on pressure, or on walking. The swelling enlarged day by day, but became less painful; and night sweats came on.

On admission he had a cachectic appearance; and in the right iliac region was a hard swelling extending into the umbilical region; it descended also below Poupart's ligament on the right side; the inferior part was firmer than the upper; the pain was increased by pressure, and during defecation. The respiration was difficult; but the chest was normal; the urine was healthy, but there was pain after passing it. The appetite was tolerably good.

He was ordered to take a dose of castor-oil; four leeches were applied to the tumour; and Dover's powder with grey powder were given night and morning.

Free action on the bowels took place, which lessened the abdominal tumour, in fact it had almost disappeared; but the tumour in the thigh remained hard and tender. It became red, more swollen, and crepitant. Severe pain in the thigh then came on, and a free incision was made into the abscess; about a pint of fecal matter, with gas, was discharged. This fecal discharge with pus continued very abundant; the edges of the wound sloughed, and a second opening formed near the crest of the ileum. The patient gradually became prostrate; and for more than a month before his death he had very troublesome diarrhoea. He died about three months after admission.

On inspection of the abdomen the general peritoneal surface was found to be healthy; the small intestines were collapsed. There was an old inguinal sac on the right side quite free and empty. Several coils of small intestine, the lower parts of the ileum, were firmly adherent on the inner side of the cæcum, at the brim of the pelvis; and the cæcum itself formed the anterior surface of a firm tumour. On carefully removing the cæcum and intestine, it was found that the posterior wall of the cæcum was destroyed by carcinomatous ulceration, and offensive fecal matter was poured out from beneath the iliac fascia; and the abscess extended downwards to the opening on the thigh. There was also an irregular nodulated growth, extending from the mucous membrane of the cæcum, anteriorly attached near the valve, and surrounding the intestine; the edges were exceedingly vascular, but not flocculent. The section of the thicker portion near the ileum presented yellowish white medullary structure; and consisted of an aggregation of

large nuclei, evidently medullary cancer; near the margin, beautiful capillaries were observed distended with blood. At the posterior part some of the cellular tissue was infiltrated. The coil of ileum which was adherent to the cæcum had an irregular transverse opening into it, and was much injected. The remaining part of the intestinal canal was healthy, so also the mesenteric and lumbar glands. The liver was pale and somewhat fatty. The kidneys and bladder were healthy; so also the thoracic viscera.

The commencement of this disease was different from ordinary cæcal mischief. There was at first a small painful tumour in the abdomen, which had more resemblance to cancerous growth, or diseased gland, than cæcal disease of a simple inflammatory character; the disease was slow in its progress, hectic was developed, and the patient ultimately sank from exhaustion. The course of the disease beneath the iliac fascia was that followed in most cases of suppuration in the iliac fossa. In other cases which I have witnessed of ulceration commencing at the posterior surface of the cæcum, and leading to extravasation into the cellular tissue of the iliac fossa, the disease has generally been of a cancerous character.

CASE CLVII.—*Carcinoma of the Cæcum, of the Omentum, and of the Lumbar Glands. Faecal Abscess.*—James P—, æt. 21, was admitted into Guy's Hospital under the care of one of my colleagues. Three months before admission he discovered a small swelling in the right iliac region; when first noticed it was only the size of a marble, but gradually increased so as to fill the whole of that region.

On admission he was pale, nervously agitated, and had a strumous aspect; there was considerable febrile excitement, and he suffered from pain in the right iliac region, where a hard tumour could be felt extending into the lumbar region. At the left axilla was a small dark-coloured tumour, apparently cancerous; he had no difficulty in passing his motions, and his urine was healthy. The pain in the abdomen became more severe, and he died a few weeks after admission.

Inspection.—The lungs were emphysematous; the bronchial glands were not enlarged. In the abdomen, the peritoneum was acutely inflamed; the cæcum was perforated, so also the termination of the ileum, and the openings communicated with a faecal abscess at that part. The walls of the abscess were infiltrated with medullary cancer. The omentum was spread in front of the intestine and was adherent, and it formed part of the walls of the abscess; the lumbar glands were also infiltrated. The liver and kidneys were healthy.

This was apparently an instance of carcinomatous disease, but its precise origin or point of development was doubtful—whether

it began in the lumbar glands or in the cæcum, probably the latter. The detection of the gland similarly diseased in the axilla, very much assisted in forming a correct diagnosis.

CASE CLVIII.—*Colloid Cancer of the Cæcum. Jaundice.*—Ann C—, æt. 28, was admitted into Guy's Hospital July 31st, 1860. She had resided in service at Woolwich; and her previous health had been tolerably good, but the bowels had always been irregular. On July 22nd she was seized with vomiting, which continued till the time of admission, but was relieved by milk and lime water. The vomiting came on in the evening, and was preceded by pain and soreness in the mouth; pain then came on in the right side. Since July there had been no appearance of the menses; previously she had suffered from dysmenorrhœa. On admission she was pale, and had an anxious and distressed expression; she was nervous and easily excited. The pulse was compressible and the heart irritable; the chest was healthy, and she had no cough. The abdominal muscles were rigid; and immediately beneath the skin a defined, hard, and elongated tumour, placed somewhat transversely, could be felt in the region of the cæcum and ascending colon; the pain on pressure was severe, but was especially complained of when the pressure was sudden. The part was dull on percussion; the bowels were easily acted upon. After food flatulent distension came on, and she often complained of severe pain in the region of the cæcum directly after swallowing fluids. She was nervous and excitable, and the pain was manifestly increased when attention was directed to the part. Dover's powder, and the steel and aloetic pill afforded considerable relief, and she left the hospital on November 13th. She returned, however, in six weeks, much worse; jaundice had come on a few days previously, and she was very ill. The mind was oppressed, the pulse was very compressible, and she died in ten days from this hepatic complication.

Inspection.—The thoracic viscera were healthy. In the posterior wall of the right lumbar and ileo-hypogastric regions the peritoneum was studded with small yellowish tubercles of cancerous matter. The stomach was drawn down by the great omentum, so as to be visible below the liver. The ileum near the cæcum became gradually thickened and rigid, from the deposit of cancerous matter of the appearance and consistence of old honey in a crystalline state. The ileo-cæcal valve was obliterated; the cæcum and ascending colon were very thick and rigid; the peritoneum over them was involved in cancerous disease, and the muscular coat could not be distinguished. The thickening of the whole coat increased to an inch just above the cæcum, and the calibre of the intestine was much diminished in size; at this part also the mucous membrane was ulcerated for the space of several square inches. The mesenteric glands were enlarged and infiltrated with cancerous matter. The glands about the head of the pancreas compressed the common bile duct. The lumbar glands also were infiltrated. The kidney contained some cancerous tubercles. The right ovary was enlarged to the size of an egg, and was roughened and tubercular on the surface. The peritoneal veins at that part of the pelvis were enlarged. The walls of the cæcum and the last six

inches of the ileum presented a beautiful specimen of colloid cancer. The mucous membrane was nearly half an inch in thickness, and of a gummy appearance; it presented delicate fibres forming cellular spaces, which were filled with large nucleated cells; portions of the cellular interspaces consisted of very minute granules, as if from degenerating fibre; numerous clusters of granules were also observed. Beneath the thicker portion of the diseased mucous membrane was a red layer, consisting apparently of degenerated muscular fibre, or rather cancerous tissue, in the place of the muscular, and formed by an immense number of nucleated cells.

The cause of death in this case, was the obstruction of the bile ducts by enlarged pancreatic glands. The patient was an extremely nervous and excitable one, and at the onset of the disease it seemed doubtful whether the symptoms arose from ovarian irritation. It was, however, soon evident that there was organic disease of the cæcum or small intestine. The mischief was more circumscribed than in ordinary cæcal inflammation, and there was an absence of febrile symptoms, and of local peritonitis. It might have been doubtful whether ovarian disease had set up enlargement of the lumbar glands; but the manner in which pain came on in the tumour directly after food had been swallowed, pointed to intestinal mischief; and the tumour appeared in her emaciated state to be very superficial, immediately beneath the skin, and more distinct than enlarged glands would have been; neither was the growth in the position of mesenteric or omental tumours.

CASE CLIX.—*Appendix Cæci in the Inguinal Canal.*—James C—, æt. 16; the testes had not descended, and the appendix was adherent in the inguinal canal; the small intestine was fixed in the pelvis. The symptoms of hernia came on, and an explorative operation was performed. Peritonitis supervened, and after death purulent effusion was found in the abdominal cavity.

CASE CLX.—*Appendix Cæci adherent with the Omentum at the Internal Abdominal Ring. Supra-renal Capsular Disease.*—Henry T. M—, æt. 38, admitted into Guy's Hospital, March 1857, was a man of sallow, very anæmic appearance; and he had been losing strength for twelve months. He was exceedingly prostrate, unwilling to be disturbed or spoken to; and with the exception of occasional vomiting at the time of admission no symptoms of disease could be detected. He had no cough; the abdomen was collapsed, and he appeared to die from exhaustion. On inspection milary tubercles were found in the lungs, and some iron-gray deposit. The supra-renal

capsules were twice their natural size, and infiltrated with strumous product in a degenerating condition. The intestines were contracted and healthy throughout; one of the mesenteric glands was enlarged slightly, but there was no evident pressure on the thoracic duct. The omentum was firmly adherent in the right inguinal canal, and at the orifice the appendix was also inseparably united. It contained some semi-purulent mucus, and was slightly dilated at its extremity. This condition of the appendix was not known to have produced any symptom recognisable during life.

Intussusception of the ileum into the cæcum and ascending colon is of not unfrequent occurrence; and its symptoms might be mistaken for simple inflammatory disease of the cæcum. This subject will be again referred to in speaking of intussusception generally.

In many of the morbid changes of the cæcum to which allusion has been made, constipation is a more or less constant sign, and there is danger of yielding to the temptation of administering purgatives; these medicines rarely effect the desired object until inflammatory irritation has subsided.

CHAPTER IX.

ON DIARRHŒA.

DIARRHŒA consists in the abnormal frequency of evacuation from the bowels, as defined by Cullen, "Dejectio frequens; morbus non contagiosa; pyrexia nulla primaria:" and it arises generally from an irritated condition of the large intestine.

It manifests itself in various forms, some of which have received distinctive appellations, as *Diarrhœa crapulosa*, *biliosa*, *mucosa* or *catarrhalis*, *dysenterica*, and *choleraica*, to which might be added *nervosa*, and *colliquativa*.

Diarrhœa crapulosa is that state in which there is an unnatural fluidity and excess of fæcal excretion, in which the evacuations are healthy in character, but in excessive frequency and fluidity; in some cases very large quantities are discharged without any discomfort, but, on the contrary, with relief to the patient. This form of diarrhœa should not be checked when it is a natural discharge; but more frequently it is the sequence of irritating and undigested food; either, that too great a quantity has been taken, and a portion of it has passed into the intestine crude and partially dissolved, or from the insoluble character of portions of the food in the gastric juice, unchanged ingesta, as the woody fibre of vegetables and fruit, irritate the intestine. Again, active mental or bodily exercise immediately after a meal, which has been suitable both as to quality and quantity, may interfere with the proper solution of food, and lead to its hasty passage into the duodenum.

When the alimentary canal becomes in this way loaded with undissolved ingesta, pain of a griping and twisting character, from irregular peristaltic action and from distension, ensues.

The abdomen becomes full ; and the skin and complexion sallow ; the tongue is furred ; the pulse is compressible ; headache and giddiness are often present ; the sleep is disturbed ; the bowels act frequently and irregularly, and the motions contain undigested substances, with fluid fæces or with firm scybala. Considerable soreness is at times experienced in the course of the large intestine, and distressing tenesmus arises from the irritation of the mucous membrane of the rectum.

The term *lientery* is used to designate the condition in which the food is passed almost unacted upon, either by the gastric or intestinal secretions, and is often discharged in a very short time after having been taken ; this state arises from excessive irritability of the whole intestinal tract, with disordered secretions ; it is not unfrequent in children after protracted diarrhœa, and gastro-enteritis. It is of common occurrence among the out-patients of large hospitals ; and in not a few cases leads to fatal termination.

Bilious Diarrhœa is also a form of disease produced by the effusion of irritating substances into the intestine ; not, however, from without, but from the liver, and possibly from other glands.

The secretion of the liver becomes either excessive in quantity, or irritating in quality ; and the contents of the canal are apparently hurried onward, producing frequent loose and bilious dejections. The causes of this state are various, and sometimes the disorder of the liver is really secondary to an irritable condition of the intestine itself, due to excess, especially of stimulants. Exposure to cold and wet induces diseases of this kind, especially in the autumnal season of the year. The symptoms are somewhat similar to those previously mentioned ; the pain is slight, unless the disease become aggravated ; the tongue is furred ; the complexion is sallow ; some febrile excitement is present with frontal headache ; pain in the abdomen and in the hypochondriac region. This form of diarrhœa is sometimes epidemic, attacking considerable numbers exposed to similar exciting causes ; and when severe, and accompanied with colic or spasmodic pain in the abdomen and legs, and especially with vomiting, it constitutes English cholera, and often leads to great prostration of strength ; then, the coun-

tenance becomes haggard, the eyes appear sunken, the pulse is exceedingly compressible and failing, the tongue is brown, and the patient too frequently sinks exhausted, especially if very young, or advanced in life, or if already prostrate from other disease.

Abnormal conditions of the bile tend to produce other modifications; thus, the motions in diarrhœa are sometimes in a state of fermentation; they are watery, frothy, and only contain fluid fœcal matter; as in the following case of phthisis, in which there was probably some ulceration of the intestine. On admission into Guy's Hospital the evacuations consisted of long shreds of mucus and casts, which were composed of columnar epithelium and nuclei; after a few weeks this condition subsided under the use of cusparia, sulphuric acid, and opium, with occasional starch injections; but it was followed by very severe pain in the course of the colon, and by frothy, yeast-like evacuations. The symptoms of disease of the chest became less marked, the cough less trying, but the patient continued very prostrate, the tongue became furred and brown, and the pulse very compressible. For this state I used injections of charcoal,* ʒij to about a pint of thin barley-water, with great relief; the character of the evacuations improved, and in a short time became naturally fœcal, the pain diminished, and the strength increased. I afterwards gave her several grains of myrrh, twice or three times a day, with manifest improvement, till she left the hospital several months later.

On the contrary, diarrhœa sometimes occurs with an absence of bile in the evacuations; in jaundice this may be the case; it is so in cholera; and towards the close of chronic disease, the liver may cease to pour out its ordinary secretion; thus, in a patient slowly sinking from the exhaustion consequent on diabetes, diarrhœa existed for many weeks before death; it was sometimes checked, but only to return with greater violence, and with much irritability of the stomach. The motions were often quite white, like water, frothy from an abundance of soap.

There is, also, a form of diarrhœa arising from the inhalation of noxious effluvia, which is closely allied to that just described; the fumes of sulphuretted hydrogen gas are absorbed by the

* See Dr. Theophilus Thompson's Lectures on Phthisis.

lungs, and through their minute capillaries enter the blood ; it is circulated, and acts as a poisonous agent on that vital fluid, and if concentrated, proves rapidly fatal ; if less concentrated, it produces headache, and frequently, also diarrhœa. It appears, that not only are the secretions of the liver and alimentary canal changed, but that, in this way, the blood rids itself of the poison. So rapid is this agent in its action, that to be present for a short time, even a quarter of an hour, in a dissecting room, will, in some, produce distressing diarrhœa.

In typhoid fever, and in phthisis, ulceration of the small intestine is frequently found to be accompanied with diarrhœa ; of these we have elsewhere spoken ; in some of these cases, the large intestine is involved, but in others, when the diarrhœa has been severe, such has not been the case. It would appear that the continuity of structure with the ulcerated ileum, the irritating excreta, as well as the changed and probably accelerated peristaltic action of the small intestine, tend to produce a corresponding state of the colon, and thus to set up diarrhœa.

Catarrhal and *mucous diarrhœa* arise from a state of slight inflammatory disease, closely allied to ordinary coryza, affecting the mucous membrane of the *large* intestine ; the secretion is at first checked, but afterwards greatly increased, and a watery fœculent mucus is discharged mixed with the ordinary fœces ; this state may continue for several days, or even for a longer period ; the motions are loose, and somewhat watery, and if the rectum be affected, considerable tenesmus is produced ; the pain and febrile excitement are slight, but the strength of the patient is reduced, and he is unequal to his usual duties ; the tongue is clean, the pulse is compressible ; the bladder sometimes sympathises with this irritation, and a frequent desire to pass urine is induced ; in little girls, also, a muco-purulent secretion often takes place from the vulva ; redness of the parts is produced with smarting pain, and the idea has sometimes been suggested that the child has been cruelly treated.

In this form of diarrhœa the evacuations contain a considerable quantity of mucus ; this is especially manifested when irritation occurs very low down in the rectum, or is set up by hæmorrhoids ; and the mucus will sometimes pass both before and after the dejection.

This catarrhal diarrhœa not unfrequently becomes a chronic

disease, the more severe symptoms cease, but still the bowels do not act in their normal manner ; constipation often ensues, and afterwards a fresh looseness of the bowels, and this alternation is oftentimes repeated, or the more solid motions are followed by a discharge of mucus coating the fæces ; sometimes the mucus is passed in considerably quantity after the evacuation, or it forms an elongated flake or cast of the intestine. I have observed this condition following severe disease of the intestines of a dysenteric character, and it is sometimes associated with a state of chronic congestion of the liver ; again, it is often perpetuated by the presence of hæmorrhoids, and by ovarian disease.

In infants it closely resembles gastro-enteritis, or is, perhaps, rather identical with it, but differing in degree, as a greater or less part of the alimentary canal is diseased ; in them the whole tract sometimes becomes rapidly affected, and great, if not fatal prostration, rapidly ensues. (See Muco-Enteritis.)

As with bilious diarrhœa, before mentioned, it is in very young or aged subjects that catarrhal diarrhœa, or catarrhal inflammation of the large intestines, leads to more serious disease, as well as amongst those in whom chronic or more exhausting disease has existed.

There are several *pathological* changes observed in the intestines of patients who have died from diarrhœa ; but in many instances, where severe diarrhœa of the forms already described has existed and led to fatal results, the appearance of the mucous membrane has been normal, its congestion has entirely disappeared, and a thin mucus is found upon the membrane ; or, secondly, a vivid injection is presented in more or less isolated patches.

Thirdly. When the diarrhœa has been chronic, the mucous membrane is not unfrequently covered by a thick layer of mucus, and presents a gray colour. I have frequently examined membranes thus changed (as before described ; see Duodenum and Cæcum,) and have observed that the gray colour arises from minute particles of dark pigmental matter deposited in the substance of the mucous membrane. The continued congestion of the membrane probably gives rise to effusion of colouring matter from the blood, changed hæmatine.

In the large intestine, this pigmental deposit is found in

minute circles around the follicles ; the mucous membrane, and also the connecting cellular tissue become thickened.

A fourth condition is observed after long-continued diarrhœa, namely, that of minute ulceration, probably follicular, extending nearly the whole length of the colon. (Such a preparation is found in the Museum of Guy's.) These ulcerations are about one sixteenth of an inch in diameter, and present a minute black zone around each of them. This state would be regarded by many as the result of dysenteric disease.

Dysenteric Diarrhœa.—Purging is the most marked symptom of dysentery, and the lesser degrees of irritation which we have considered under the term of catarrhal diarrhœa might be regarded as a form of dysentery of the mildest character. In dysentery, however, the diseased mucous membrane rapidly passes into a state of ulceration, and blood is discharged with the fæcal excreta.

In *Choleraic Diarrhœa* a thin, very abundant watery mucus is discharged from the alimentary canal. The evacuation may have very little colour, and have the appearance of rice-water. It is often alkaline in character, and consists of nuclei and epithelial cells in various degrees of development. After death the membrane is found to be entire, and pale or sodden, the solitary and Peyer's glands are enlarged. In many cases of uncomplicated cholera which I have examined, no further morbid appearance was presented.

The symptoms are those of rapid prostration, with pallor and sunken eye ; the pulse is compressible, the tongue is cool, and the voice is often scarcely audible ; the abdomen is collapsed, and the urine is scanty in quantity ; the stomach is often exceedingly irritable, so that everything is at once rejected from it ; the alvine evacuations are generally frequent, and of the character before mentioned ; and severe cramps in the legs and in the abdomen are often present. This state may pass into one of profound collapse ; but as the prostration subsides, in favorable cases, I have never observed the febrile excitement which is secondary to true cholera ; even after a single evacuation of the character of rice-water, profound prostration may ensue.

Another kind of diarrhœa is that which has been correctly called *Serous*, and which is frequently observed in albuminuria.

A dropsical condition of the mucous membrane is induced, and the serous exudation from the overcharged capillaries leads to watery discharge into the colon, and thus to diarrhœa. This state of the mucous membrane is precisely analogous to the œdema of the lungs, and to anasarca of the cellular tissue in renal disease. So frequently is diarrhœa present in these cases, that it may almost be regarded as a symptom of the disease, and when moderate is beneficial in its results. It is the action we often seek to produce artificially by powerful hydragogue cathartics, so as to diminish the quantity of urea circulating in the blood, and to relieve the oppressed kidney; for these fluid evacuations contain urea, as does the gastric juice and the mucus discharged from the lungs.

Another class, which can scarcely be placed among the cases previously mentioned, are those arising from fright, from excessive mental agitation, from want of food, and from exhausting disease; the former cases are of mental origin, the latter constitute what is sometimes called "colliquative diarrhœa;" and the condition of the mucous membrane corresponds to that of the skin, from which profuse partial sweats break out.

In fright the capillaries of the face become blanched, and the blood leaves the whole of the surface; the cavities of the heart are increasingly distended, hence the discomfort there experienced, and the mucous membrane of the intestine is probably also engorged; therefore the discharge from the mucous membrane is to a certain extent beneficial in relieving internal congestion. The intimate connection of the sympathetic nerve with the centres of thought and feeling is the probable explanation of these instances of diarrhœa following mental agitation.

In scurvy, purpura, starvation, &c., the altered character of the blood, and its inadequate capability of sustaining the changes of nutrition, lead to the effusion of serum, or blood, into the mucous membrane, or into the canal itself, corresponding to the effusion into the skin. In some fatal cases of purpura, the whole of the mucous membrane of the alimentary canal is studded with spots of ecchymosis. An interesting case of this kind occurred at Guy's Hospital in 1856, in a young man who had been starved to death. (Case CLXI.)

Discharge of blood, or melæna.—Obstruction of the portal circulation, either from pulmonary, from cardiac, or from hepatic disease, leads to great engorgement of the mucous membrane of the whole alimentary canal; and this congestion may cause hæmorrhage from the bowels. In examining the mucous membrane in these cases, it is very common to find points of ecchymosis, and capillary vessels of the membrane much distended. Under a low magnifying power we find the capillaries beautifully injected, with extravasated blood between them, still, however, restrained by the basement or bounding membrane: if the rupture of this membrane occur blood is extravasated. The discharge of blood may be a symptom of various diseases; thus, ulceration is a frequent cause of hæmorrhage from the bowels, and the ulcer may be located in any part of the canal, as in the stomach and duodenum; in the small intestine in fever and in phthisis; in the colon in dysentery, &c.

The blood does not always present the same appearance, if it arise from hæmorrhoidal vessels the blood will be florid, and precede or follow the dejection; if higher in the canal it is incorporated with the fæces; and when it has traversed a considerable portion of the canal, it becomes discoloured by the secretions from the membrane. This is the case, to some extent, when the blood is poured into the cæcum, but is especially so whenever it has been extravasated into the stomach; the acids of the gastric juice act upon the effused blood, which becomes blackened, and is discharged from the intestine as a pitchy fluid, constituting true melæna.

The *symptoms* of diarrhœa have, perhaps, been sufficiently described in mentioning its several forms; and they vary according to the cause. In the simplest form there is neither pain nor constitutional disturbance; in more aggravated cases there may be severe colic, and febrile excitement; and generally, unless there be hepatic disturbance and derangement of the whole mucous tract, the tongue is clean, it is then furred and injected, and in typhoid prostration assumes a brownish colour. The pulse is compressible, because the nerve of organic life sympathises; and the consequent prostration is often very alarming, especially in infants and aged persons, and in some cases it leads to a fatal result.

It is important carefully to mark the character of the evacuations; first, as to the admixture of undigested substances; secondly, as to *fluidity* of the evacuations; a simple fluid state, with normal excreta, indicates irritation of the mucous membrane in a slight degree; thirdly, the *presence of mucus* is evidence of more severe irritation of the colon; and it is sometimes found in excessive quantity when the evacuation is poured from one vessel into another; fourthly, if more acute disease of the colon exist, detached portions of fæces are found floating on the fluid, which from the rapidity of its discharge, and possibly also from intestinal changes, is often frothy, from the admixture of air; fifthly, thin watery fluid may be discharged in severe diarrhœa, with scybala, and with sedimentary portions of fæcal matter;* sixthly, thin fluid, almost like clear water, may be discharged, as in some cases of albuminuria, from an œdematous condition of the membrane, or like rice-water in choleraic diarrhœa, or like soap-suds when with colliquative diarrhœa the hepatic secretion is also checked; seventhly, the fæces are sometimes discharged in a state indicative of fermentative action, and a frothy surface is produced of the appearance of yeast, and the whole discharge closely resembles the matters occasionally ejected in obstructive disease at the pylorus; eighthly, as to the *colour* of the evacuation, we have evidence thereby of the excess and of the paucity of *bile*, sometimes the stool being of a deep brown colour, at others almost as pale as chalk; ninthly, the colour may be changed by the admixture of such substances as logwood administered medicinally, or blackened by steel medicines, the sulphuret of iron having been formed; and tenthly, the colour is a guide to the detection of blood. Blood in the alvine discharges may be only observable by microscopical examination; but if in larger quantity, the colour varies from the ordinary appearance of blood to the black pitchy stool of melæna, as we have before mentioned, according to the position of the hæmorrhage in the canal. The green colour of the discharges in the severe diarrhœa of children, we believe, with Dr. Golding Bird, to be altered blood from an irritated and perhaps aphthous surface. Again, in severe dysentery,

* Dr. Osborne "On the Examination of the Fæces." 'Dublin Quar.,' 1853.

thin watery fluid, like the washing of beef, is sometimes discharged, which consists of blood with mucus, and of imperfect epithelial elements. To these dysenteric evacuations we shall have again to refer. Lastly, the *odour* of the *fæces* is not altogether unimportant; sometimes they are intolerably fetid from rapid degenerative changes, at other times they have scarcely any odour. In many instances the microscope enables us to detect an excess of mucus, the presence of blood, the rapid discharge of epithelial elements and nuclei, and other organic and inorganic substances, which the unassisted eye would in vain search for. We have elsewhere referred to the occasional presence of phosphatic crystals upon the mucous membrane of the intestines, and they are sometimes found in the alvine discharges, in simple as well as in typhoid diarrhœa. The presence of fatty matters in the evacuations was first noticed by Dr. Bright, in connexion with disease of the pancreas; and the observations more recently made in reference to the physiological effects of the pancreatic fluid have directed increased attention to the subject, but we sometimes find oleaginuous substances discharged after the administration of large quantities of milk and of cod-liver oil; thus in one case masses of fat as large as filberts were sent to me by a patient affected with phthisis, who had partaken of milk very freely; still, the observation has been confirmed by subsequent observers, that fatty matters are discharged in the alvine evacuations sometimes in disease of the pancreas, and sometimes in extensive disease of the mesenteric glands.

The *causes* of diarrhœa have been partially referred to. The most common cause of ordinary diarrhœa is exposure to cold and wet; standing in damp places; allowing the legs and loins to become damped and chilled; sitting down upon the ground, and falling asleep in the open air; injudicious bathing; the habit of leaving off flannel garments in hot weather, by which perspiration more rapidly evaporates, and the blood is driven from the surface towards the internal organs.

2nd. Improper and indigestible food, unripe fruit, and an *excess* of uncooked fruit; salads, pastries, and much that modern cookery produces, especially when an excess in quantity is combined with an injurious quality.

In infants a fertile source of diarrhœa, often passing into severe gastro-enteritis, is the administration of unsuitable food, the injurious effects of which are greatly increased by exposure to cold. In hospital and dispensary practice, this cause of disease is observed to a frightful extent; at seven or eight months, even while the infant is, in a great measure, nourished by the breast of the mother, meat, raw vegetables, and fruits, sweets, almost *ad libitum*, are given; and a few months later we often find, that before the child has the power of mastication, the mother gives the food of which she herself partakes, sometimes adding malt liquors and ardent spirits. The consequences of this dietary are such as might be anticipated; the food passes onwards undigested, and severe gastro-enteritis is induced; and the malady is aggravated by a want of cleanliness, and by exposure to night air and dampness. The mortality in London from these causes is exceedingly great. In other infants the food, although in itself proper, is unsuited to the condition then existing, and perpetuates diarrhœa; or it may be, that the milk of the mother disagrees with the child, from the impairment of her health. In such subjects we occasionally find, that an alteration in the character of the gastric juice of the infant leads to coagulation of the milk, and to severe diarrhœa, with colic, &c., the stools containing portions of curdled and undigested milk, namely, oleaginous matter mixed with casein.

3rd. Diarrhœa is set up by exhaustion, either from want of food, starvation, and its attendants of misery, or as the consequence of chronic disease. This form of diarrhœa is sometimes observed in women who have nursed their infants too long. Enfeebled by bearing children rapidly, their strength is additionally taxed by nursing for twelve, fifteen, or eighteen months without proper nourishment and invigorating air. The whole mucous membrane is affected; the nerve of organic life shows its ebbing powers; the blanched cheek, the dilated pupil, the desponding countenance, and impulses of a mind verging on insanity, are symptomatic of this condition. There is intense pain in the head, the heart is enfeebled, the pulse sharp, and sometimes irregular; there is distressing sensation of exhaustion at the scrobiculus cordis, with severe pain in the back, and with this a very slight irregularity of food will sometimes set up

diarrhœa and vomiting. Cancerous and strumous disease of the mesenteric glands, obstruction of the thoracic duct, chronic disease of the pancreas, diabetes, &c., sometimes have uncontrollable diarrhœa as one of their latest symptoms.

4th. *Epidemic causes*.—At some seasons of the year, in our own climate during the spring and autumn months, diarrhœa of varying severity is set up, and appears to arise from the condition of the atmosphere, perhaps from germs of vegetable or animal growth.

5th. *Endemic causes* are more numerous, and with them may be classed the diarrhœa arising from offensive drains, from decaying animal and vegetable matters. Causes of this kind operate with greater severity upon the young and enfeebled, upon the strumous and ill-nourished. Many infants are thus affected with diarrhœa, and with severe general gastro-enteritis. Dr. Snow has shown that the impure water supplied some time ago in several districts of London, containing, as he stated, the refuse of sewers and closets, led to severe diarrhœa, if not, as he believed, to cholera. Again, a general dampness of locality, as from a clay subsoil, will set up, or will increase and perpetuate diarrhœa. We have witnessed the removal into dry bracing air followed by cessation of the disease, and the return to the same district repeatedly cause its recurrence.

6th. *Excessive secretion* of bile, and other diseases of the liver, as well as disease of other intestinal glands, set up diarrhœa.

7th. Other causes are, strumous disease of the mucous membrane of the intestine and the mesenteric glands. 8th. Œdema and long-continued congestion of the mucous membrane. 9th. Mental agitation and fright. 10th. Ulceration of the small and large intestine, as in fever, phthisis, &c. 11th. Cancerous diseases. 12th. Purpura and scurvy. 13th. Large draughts of water. 14th. Miasmatic disease. 15th. Poisons.

Prognosis.—Diarrhœa is never altogether free from danger in aged persons, or in very young children; but the prognosis differs according to its cause and character. If associated with chronic disease, or an enfeebled condition of the system, it is often the immediate precursor of death; but when the cause can be removed, and the subject is young, however severe the case may be, we should encourage the prospect of recovery.

Many of this class, who have appeared quite *in extremis*, have gradually and almost miraculously recovered.

The prognosis is unfavorable, when diarrhœa has been long continued, and is very severe in its character; in some of these cases scarcely any treatment appears to arrest the purging, and the patient gradually sinks into a typhoid condition.

It may appear unnecessary to say anything in reference to the *diagnosis* of diarrhœa; it is well, always, if possible, to ascertain personally the character of the evacuations; but in some there may be apparent diarrhœa, without the reality; I have seen starch enemata used, when patients were greatly exhausted, and on inspection, found the intestine loaded with solid fœcal matter; and, in spinal disease, a weak sphincter and with involuntary defecation is often mistaken for diarrhœa, and I have known astringents continued for several months ineffectively, whereas rest to the spine quickly relieved the malady. Or again, a single loose motion may constitute diarrhœa, and require immediate attention; for the character rather than the quantity should be our guide.

Treatment.—The primary object must be to ascertain the character of the diarrhœa, and to remove, if possible, its cause.

If food be improper, to change it, and administer such as shall be of the least irritating kind.

If the air be impure, to remove to a healthy atmosphere.

If the mucous membrane and the secretions be disordered, to try and restore them to a healthy state.

To check the diarrhœa by various astringents and by rest.

Warmth.—Warm baths, warmth applied to the feet, and flannel to the abdominal parietes, a warm but pure air, &c., assist in checking many of the simpler forms, and in diminishing those arising from chronic disease. Local warmth may be attained by the application of a hot fomentation, or poultice, to the abdomen, or by such rubefacients as a mustard poultice, or turpentine embrocation.

Food.—In diarrhœa the least irritating and the most easily digestible kinds of nourishment are advisable. Many of the forms of amylaceous aliment, arrowroot, sago, are of this kind, and may be given made with milk; these are in themselves soothing applications to irritated mucous membranes,

whilst they serve as nourishment to the system. Milk, rice, soaked bread and toast, lightly-boiled puddings of flour and eggs, &c., may be also taken with advantage, and in chronic diarrhœa suet and milk is often of great benefit.

The avoidance of stimulants, of rich and greasy food, of highly seasoned dishes, of vegetables, especially when uncooked, of fruits, &c., is essential; and it is well in many cases to abstain for a short time from solid animal food altogether. That form of animal food which is least irritating is chicken, sweet-bread, and some forms of fish, as sole, cod, and whiting; then venison, mutton, and beef; but much depends on the mode in which these viands are dressed. When dried, salted, and cold, they require a much longer period for their digestion, and portions often pass into the intestine undissolved. Beef-tea sometimes appears to increase diarrhœa, when veal and mutton broth can be taken with benefit.

Rest, and the avoidance of muscular excitement and sudden movements, are very important in checking diarrhœa; and in many instances, especially in severe cases, a recumbent posture should be maintained. In the erect position the gravitation of fluids increases their rapid movement over the irritated mucous membrane.

Pure and dry air is very desirable; many patients at once recover when removed from a damp atmosphere to a dry and bracing one; and when the contamination of decomposing animal and vegetable substances is setting up the disease, removal is still more important, and is often essential to permanent restoration. In miasmatic districts, diarrhœa may not only be rendered paroxysmal, but be perpetuated by the marsh poison.

Many cases of diarrhœa will be cured by this attention to warmth and diet, to rest and pure air; but other means often promote the comfort and favour the restoration to health.

If the large intestine, and especially the rectum, be affected, much benefit is derived from *injections* or *enemata*. These are of various characters, simple starch, thin gruel, and barley-water; and to these we may add tincture of opium and biborate of soda. Or the enemata may be more astringent, as decoction of oak bark, with tragacanth; or a very dilute solution of nitrate of silver may be used.

To restore the diseased mucous membrane and to correct secretions.—The alkalies are of very great service in diminishing congestion, as well as in rendering the secretions less irritating. Solution of potash, lime-water, chalk, some salines, as chlorate of potash, bicarbonate of potash, and nitrate of bismuth, act in this manner.

When the hepatic secretions are disordered, as shown by furred tongue, and pale evacuations, the moderate use of mercurials is of value, as grey powder or calomel, combined with Dover's powder, with soda or with opium; but we should strongly urge that mercurials be very carefully administered, because in many forms of diarrhœa they tend greatly to aggravate the disease. It is only in some cases, even with a foul tongue, and deficient hepatic secretions, that we would recommend their use.

Demulcents.—These act by directly sheathing the mucous membrane; the most important are those mentioned as food, but others are of considerable utility, as acacia, tragacanth, linseed, liquorice, glycerine, spermaceti, &c.

Castor-oil, Linseed-oil.—These are of great value, when improper food, retained secretions, and scybala irritate the alimentary canal. They are combined with great advantage with the compound tincture of rhubarb, and sometimes with a small dose of opium, \mathfrak{m} v or x. These remedies are of most service in some forms of dysenteric diarrhœa, when scybala irritate the mucous membrane.

Ipecacuanha is a remedy which acts, apparently, on all the mucous membranes, and is as valuable in disease of the alimentary as of the respiratory mucous membrane. *Ipecacuanha* not only increases the quantity of mucus from the membrane, but mitigates inflammatory congestion. It is of great service in the dysenteric diarrhœa of adults, and equally so in the diarrhœa of infants. In the former, Dover's powder is a valuable form of administration, or the *ipecacuanha* may be combined with astringents, as in the compound decoction of *krameria*, and the compound logwood mixture of the Guy's Pharmacopœia.*

Astringents and Desiccants.—These may be divided into

* 'Decoctum *krameria* compositum: *Krameria* rad. \mathfrak{z} ix, aquæ ferventis \mathfrak{z} xviii; decoque ad \mathfrak{z} xv et cola; deinde adde vini *ipecacuanha* \mathfrak{z} vj, tinct. catechu \mathfrak{z} vj, syrapi \mathfrak{z} iss.'

several classes. The *saline*, as chalk ; the *vegetable*, as krameria, kino, catechu, logwood, Indian bael, cusparia, opium ; *metallic*, as sulphate of copper, acetate of lead, nitrate of silver, nitrate of bismuth, &c.

Opium acts not only as an astringent, but also as a narcotic ; it diminishes the secretion from the mucous membrane, and the peristaltic movement of the intestine, and it relieves the pain of colic. It is of great value in diarrhœa, and may be combined with other remedies in the simple forms, as with chalk and ipecacuanha ; but when irritating ingesta and disordered secretions perpetuate diarrhœa, opium and astringents are not appropriate remedies. When the disease is chronic, opium may be given, with the more active vegetable astringents, catechu, krameria, and logwood, and sometimes very advantageously with quinine.

The metallic astringents are combined in a similar manner with opium and ipecacuanha, but are more frequently used in chronic dysentery, and in strumous ulceration of the intestine in phthisis, than in simple diarrhœa.

Mineral Acids.—Much has been said and written in reference to the use of dilute sulphuric acid in diarrhœa ;* and its use has certainly been attended with benefit, although not to the extent we were led to suppose. Both dilute sulphuric acid, and dilute nitric acid, are of value after the more severe symptoms have passed off ; they act at first possibly by checking chemical and fermentative changes, and afterwards as tonics to the relaxed mucous membrane. Combined with slightly astringent and mucilaginous tonics, as with cusparia and simaruba, they are of great service in some cases.

When there is much pain, we may associate narcotics with other remedies before mentioned. Chloric-ether, also, sometimes affords great relief, when administered with mucilaginous remedies ; so also the tincture of hcnbane ; in other cases, simple carminative medicines are sufficient to relieve the pain, as ginger, cardamoms, &c., especially where the diarrhœa is associated with flatulent colic.

* Dr. W. Griffith and Dr. Fuller, "On the Use of Sulphuric Acid in Diarrhœa ;" 'Lancet' and 'Medical Times and Gazette.'

In the colliquative diarrhœa of weaned children, Dr. I. F. Weisse has strongly advocated the administration of raw meat, scraped and reduced to a pulp; but our experience in this remedy does not lead us to recommend its use.

Leeches.—The application of leeches to the anus is a remedy which greatly relieves inflammatory congestion of the mucous membrane of the large intestine, but it is one which we should scarcely recommend, unless the disease assume a severe and dysenteric character.

Suppositories, composed of the compound soap pill, are often of great service when there is distressing tenesmus which disturbs the rest of the patient; and when it is undesirable to administer an opiate by the mouth, and inconvenient to use an enema.

In chronic mucous discharge from the bowels, we must first seek to rectify the disease of the liver, if such exist, by mild alteratives; by taraxacum, and by nitro-muriatic acid. These remedies, also, assist in relieving the chronic congestion and inflammation of the intestine, and are more effective than astringents. It is well, however, to be assured that no polypoid growth, nor disease of the rectum and sigmoid flexure, is setting up the disease.

If astringents be required in these instances, the oxide and nitrate of silver, sulphate of copper with opium, or the vegetable astringents just mentioned, may be used; and as enemata, the solution of nitrate of silver (gr. x—xv to Oj.*), the infusion of quassia, the decoction of oak bark, and the decoction of poppies, may be employed with advantage.

In the treatment of choleraic diarrhœa, rest in the recumbent position, warmth to the abdomen and the feet, and gentle friction, if muscular spasm distress the patient, are valuable remedies. A full dose of chalk and opium with catechu and aromatic spirit of ammonia should be given, and repeated in two or three hours, if necessary. Demulcent nutriment, as mutton broth and arrowroot, may be allowed; and if vomiting supervene, ice or cold water will be beneficial. Dilute sulphuric acid has been sometimes used with great advantage, and by some

* Trousseau.

calomel has been freely given in these cases, especially when vomiting has come on. When the collapse of true cholera has attacked the patient, general experience does not favour the free use of either opium or brandy; but to enter fully into the treatment of cholera is foreign to our purpose.

CASE CLXI.—*Inanition. Diarrhœa.*—John M—, æt. 26, was admitted into Guy's Hospital December 16th, 1856, and died December 21st. He had been a sailor, and stated that he had had dysentery, but this was not satisfactorily ascertained on account of his prostrate condition. It appeared that he had been on board an American vessel from China to Liverpool, and arrived at the latter place on December 6th; he then came up to London. He informed the nurse that there had been a mutiny on board, and that he had been put in irons in the hold. He was in the most emaciated state; the voice was scarcely perceptible; the pulse was exceedingly compressible, and the tongue and mouth presented yellowish-white aphthous patches; he had no vomiting, but the stools escaped from him, and were white and very offensive; the respiration was easy, and the mind perfectly conscious. Milk was ordered. The following day he was better, but sank on the third day after admission, and was sensible till nearly the last.

Inspection, December 22, 1856. There were ecchymoses on both thighs, and old cicatrices on the wrist and leg. The brain was less firm than normal, and the lungs were collapsed and healthy. The heart was small. The liver was healthy. The gall-bladder was not distended, and the spleen and kidneys were healthy. The stomach presented gastric solution at the cardiac portion. The small intestines were healthy. The large intestine was throughout of a gray colour, and was filled with dry, white fæces. At the root of the mesentery were several white strumous masses in the glands, but it could not be found that the thoracic duct was obstructed. The urinary bladder was distended.

This case presents us with a well-marked instance of a man dying from the effect of starvation. The diarrhœa was probably the result of want of nourishment, of good air, and of light, &c.; so that the supply and conditions necessary for reparation having been cut off, the whole body wasted, and the spark of life gradually expired.

CASE CLXII.—*Chronic Diarrhœa. Hysteria. Great relief from Tincture of Iron.*—Georgiana B—, æt. 40, a single woman, who had resided in the Commercial Road, and had supported herself by her needle, applied at Guy's Hospital May 23rd, 1860, and was admitted under my care. She stated that for two or three years she had suffered from ulceration of the womb, and the speculum had been repeatedly passed. During eighteen months she

had been affected with diarrhœa, and when she had mental anxiety the disease increased in severity, but it had been partially relieved by chalk medicine. Menstruation was regular as to time; the urine was passed well, but afterwards she suffered from "bearing down" pain. The slightest exertion produced perspiration. She was a tall woman, extremely nervous, the eyes sunken, the countenance dejected, the hair turning gray; the skin was hot, but moist; the tongue was whitish; the pulse was 90, small and hard; respiration 22. The heart and lungs were normal. She complained of great pain in the abdomen, on the right side, below the liver, in the region of the ascending colon; there was tympanitic distension in the same region; the bowels were opened six times in twenty-four hours, but there was no evidence that blood had been passed. There was pain on passing water. She had not taken meat during several months. The compound krameria mixture and the compound logwood mixture of Guy's were tried with very little effect; starch injections afforded slight relief; wine also was given. Dr. Oldham could not detect any uterine disease. Nitrate of silver, gr. $\frac{1}{4}$, and opium, gr. $\frac{1}{2}$, were given three times a day. On June 23rd the bowels were not quite so loose; the patient had tried to take a little meat, but it produced pain at the stomach; a starch injection, with gr. xv of borax, also afforded slight relief. The tincture of iron was then given, and after it had been continued for three weeks, although the diarrhœa had not ceased, the patient felt stronger and much improved in health. The same treatment was continued for six weeks more; the countenance had then become less pallid, the pulse more firm; she was able to take meat diet. The diarrhœa had in a great measure subsided. She left the hospital convalescent, stating that she had not been so well for eight years.

This case appeared to be one of passive mucous diarrhœa in a very hysterical subject, and the uterine irritation had tended to perpetuate the disease. Astringents were less efficacious than preparations of steel, which diminished the nervous irritability and gave tone and strength to the whole system. The regulated and more generous diet which the patient had must not be overlooked; and by persuasive measures she was induced to take a meat diet, which lessened the fluid contents of the colon, and thereby increased the consistence of the alvine discharges.

CHAPTER X.

ON COLITIS AND DYSENTERY.

Numerous authors, as Sydenham, Annesley, Parkes, Ballinghall, Sir J. McGrigor, and Morehead, have described the terrible forms of this disease as they are manifested in tropical climates, and in military campaigns, and as formerly seen amongst ourselves. In our own country, however, it has very much diminished in severity and in frequency, so that acute dysentery is rarely seen amongst us, unless contracted in foreign climes, and then brought to our shores. Still true dysentery occurs more commonly than some of very great experience will admit; sometimes it is the only disease from which the patient suffers, and is quickly fatal; or it is found in association with a general inflammatory condition of other mucous membranes; or lastly, it causes the sudden termination, or it aggravates varied forms of chronic disease.

Particular localities and periods of the year cause the manifestation of this disease, and its complication with others in a very marked degree. This fact is shown by the observations of Dr. Latham and Dr. Baly at the Millbank prison; and the greater humidity of the atmosphere is probably the reason of the more frequent occurrence of dysentery in the hospitals in Southwark, than elsewhere in London. The complication of dysentery with other diseases is a very important consideration in their prognosis and treatment: thus, incipient phthisis may become altogether hopeless, and in a very short time fatal, not from the severity of the pulmonary affection, nor from strumous disease of the intestines, but from acute inflammation of the mucous membrane of the colon.

Abercrombie defined diarrhoea as purging, arising from irrita-

ting substances in the canal, and from secretions poured into it; and dysentery, as acute inflammation originating in the mucous membrane of the large intestine. This distinction is probably, to a considerable extent, correct; but some forms of disease usually considered as diarrhœa, arise from catarrhal inflammation of the colon and small intestine, and after death may present scarcely any trace of abnormal change. Dysentery is generally limited to the colon, and when severe the disease rapidly passes into ulceration and sloughing, unless from its extent, or the previous condition of the patient, it prove fatal at an anterior stage, as in several of the fatal cases recorded in this chapter, which terminated before extensive ulceration had taken place.

Dr. Lyon, in his Crimean report, has divided dysentery into two forms, the exudative and the follicular. The former is, however, probably the earlier stage, or that preceding ulceration and sloughing. The cases which have come under my own observation may be divided, practically, into three classes:—

1. Those instances in which acute disease of the colon is the primary disease, and sometimes terminates fatally in a short time.

2. Those cases in which dysentery is associated with inflammation of other membranes and organs, as when bronchitis, laryngitis, and pneumonia arise at the same time, and are produced apparently by the same cause as the disease of the intestine; in some instances the disease is closely allied to pyæmia.

3. Those cases in which inflammation of the colon has hastened the fatal termination of other more chronic disease.

The dysenteric process is well described by Rokitansky, who divides it into four stages, and considers that it consists in inflammation of the mucous membrane of the colon, terminating in severe cases in sphacelus. Dr. Parkes believes that, in true dysentery, ulceration is always present, and attaches great importance to the affection of the glands; whilst Dr. Baly describes the process as sloughing, rather than ulceration. Are we then to look upon inflammation of the colon, in which there is no destruction of the mucous membrane, as true dysentery? Most persons will acknowledge, that death may take place prior to the ulceration or sloughing, although we rarely, if ever, find the

mucous membrane entire; it is probable that the diseased condition is closely allied to that of the pharynx in diphtherite; and that in severe cases the membrane rapidly sloughs, without antecedent ulceration.

In the earliest stage of dysentery, the mucous membrane becomes injected, œdematous and thickened; the mucus is scanty, and the fæces become adherent; this condition may be universal in the colon, or limited to the rectum, to the sigmoid flexure, or to the cæcum. The solitary glands will be found to be distinct, prominent, and enlarged.

The secretion from the membrane then becomes further changed, and a thin exudation, consisting of epithelium with a considerable quantity of granular amorphous matter, coats the intestine. This exudation is found in patches, or in lines, or it is spread generally, upon the surface. It has been described as dipping into the follicles; an appearance which I have myself observed, and it may also be seen closely incorporated with the surface of the membrane, so that it can only be separated by considerable violence. The colour of the exudation is generally a greenish yellow; but it varies somewhat according to the character of the fæces. On scraping off the effusion from the surface, the membrane beneath is found intensely congested, and often superficially ulcerated; or, there may be merely minute circular patches of ulceration, with portions of the false membrane adhering at that part. This tendency to ulcerate, or to slough, resembles diphtheritic disease in the pharynx and nares; but the character of the false membrane in the throat is sometimes more fibrinous than that developed in the intestine. The muscular coat of the colon appears thickened in this stage of dysentery, probably because it is contracted; and the submucous cellular tissue is often whitish and distinct from inflammatory œdema. Dr. Baly, whilst describing this epithelial degeneration, states, that in most cases these minute adherent coverings on the surface of superficial erosions, or small ulcers, consist of thin sloughs of the mucous membrane. He believes that, in all cases, the destruction of the mucous membrane consists in a process of mortification and sloughing, and not of simple ulceration; and that the disease commences in the solitary glands of the intestines. Other parts, however, beside the solitary glands, are

found to be diseased ; but whether primarily or by extension, is matter of opinion. Many instances of *diarrhœa* are observed, in which, after death, the solitary glands were found enlarged, or minute points of ulceration were presented ; the whole colon may be studded over with minute ulcers, arising apparently in the glands, as is well shown in a specimen in the Guy's Museum, to which reference has before been made. Dr. Baly would probably consider these to be instances of the dysenteric process in its mildest form, and that in other instances, previously alluded to, more acute changes had spread from the glands to the general surface of the membrane.

Dr. Morehead has less frequently observed diphtheritic membrane effused in dysentery, and believes that the mucous follicles are more frequently affected than the solitary glands*.

In the third stage, we find ulceration, sometimes merely as minute circular ulcers, but generally of a more extensive character ; the ulcers are often oval in form, and placed in the transverse axis of the intestine ; and their edges are raised and injected, irregular and undermined ; and their base is formed by the cellular or muscular coats. These ulcerations gradually extend so as to coalesce, till at last nearly the whole of the mucous surface is destroyed, except here and there prominent isolated portions, which become intensely congested, and resemble polypoid growths. In severe cases the whole colon, from the cæcum to the rectum, is in this condition, or greater spaces intervene between the ulcers, which are only found in the rectum, or in the sigmoid flexure, or in the cæcum ; but generally the rectum and the sigmoid flexure are the parts most severely implicated. It sometimes happens, that the ulceration extends through the muscular and the peritoneal coat, leading to fatal peritonitis from perforation, but this is a rare occurrence ; or, the coats of the intestine become sinuous abscesses, so that on dividing a prominent portion of mucous membrane between two ulcers, several drachms of pus escape. This extensive suppuration is very different from the small local collections of pus, which sometimes form in the substance of the mucous membrane after follicular and glandular inflammation,

* Morehead 'On Diseases of India.'

where small eminences about half an inch in diameter are observed, covered by thin layers of mucous membrane. The latter is a less general and less severe form of inflammation of the colon, but one which I have several times observed.

The liver is generally found to be congested, and the gall-bladder to be partially distended with bile, either inspissated or orange-coloured mucus. The mesenteric glands are often somewhat congested; and in some instances the ileum and the jejunum are both diseased, becoming intensely injected, partially ulcerated, and covered with diphtheritic membrane; Peyer's glands are also found to be congested and enlarged, although never raised and infiltrated with deposit, as in typhoid fever.

Dr. Morehead gives an instance of faecal abscess, in the right iliac fossa, from perforation of the caecum in dysentery.

If the acute symptoms have subsided, the injection is less deep in colour, and often becomes grey; the edges of the ulcers assume a rounded and less prominent aspect; their surface has a smooth and fibrous appearance; ulcerative action has been checked, and cicatrization has commenced. The healing process may go on, so that the cicatrix has an irregular, puckered appearance. The base of the cicatrix is formed by fibro-cellular tissue, but the gland structure is not reproduced. The contraction of the cicatrix sometimes produces considerable constriction of the intestine, and occasionally tends to fatal obstruction. Very frequently above the cicatrix all the coats become hypertrophied, showing that there has been much impediment.

Dr. Wilks has informed me of a case in which the cicatrix presented a growth at its margin, evidently of a carcinomatous character, indicating a greater tendency to heterologous deposit in the new tissue.

In a fourth stage of dysentery the mucous membrane presents a grey, ashy appearance, and considerable portions of it constitute ragged and semi-detached sloughs.

Dysentery often terminates in complete recovery; at other times death quickly supervenes from the severity of the disease; or, thirdly, it passes into a chronic state, which may continue with intermissions for several years.

Other *sequelæ* of dysentery are—perforation of the intestine

and fatal peritonitis; faecal abscess; gradually increasing exhaustion from the destruction of the mucous membrane, and continued purging; constipation, arising from the contraction of cicatrices, leading to a very troublesome and irregular condition of the bowels, and sometimes to fatal obstruction; pyæmia and suppuration in the substance of the liver, from the absorption of pus, as described by Dr. Budd, in his work on Diseases of the Liver. This last result I have only once observed at Guy's, in simple English dysentery, which is nearly in accordance with the experience of Dr. Baly, at the Millbank Penitentiary, and shows that whilst the disease may be the same in its general character and pathology with tropical dysentery, there is some modifying cause. Dr. Mayne, in a very interesting article in the Dublin 'Quarterly Review,' also mentions a state resembling phlegmatia dolens as the occasional sequence of dysentery, and speaks of spontaneous salivation as sometimes supervening after the disease.

Symptoms.—A sensation of coldness in the loins, chilliness, or actual rigor, is followed by a loose evacuation from the bowels; this alvine discharge is repeated, and the evacuations become scanty, and are often accompanied with tenesmus, or a forcing sensation as if the intestine retained its faecal contents. With this, there may be slight pain or soreness in the iliac region or position of the transverse colon, and even severe griping; febrile disturbance supervenes, and the tongue has a whitish fur, or it is injected at the tip and edges, but the latter symptoms are often absent. In mild cases the energies and mental activity are unimpaired, but the face becomes pallid, and the strength is not equal to accustomed duty.

This condition may continue for several days, the patient becoming more feeble, the motions watery, or they contain mucus and blood, and scybala are passed with pain. The countenance becomes haggard, and expressive of distress, the skin clammy, and the pulse compressible; the abdomen is collapsed, and tolerant of pressure; the pain is paroxysmal, and the colic occasionally very severe, and vomiting sometimes supervenes.

If the disease continue unchecked, the strength fails, the pulse is rapid and compressible, and the eyes are sunken; the tongue is brown, or it is dry and cracked; the motions are

passed involuntarily, and often are of a greenish colour, or they resemble the washing of meat; the lower extremities are cold, and the hands and face are covered with a clammy sweat; occasional cramps, with hiccup and subsultus tendinum, come on; the patient is sensible, but speaks in a feeble tone of voice, and at last dies, more suddenly, perhaps, than those around him had anticipated.

These symptoms may only extend over a very few days or hours, and sometimes are accompanied with much febrile disturbance, with tenderness of the abdomen, and with a furred and brown, or red, dry and glazed tongue. If the severity of the disease abate, the bowels become more composed, and the patient rallies; or the diseased condition returns with greater severity; for the first exhaustion is scarcely recovered from before the strength is still further reduced, and in this way the malady may extend over weeks, or months. The patient then has a peculiar and characteristic appearance; he is much emaciated; he has a sallow complexion; his mind is active, but physical strength fails gradually, till at last he is obliged entirely to take to his bed from a sense of utter prostration.

In the dysentery of the tropics, or the equally severe disease observed in the hardships of war and after famine, the prostration is more rapid, the disease more quickly fatal, and in some instances has been accompanied with violent delirium.

In reviewing the symptoms of inflammation of the colon in its severe forms, we have been struck with the occasional absence of febrile symptoms, as indicated by a hot skin, furred tongue, or excited pulse. The tongue, unless it be itself affected with local disease, indicates the condition of the whole nutritive function, rather than that of any isolated portions of the alimentary canal.

If the peritoneum become inflamed, or if the extension of disease to the muscular coat produce irregular muscular contraction, pain often of a severe, griping, and of a paroxysmal character is the result. The amount of pain appears to be greater in disease of the small than of the large intestine, perhaps on account of its greater mobility.

The character of the evacuations deserves particular attention. They may consist of fluid *fæces* containing *scybalæ*, or only of

and fatal peritonitis; fæcal abscess; gradually increasing exhaustion from the destruction of the mucous membrane, and continued purging; constipation, arising from the contraction of cicatrices, leading to a very troublesome and irregular condition of the bowels, and sometimes to fatal obstruction; pyæmia and suppuration in the substance of the liver, from the absorption of pus, as described by Dr. Budd, in his work on Diseases of the Liver. This last result I have only once observed at Guy's, in simple English dysentery, which is nearly in accordance with the experience of Dr. Baly, at the Millbank Penitentiary, and shows that whilst the disease may be the same in its general character and pathology with tropical dysentery, there is some modifying cause. Dr. Mayne, in a very interesting article in the Dublin 'Quarterly Review,' also mentions a state resembling phlegmatia dolens as the occasional sequence of dysentery, and speaks of spontaneous salivation as sometimes supervening after the disease.

Symptoms.—A sensation of coldness in the loins, chilliness, or actual rigor, is followed by a loose evacuation from the bowels; this alvine discharge is repeated, and the evacuations become scanty, and are often accompanied with tenesmus, or a forcing sensation as if the intestine retained its fæcal contents. With this, there may be slight pain or soreness in the iliac region or position of the transverse colon, and even severe griping; febrile disturbance supervenes, and the tongue has a whitish fur, or it is injected at the tip and edges, but the latter symptoms are often absent. In mild cases the energies and mental activity are unimpaired, but the face becomes pallid, and the strength is not equal to accustomed duty.

This condition may continue for several days, the patient becoming more feeble, the motions watery, or they contain mucus and blood, and scybala are passed with pain. The countenance becomes haggard, and expressive of distress, the skin clammy, and the pulse compressible; the abdomen is collapsed, and tolerant of pressure; the pain is paroxysmal, and the colic occasionally very severe, and vomiting sometimes supervenes.

If the disease continue unchecked, the strength fails, the pulse is rapid and compressible, and the eyes are sunken; the tongue is brown, or it is dry and cracked; the motions are

passed involuntarily, and often are of a greenish colour, or they resemble the washing of meat; the lower extremities are cold, and the hands and face are covered with a clammy sweat; occasional cramps, with hiccup and subsultus tendinum, come on; the patient is sensible, but speaks in a feeble tone of voice, and at last dies, more suddenly, perhaps, than those around him had anticipated.

These symptoms may only extend over a very few days or hours, and sometimes are accompanied with much febrile disturbance, with tenderness of the abdomen, and with a furred and brown, or red, dry and glazed tongue. If the severity of the disease abate, the bowels become more composed, and the patient rallies; or the diseased condition returns with greater severity; for the first exhaustion is scarcely recovered from before the strength is still further reduced, and in this way the malady may extend over weeks, or months. The patient then has a peculiar and characteristic appearance; he is much emaciated; he has a sallow complexion; his mind is active, but physical strength fails gradually, till at last he is obliged entirely to take to his bed from a sense of utter prostration.

In the dysentery of the tropics, or the equally severe disease observed in the hardships of war and after famine, the prostration is more rapid, the disease more quickly fatal, and in some instances has been accompanied with violent delirium.

In reviewing the symptoms of inflammation of the colon in its severe forms, we have been struck with the occasional absence of febrile symptoms, as indicated by a hot skin, furred tongue, or excited pulse. The tongue, unless it be itself affected with local disease, indicates the condition of the whole nutritive function, rather than that of any isolated portions of the alimentary canal.

If the peritoneum become inflamed, or if the extension of disease to the muscular coat produce irregular muscular contraction, pain often of a severe, griping, and of a paroxysmal character is the result. The amount of pain appears to be greater in disease of the small than of the large intestine, perhaps on account of its greater mobility.

The character of the evacuations deserves particular attention. They may consist of fluid *fæces* containing *scybalæ*, or only of

Dysentery often prevails after scarcity of food, and terrible manifestations of this fact have occurred in Ireland. The observations there made by some of its most accomplished physicians, and the valuable reports they have furnished, give some of the most faithful descriptions which we possess of dysentery in its worst forms. I refer to the labours of Drs. Mayne and Malcolm, Lalor, Harty, and Young, &c.

In the active life of a soldier, especially in the tropics and in marshy districts, these causes operate in all their intensity; and the disease thus manifested is more fatal than the field of battle itself. Such has been the testimony of nearly all the writers on military surgery and medicine.

Inflammation of the colon is, however, set up without any miasmatic influence; thus, from poisons, inflammation of the stomach and the colon may be produced without the intervening small intestine being affected; so also irritating ingesta and excretions appear in some instances, to be the sole cause of the complaint. Cold drinks in excess during summer heat also tend to produce dysentery.

Many believe in the contagious character of the disease. They have found it extend from one to another, to attendants, or from house to house, &c.; but it must be remembered, 1st, that in many of these cases there is a general pervading influence, miasmatic or otherwise; 2nd, that the effluvia from the dysenteric discharges are exceedingly offensive; 3rd, that animal effluvia are of themselves sufficient to induce the complaint; 4th, that the depressing effect of night-watching, and of witnessing the rapidly fatal termination of the disease, tends also to induce the complaint.

The occasional similarity of dysentery to typhoid fever, or the diseases taking place at the same time and in the same patient, do not warrant us in considering it as of the same character or type of disease as typhoid fever, namely, a blood disease associated with special changes in the glands of the intestine. There are few diseases that have not more or less of a constitutional origin, and dysentery is doubtless of that character; but in some there is evidence of poison contaminating the blood, leading to a special train of symptoms, as in smallpox, or in pyæmia. There does not appear to be warrant for supposing

that such is the case in dysentery, more than in cancrum oris, ulcerative stomatitis, and diphtheritic inflammation of the pharynx. Some cases of dysentery, as of pneumonia, doubtless arise from a diseased condition of blood, whilst others are of a different character; thus, pneumonia may be produced by direct exposure to cold; or again, the blood may be in such a condition from pyæmia and other septic changes, that pneumonia is induced without any other exciting cause. Some of the cases of dysentery here recorded appeared to be produced by irritation extending from the rectum; others from direct irritation caused by improper food; a third class from a morbid condition of the whole circulating fluid.

As far as our observations extend, we are disposed to agree with the statements of Dr. Harty, in the 'Dublin Quarterly Review'—1st, that genuine and simple dysentery is unattended with idiopathic fever, and is never of itself contagious; 2nd, that every form of the disease when epidemic is a combination of the simple dysentery either with intermittent, remittent, or continued fever; and 3rd, that the combination of dysentery with fever is alone contagious.

The *Prognosis* of dysentery is unfavorable when purging of blood, and thin serous offensive discharge like the washing of meat continue. When there is involuntary discharge from the bowels, with great tenderness of the abdomen, vomiting, red and glazed tongue, typhoid symptoms, irregular pulse, refusal of food, restlessness, and great prostration of strength, or when these symptoms have continued for a considerable time, and one relapse has followed another, without the patient in the interval regaining strength, but retaining the same sallow and haggard expression; when, also, there are cerebral symptoms, as coma, delirium, convulsions, we judge unfavorably. On the contrary, when, on the removal of the causes of the disease, the evacuations assume a healthy appearance, and contain bile, when the pulse remains firm, and the prostration becomes less, when the tenesmus ceases, and the patient gains strength, we may give a cheering prognosis.

Diagnosis.—There are several conditions which may, unless due care be used, be mistaken for dysentery.

1. The *discharge of blood from hæmorrhoids*, accompanied

with diarrhœa and prostration. Ordinary care alone is necessary to guard against this error.

2. *Disease of the rectum.*—The lower part of the rectum sometimes become ulcerated, and leads to a discharge of mucus and blood with tenesmus, and with great anxiety and sympathetic nervous disturbance to the patient. It will be found in many of these cases, that a small quantity of mucus or pus escapes from the rectum at irregular intervals. In organic constriction of the rectum the bowels are sometimes in a loose condition.

3. *Polypus in the rectum* will produce like symptoms. Many of these growths can be detected by examination per rectum; others, however, are beyond the reach of the finger, and we can then only decide by the difference in other symptoms. In these instances healthy fæces are passed, but coated or followed by mucus or pus, and the constitutional disturbance is less severe.

4. *Fibro-cellular ulceration.*—By this we mean a condition of rectum which is not cancerous, but in which the mucous membrane is ulcerated and the coats of the intestine are converted into dense fibrous tissue.

5. Some of the forms of *diarrhœa* resemble the earlier symptoms of dysentery, but they are generally distinguished by the absence of blood with mucus. In some cases, however, where there is disease of the mesenteric glands, with persistent diarrhœa, and prostration of strength, we are apt to believe that ulceration of the colon exists; the stools however are different; there is less of mucus, and no blood can be detected even by the microscopical examination of the evacuation.

6. A feeble condition of the *sphincter ani*, and of the muscular coat of the intestine, leading to the involuntary discharge of fluid fæces, might induce the belief in an abnormal condition of the mucous membrane of the colon.

Treatment.—Amongst the most important points for consideration in the treatment of dysentery are the removal of all exciting causes, and of the general conditions in which the disease has had its origin. The patient should be clothed in flannel, and the temperature of the body maintained, dampness and cold carefully warded off, improper diet avoided, and, if possible, good and pure air inhaled.

Demulcents administered by the mouth have many feet over

which to pass before reaching the large intestine, but they act very beneficially by rendering the excreta poured into the colon less irritating, and thus soothing the diseased membrane. In this way arrowroot, linseed tea, mutton and veal broth, milk with suet, rice milk, tapioca, &c., may be both grateful to the patient, and act remedially, as before mentioned; fluid drinks should not be taken in great excess; water alone has been used, but it is inferior to demulcent drinks. By these simple means many attacks may be relieved, and the patients speedily recover.

The diet calls for much attention, equally in the early stages and milder forms, as in those of a more severe character; but in the former it is often more difficult to persuade the patient to follow our directions; being free from pain, the tongue clean, the appetite craving, he does not at once see the importance of using proper care. Any imprudence perpetuates and aggravates the disease. Malt liquors and spirits should be abstained from. Meat in a solid form is generally better avoided in the acute disease, but when taken should be in an easily assimilable form, and neither richly dressed nor highly seasoned.

In protracted cases of dysentery, it is necessary to take animal food, and it is very desirable in other ways to sustain the strength of the patient as far as possible. For a short time, at least, it is well to omit vegetables and fruits, especially when uncooked; but oranges, grapes, &c., by supplying to the system that which is necessary for the maintenance of sound action, without much indigestible product, may sometimes be taken with great benefit in chronic dysentery, although apples, and stone fruit, melons, salads, &c., do harm.

Rest is important, and in very many cases the patient is too ill to leave the recumbent posture; in the chronic disease, also, fatiguing muscular exertion and horse exercise must be avoided.

Other remedies in most frequent use, have been opium, astringents, both metallic and vegetable, ipecacuhana, mercurials, depletion, enemata of various kinds, &c.

Opium is of almost universal application, but cannot be given *indiscriminately*, either alone or in combination; it acts partly by its astringent properties, and partly as a narcotic, by its secondary influence upon the inflamed mucous membrane, through the sympathetic system of nerves. It is often ad-

ministered alone, but more frequently in combination with ipecacuanha, as in Dover's powder; or with astringents, as in the compound krameria mixture, or compound logwood mixture of Guy's; or in combination with mercurials, as calomel, and grey powder; and lastly, in enemata. When the excreta are depraved, and cause irritation of the intestine by their presence, no beneficial result is attained by seeking to check their removal by astringents.

Mercurials.—Calomel has been very largely and notoriously administered in dysentery. In India 3j doses are described by Annesley as productive of essential benefit; others have now given up its use altogether, in the severe dysenteries of the East. Its administration, in more moderate doses, is often advantageous. In the earliest stages, where the motions are of unhealthy character, pale, or containing hard scybalous masses, it may be well to give a dose of calomel, or of grey powder, followed by castor oil, guarded by opium; or the opium may be combined with the mercurial, as calomel with opium, or grey powder with Dover's powder; in this way offending substances may be removed, and acrid excretions rectified and corrected; but it is, we believe, unwise and uncalled-for, to persist in the use of mercurials. We have no facts to show that the inflammation of the mucous membrane is diminished by its action, but rather that it is increased, and ulceration accelerated, although the abdominal glands may be stimulated to a more healthy action.

Purgatives.—Almost the same may be said of these remedies, as of mercury. In the earlier stages of dysentery, and dysenteric diarrhoea, when oftentimes, irritating and crude materials are retained, and equally irritating secretions are poured out, purgatives are of great value; castor oil, and linseed oil, with tincture of rhubarb, and with small doses of tincture of opium, are old and valuable remedies for this purpose. They may be repeated several times during the day, with manifest improvement of the alvine evacuations, and relief to the tenesmus and the pain. This administration of a laxative draught, followed by a full dose of Dover's powder, or chalk with opium is the best treatment for ordinary dysentery at its onset. Violent purgatives are, however, neither beneficial nor warrantable.

Ipecacuanha was introduced into Europe as a remedy for dysentery, nearly 170 years ago, and has been used with success since that time; large doses have been given without producing vomiting, as by Sir J. Pringle, in gr. v, to ʒj, doses, and by Mr. Twining, with extract of gentian; but much smaller doses are equally effective. *Ipecacuanha* appears to have the same beneficial action in inflammation of the alimentary as of the respiratory mucous membrane, in relieving the congestion, and in restoring healthy secretion. It has been given in various combinations, frequently with opium, and with vegetable astringents, as kino, krameria, logwood, catechu, &c.

Astringents are of greater value in chronic than in acute dysentery; sulphate of copper, acetate of lead, oxide and nitrate of silver, nitrate of bismuth, sulphate and perntrate of iron, have all been used; several of those named act by their direct astringent property, others become absorbed and are sedatives to the mucous membrane; some of these remedies have been used in combination with opium, as copper, lead; others, with considerable success, as injections, and in that way applied directly to the diseased surface; thus nitrate of silver has been used as an enema by Trousseau, in the proportion of gr. x to xij to a half pint of distilled water; so also, borax in about ʒj doses; in some cases I have used charcoal, ʒij, with Oss of thin. gruel, with evident relief.

Vegetable astringents are valuable remedies in the chronic disease, and even in the acute stage after the removal of irritating excreta and scybala, as krameria, kino, logwood, and catechu, tormentilla, and simaruba; they are of especial service when given in combination, as in the krameria and logwood mixture of Guy's. *Cusparia* is a more stimulating astringent and tonic, and is often of great benefit after the more active symptoms have subsided.

Tannin and gallic acid have been used, but I have been disappointed in their action. The rind of the pomegranate root is a favorite remedy with some, but is a less effective astringent than others just mentioned. I have tried the Indian bael in many instances of chronic dysentery, but I have not seen it followed by the benefits which were anticipated from the high encomiums given to it by practitioners in India. The solution

of potash, as well as the bicarbonate of potash, and the corresponding preparations of soda, in the acute disease diminish intestinal irritation; they are best given, however, in combination with opium, or henbane and astringents.

Earthy astringents.—Chalk, alone or combined with opium and astringents, is one of the ordinary remedies at the outset of dysenteric diarrhœa, and is frequently sufficient to check the purging; but in many cases no form of astringent nor sedative will serve to restrain the purging, whilst irritating excreta and other substances are retained; and in others, so great is the extent of the disease, that it is futile to attempt to check it by a grain or two of medicine which acts locally, whilst the disease extends over many feet of intestine. By demulcents, by removing the cause of the disease, by rest, by correcting the general disturbance, and by sustaining the patient during the degenerative changes which are in operation, we must seek to shorten the morbid process and restore to health.

Mineral acids.—After the urgency of the symptoms have subsided, mineral acids, both sulphuric and nitric, appear to act beneficially on the mucous membrane in restoring healthy vigour. I have not found the benefit from dilute sulphuric acid that some practitioners have observed; still it is a remedy of value, and well deserves a trial; and in chronic dysentery the nitric and nitro-hydrochloric acids are often of service when combined with opium, and with the milder vegetable astringents.

Quinine.—The combination of quinine with Dover's powder, or with mineral acids and astringents, is productive of considerable relief in those cases of chronic dysentery which have been perpetuated and aggravated by residence in miasmatic districts.

Depletion.—I have never observed a case of dysentery where venesection could have been tried with probable success; but local depletion is often of great service, either applied to the anterior surface of the abdomen, or to the anus. The latter more directly acts on the intestine, on account of the connexion of the inferior hæmorrhoidal with the inferior mesenteric veins; ten or fifteen leeches may thus be applied with relief to the pain and to the violent tenesmus, &c.

Warm enemata often afford very great relief by washing out

the lower part of the rectum, and by soothing the inflamed membrane; in this way gruel, thin starch, and barley water may be used with much relief; but their efficiency is increased by the addition of tincture of opium. Borax and nitrate of silver are used, as we have before mentioned, and sometimes charcoal; these agents and vegetable astringents, as oak bark, have been more especially tried in chronic dysentery.

Suppositories, composed of opiates or anodynes, afford much relief to the tenesmus, but are less effective than enemata.

In the constipation which follows the cicatrization of dysenteric ulcers, much relief is afforded by the use of sedative with aperient medicine, as by the colocynth and henbane of the Edinburgh Pharmacopœia with ipecacuanha, and by soap with rhubarb and capsicum.

A change to more genial climate is often productive of the most beneficial result; this applies especially to residents in India, and to those who have contracted dysentery in our own country in damp and in miasmatic districts.

The following fatal cases of dysentery were, with one or two exceptions, produced in our own country, and will illustrate the causes of death in this disease:

Harriet S—, æt. 28 years. Acute inflammation of the whole of the colon.

Elizabeth H—, æt. 7. Acute inflammation of the whole of the colon.

Ann H—, æt. 60. Inflammation of the colon after hernia.

Edward B—, æt. 39. Inflammation of colon; perforation; fæcal abscess.

James T—, æt. 52. Inflammation of colon; perforation; submucous suppuration; pus in the vena portæ; inflammation of the liver.

Sarah W—, æt. 34. Dysentery; perforation of the colon.

Thomas R—, æt. 31. Chronic bronchitic phthisis; cirrhosis; contracted abscess of liver; chronic dysentery; and chronic peritonitis.

Thomas D—, æt. 25. Chronic dysentery; hepatic abscess; pyæmia; abscess in the brain.

John I—, æt. 32. Dysentery; abscess of liver; perforation of the diaphragm; empyema.

Evan T—, æt. 33. Cicatrization and contraction of the rectum, and sigmoid flexure, after dysentery.

Mr. W—, about 35. Dysentery; cicatrization; contraction; abscess near the crest of the ileum.

These cases indicate—

I. That dysentery of a most severe form arises in our own country, and is not of unfrequent occurrence.

II. That the cause of death in some is the extent and severity of the affection.

III. That others die from perforation and faecal abscess.

IV. That pyæmia follows some of the worst forms of English dysentery.

V. That the constriction of the intestine sometimes leads to abscess in the parietes and to artificial anus.

VI. That in the worst cases astringents and opiates are ineffective.

VII. That injections and demulcent remedies afford considerable relief, and in mild cases will alone be sufficient; but are inferior in their efficacy to astringents and opium.

VIII. That rest even in mild cases is desirable.

Many cases have presented themselves, in my own practice and spheres of observation, where these means checked the purging, and restored to health.

IX. That as far as can be judged, mercurial preparations, if continued, would have been injurious.

CASE CLXIII.—*Inflammation of Colon. Aphtha or Muguet of the Pharynx.*—Harriet S—, æt. 28, was admitted July 18th, 1855, under my care.

She was a pale, thin woman, who had resided in the Borough, the wife of a butcher, and she had been out of health for six months, with occasional diarrhoea; but during seven weeks she had become much worse. Her illness came on after exposure to cold on entering a new house; she had great lassitude, weariness, and severe pain in the limbs. These symptoms increased, and she was admitted into St. Thomas's Hospital, and whilst there she passed blood and mucus from the bowels. She stated that her family were healthy, but that she had previously had an attack of inflammation of the bowels. On admission she had the appearance of a patient suffering from acute pneumonic phthisis; the lips and nostrils were thin; the cheeks were sunken; the skin was hot and clammy; the conjunctiva and skin generally were of a very slight jaundiced colour. Along the gums was a pale red line; the tongue was coated with a yellow fur, dry and brown, and at its centre there were several

apthous patches; the throat was dry and injected, and she complained of difficulty in swallowing, and it was painful to her to speak much. The pulse was 112, soft and full. The chest was well developed; the respiration was coarse at the apices, but otherwise normal. In the abdomen a large mass of the form of the liver could be felt, extending nearly to the crest of the ileum, and across the umbilical region. The abdomen was flaccid, but slightly tender on manipulating the right hypochondriac region. The motions were passed involuntarily, and were slimy and of a green colour; three or four were passed in an hour. She had no cough nor expectoration; the catamenia were irregular. She was ordered to take the compound krameria mixture with chloric ether $\text{m} \times$ three times a day, to have wine with arrow-root, and as a suppository, gr. x of the compound soap pill.

July 19th.—The skin was moist, and the tongue was thickly coated with a dry fur. She had had no sleep in the night. There were sordes on the teeth, and the breath was offensive; the pulse 112, compressible. The motions were offensive; the legs cold; the abdomen slightly tender. To continue the medicine, and to have a poppy poultice applied to the abdomen.

21st.—The poppy poultice afforded temporary relief, but the purging continued, and the patient was more prostrate. To take opium gr. j at once, and to add to the mixture $\text{m} \times$ of the tincture of opium.

23rd.—Patient weaker, drowsy from the action of the opium; pulse 118, feeble and compressible. She became more and more prostrate till death. The administration of lead and opium, and chloroform, afforded slight relief.

On inspection, the pharynx, the posterior part of the tongue, the tonsils, and the anterior part of the epiglottis were covered with a yellowish-white crust; this was very adherent to the lateral portions of the root of the tongue; and on the glossal surface of the epiglottis this membrane was so fixed that it could not be washed off by water. The surface of the epiglottis beneath it was much injected, but only towards the tongue. On examining a portion of this substance, it was found to consist of delicate interlacing torulæ, some jointed, and much resembling the torula cerevisiæ; it was mixed with particles of fat. The larynx, internal surface of the epiglottis, trachea, and bronchi were all free from disease. The lungs were healthy, so also the pleura, and were quite free from adhesions. The muguet did not extend to the œsophagus, excepting at its upper part. The heart and pericardium were healthy, but there was an excess of fat external to the heart.

Abdomen.—There were old adhesions of the omentum. The stomach was healthy, and on microscopical inspection its follicles were full of secreting cells and granules, with a small quantity of highly refracting particles. The liver extended to the right iliac region; it was exceedingly fatty, of pale-yellow colour, and *lighter* than water. The gall-bladder was contracted and empty; the weight of the liver 8lb. 6oz. *Spleen,* corpuscles distinct, its weight 6½oz.

The peritoneum was healthy; there was a moderate amount of fat in the mesentery, and the glands near the cæcum were enlarged and somewhat

swollen. The intestines were moderately distended. On opening the colon it was found to contain fluid fæces; the whole mucous membrane was covered by an adherent whitish layer, having a granular, almost villous appearance; it was of a yellowish-red colour, which was more marked towards the rectum; the mucous membrane was swollen, and in some parts presented small aphthous ulcers about a quarter of an inch in diameter. The submucous cellular tissue was white and thickened; the muscular coat also appeared very thick and distinct. This false membrane consisted of a blastema containing granules, highly refracting particles, and some cells, but no well-marked cells nor epithelium. The last foot of the ileum was much injected, and presented several irregular ulcers, but no membrane similar to that in the cæcum and colon.

The kidneys were pale and large, their weight 11oz. The uriniferous tubes contained granules and oil particles.

This case of acute inflammation of the colon came on gradually. For six months the patient had attacks of diarrhœa, but for seven weeks the symptoms were severe, and of the character of dysentery; the motions contained blood and mucus, and became afterwards green and slimy. On admission into Guy's Hospital she was in a typhoid condition, and almost dying. Her general appearance resembled a case of pneumonic phthisis with dysentery, but there were no physical signs of phthisis. In the treatment the patient was too prostrate to bear the application of leeches, otherwise it might have been desirable. The astringents which were administered afforded only very temporary relief, and opium quickly produced torpor of the brain.

CASE CLXIV.—*Diphtherite of the Colon. Dysentery. Chorea.*—Elizabeth H—, æt. 7 years, was admitted into Guy's, February, 1855. She was a dark, strumous child, who five weeks previously, without apparent cause, became affected with chorea; she improved under the use of sulphate of zinc gradually increased, and purgatives had been required, for the bowels were generally constipated. A few days before death diarrhœa came on, with prostration, and with symptoms very much resembling Asiatic cholera; the motions consisted at first of blood and mucus, but afterwards of thin and watery fluid. On the seventh day after the onset of the purging, the child died.

Inspection.—The eyes were much sunken. The cerebral veins were full of partially decolorized clot, and the ventricles of the brain contained more than the normal quantity of fluid. The lungs appeared healthy, except a circumscribed patch at the middle of the left lung, where was a strumous mass about the size of a hazel nut, and some tubercles around it. The inner aspect of the mitral valve was fringed with minute vegetations, firm, semi-transparent, and surrounding the edge of the valve; the largest of them

was of the size of a pin's head. The inner surface of the tricuspid was slightly roughened. The pericardium was healthy. The weight of the heart was $3\frac{1}{2}$ ozs.

The stomach was healthy. In the jejunum a few of Peyer's patches were visible and were injected; in the ileum they were very distinct, and near the cæcum were covered with a delicate inflammatory deposit of lymph. The large intestine was diseased throughout. The whole of the mucous membrane was of a dark-green colour, covered with firm granular deposit of lymph. The disease increased in severity from the cæcum downwards. The cæcum was acutely inflamed, being of a red colour, and the mucous membrane entire. Lower down the mucous membrane became green, and was covered with inflammatory deposit, and in the rectum the inner surface was raised into folds or irregularly shaped eminences. In the descending colon, when the adventitious product was removed, the tissue was seen to be swollen, full of blood, and in some parts superficially ulcerated. The muscular coat was much thickened. The kidneys were healthy.

This case was supposed to be one of Asiatic cholera, and, unless a careful inspection had been made, might have been so recorded. The suddenness of the diarrhoea, the rapid collapse and prostration, closely resembled that disease. Other cases of cholera had occurred about the time. The treatment used—starch and opium injections, &c.—did not appear to have any effect upon the malady.

CASE CLXV.—*Inflammation of the Colon. Hernia.*—Ann H—, æt. 60, was a single woman, a servant, who had resided at Peckham, and she was admitted into Guy's Hospital February 6th, 1857. For ten years she had had hernia, but her general health was good. Five days before admission the hernia came down, she was seen by a practitioner, and after taking croton oil, various purgatives, shot, &c., was admitted into Guy's. She was much depressed and collapsed; there were constipation, stercoraceous vomiting, great pain in the abdomen, and some tenderness. A tolerably large femoral hernia was found in the right inguinal region. A grain of opium was given at once, and the hernia was reduced by taxis without difficulty. On the following day, the 7th, the bowels were opened, and the patient felt better. A grain of calomel and one of opium were administered, and repeated every four hours; and she was allowed beef tea.

At night she had no sleep on account of pain, and the opium was repeated.

On the third day, the 8th, the bowels were relaxed, the tongue furred; she had much pain in the abdomen; the pulse was feeble and frequent. A chloroform draught was administered every four hours.

On the fourth day, the 9th, the purging continued, the stools consisted of mucus and fluid fæces, light in colour, and very offensive; the patient was prostrate, but she had less pain. An enema of soap and opium was administered.

On the fifth day, the 10th, the abdomen was slightly tender. She had

slept better. The diarrhœa continued, but she had no tenesmus. The enema was repeated. On the 11th the compound krameria mixture was given every six hours.

On the seventh day, the 12th, the diarrhœa still continued, the patient became more prostrate and was unable to sleep; the motions consisted of mucus and blood. On the following days the purging continued, and the prostration became at last extreme, the tongue dry, the pulse more feeble. Tormentilla with opium and tannic acid, and acetate of lead with opium, were given; but on the 28th, the fourteenth day after the hernia had been reduced, and the nineteenth after the commencement of the illness, she died. For some time before death there was low muttering delirium, cold extremities, and an almost imperceptible pulse.

Inspection was made a few hours after death. The body was much wasted, and the eyes were sunken. The lungs and pleura were quite healthy, and the lungs were collapsed; the heart was small.

Abdomen.—The peritoneum was healthy; the œsophagus was normal, so also was the stomach; the jejunum was congested, the valvulæ conniventes covered with mucus. Eight feet from the cæcum about four inches of the ileum was of a deep purple colour, and had evidently been strangulated; it appeared to be recovering. Below the strangulation the ileum became much more congested, and for four feet from the cæcum the intestine was acutely inflamed; the mucous membrane was covered by a yellowish-green adherent exudation, nearly a line in thickness, which was with difficulty removed. The valvulæ conniventes were exceedingly prominent, rigid, and erect. The whole of the mucous membrane was much thickened, the muscular coat more than a line in thickness, and the areolar tissue very œdematous.

On making a section of this portion of intestine (see Plate III, fig. 1), the exudation was found to consist of cells and granules, and appeared to be quite continuous with the follicles of the mucous membrane (Leiberkühn's); these follicles were exceedingly distinct, and were evidently distended to the utmost. It appeared that the secretion from these follicles was changed by the diseased action, that it was excessive in quantity, and abnormal in quality; the form of the follicles was retained by the exudation on the surface, as if it had been exuded rapidly. The submucous cellular tissue was much thickened from inflammatory œdema.

The colon from the cæcum to the rectum was still more diseased; in the cæcum the mucous membrane had a reddish-gray colour, with minute highly injected points studding the surface; some of these points presented a darker coloured centre, others were scarlet. In the place of others were minute ulcers, the mucous membrane being eroded; the edges of the ulcers were injected, their surface whitish-grey, but there was no apparent slough. In the ascending colon more of the mucous membrane was destroyed, and elongated ulcers were found, about half an inch in length, with irregular, injected, or partially undermined margins, their base consisting of whitish lymph-like exudation; between these ulcers there were minute red points or red mucous membrane covered with exudation. The whole of the colon presented a similar appearance, even to the rectum. The appendix cæci was long, and at its extremity were about half a dozen small shot.

The examination of the mucous membrane of the cæcum showed that the minute red points were solitary glands (see Plate III, fig. 2, which represents one of these red glands, having a deeply coloured centre). The portion above the level of the mucous membrane showed distended capillaries, and there was in the centre an opening extending into the gland. In the gland were minute nuclei, and a large nucleated cell was observed. It appeared probable, from the darker colour of the congested part, that the circulation in these capillaries had nearly ceased, and that in a short time sloughing would have taken place. The adjoining mucous membrane presented an appearance somewhat similar to that found in the ileum, but the follicles were less distinct. Some crystals were observed on the surface.

The liver was healthy, so also the spleen, kidneys, bladder, uterus, ovaries, and the mesenteric glands; in fact no disease was found except in the intestines. The hernial sac was empty, and its opening nearly closed.

In this case there was acute inflammation of the whole of the colon, and of several feet of the ileum; the changes were of a degenerative kind, and rapid in their action. Was the disease dysentery, or inflammation of a different kind? In the cæcum and colon we found that the solitary glands were diseased, and that destruction of the membrane had ensued; but these glands were not the only parts affected, the follicles and all the structures were diseased, as we found with the follicles in the ileum; the mucous membrane and the muscular coat were œdematous. Very drastic purgatives and violent means had been used at the onset of the disease, and these probably contributed to the severity of the affection. The bowels began to act very shortly after the hernia had been reduced, and did not cease to act till death. As to the symptoms, they were those of dysentery; the stools consisted of blood and mucus, the prostration became gradually extreme, the pain was sometimes severe, but she never had the tenderness of peritonitis. The disease was probably, to a great extent, constitutional in its character, but it was excited by direct irritation.

The application of leeches to the anus might have afforded relief, but it must be remembered that the patient was sixty years of age, collapsed by hernia and stercoraceous vomiting; and if depletion had been used the prostration would have been increased. The application of astringents to a surface so extensively inflamed and diseased, when administered by the mouth, are inefficient remedies; they have to pass over at least twenty feet of mucous membrane before reaching the colon.

Injections are more direct, and are applied at once to the diseased surface; starch and opium was the only injection used; but even in this mode the remedy could scarcely be expected to reach the colon in its whole length.

CASE CLXVI.—Dysentery. Ulceration of Small Intestines. Perforation. Faecal Abscess. Peritonitis.—Edward B—, æt. 39, was admitted October 19th, 1853, and died November 8th. He had been a stout, hearty man, living at Walworth, and a labourer in the London Docks. His health had been unimpaired, and his habits of life regular. Six weeks before admission, he ate a considerable quantity of coarse sugar, and three days afterwards he had diarrhœa, with copious liquid evacuations, and severe colic. This state continued till admission; he had become thin and weak, having been unable to take any food since the commencement of his illness. On admission, the bowels acted every half hour. The evacuations were of a dark-brown colour, and no blood was passed. There was tenderness over the cæcum and descending colon; he had no appetite, the mouth was dry and parched, and he had considerable thirst; he had nausea, but did not vomit. The countenance was dejected, and the body emaciated and feeble.

The heart and lungs appeared to be healthy. The tongue became very red and injected. Great prostration came on, and sordes formed on the teeth, the purging continuing unabated. Kino, cusparia, opium, gallic and sulphuric acids, &c., were administered, with only slight relief.

Inspection eighteen hours after death. *Chest.*—Slight adhesion existed at the apex of the right pleura; and in the lung at the same part there was calcareous deposit and old thickening of the tissue; the lungs were otherwise healthy. The heart was flabby; the right ventricle contained firm fibrin and clot; the left side also was moderately distended. The valves were healthy, with the exception of slight atheromatous thickening about the aortic semilunar valves. The abdomen was collapsed. The intestines were contracted, but there was universal moderate injection of the peritoneum, and the coils of the intestine were united by fibrinous adhesions. On separating the abdominal parietes from the sigmoid flexure, a part of the intestine was found to be perforated, and all its coats destroyed for a considerable part of its circumference; a small faecal abscess had been formed. At the cæcum also and rectum, extravasation of fæces was only prevented by external adhesions. The large intestine was ulcerated in its whole length; at the cæcum were several transverse patches of ulceration; immediately above the cæcum the mucus and muscular coats were much thickened, apparently from older disease; beyond this, the mucous membrane presented a large slough, and nearly all the mucous membrane was destroyed. Here and there were granular masses like tubercles, or larger portions which had become intensely congested, and were raised above the ulcerated surface, giving it a polypoid appearance. The sigmoid flexure and rectum were equally ulcerated; in some parts the mucous membrane only was destroyed, in others nearly all the coats. In the small intestine, about 8 inches from the cæcum, the lining membrane was intensely congested, and Peyer's glands

presented several small aphthous ulcers. The columnar epithelium was scanty, but numerous cells like mucus were observed. The mesenteric glands were enlarged. In the stomach were several points of arborescent injection. The spleen was healthy; the liver fatty, and its weight was 4lbs.; the gall-bladder was moderately distended.

The destruction of the mucous membrane of the colon in this case was very great, both as to extent and depth; fæcal abscess had been formed subsequent to the perforation. The disease lasted nearly nine weeks, and the remedies did not at all check the symptoms. It is probable that on admission (for the disease had then continued for six weeks) considerable sloughing of the colon existed.

There was no evidence of any miasmatic influence nor noxious effluvia, but the diarrhœa and subsequent dysentery were produced by the foolish excess of the patient.

CASE CLXVII.—Ulceration of Large Intestine. Perforation. Submucous suppuration. Pus in the Portal Vein, and Inflammatory Patches in the Liver.—James T—, æt. 59, was admitted into Guy's October 12th, 1853, and died on the following day. He was a labourer in the London Docks, and had had "bowel complaint" for two months; the symptoms had gradually become worse, and a week before his death he was confined to his bed; he had repeated purging of blood, and became much emaciated.

On admission he had a tumid and tympanitic abdomen, but tolerant of pressure; the skin was of a dingy colour, and the tongue was red, glazed, and dry; he was in a prostrate condition when admitted, and died the following morning.

Inspection fifty-seven hours after death.—The body was spare. *Chest.*—The pleura at the left apex presented a little cartilaginous thickening. There was a white patch over the ventricles, the right was distended with clot, the left was empty; the mitral valve was thickened and slightly contracted.

The peritoneum was universally inflamed, injected, and covered with effused lymph. The cavity contained dirty fluid, green in colour, and of an offensive, feculent odour. On turning aside the large intestine, an opening was found above the cæcum. The whole length of the large intestine was ulcerated. These ulcers were transverse, and were generally about two inches in length. The mucous membrane was ragged, and covered with a black slough; the circumference of these ulcers was thickened. In some parts the mucous membrane was quite destroyed, and the intervening portions of mucous membrane were œdematous. The peritoneal surface of the large intestine was observed to be here and there of a yellow colour; on making a section at these parts, the subserous coat was found to be infiltrated with pus extending from the submucous coat. These abscesses were situated on the mesenteric side of the intestine. The mucous and muscular

coats of the rectum were much thickened, and at the lower part of the descending colon was a puckered portion of intestine, ecchymosed and injected.

The small intestines were healthy. The stomach presented partial injection. The spleen and kidneys were healthy. The liver presented a thickened layer of peritoneum at its lower border (attrition). Its section was yellow, and its structure soft; on the convex surface of the right lobe was an irregular congested portion about one inch in diameter, and at its centre was a branch of the portal vein filled with pus.

Glisson's capsule was thickened; no apparent disease of the trunk of the portal vein nor of the inferior mesenteric vein existed; the liver generally was fatty; weight 3lbs. 10ozs. On microscopical examination, the liver was found to contain very much fat, and Glisson's capsule was much thickened by fibrinous deposit. The thickening of the intestine around the ulcer was found to be composed of hypertrophied muscular fibre, but principally of mucous membrane, with bands of fibrous tissue passing through it.

This was one of the most severe cases of inflammation of the intestine that I have seen; large ragged abscesses extended throughout the colon, and had led to perforation and to peritonitis. The commencing suppuration in the liver, and the pus in the portal vein, were confirmatory of the views of Dr. Budd, as to the causes of abscess of the liver. When admitted into Guy's he was in a dying condition, but the disease had existed for two months.

CASE CLXVIII.—Dysentery. Perforation of Colon.—Sarah W—, *æt.* 34, was admitted into Guy's Hospital April, 1847, after having been ill for three months, and severely so for three weeks. She had resided at Huntingdon, then at Lambeth, and was the wife of a fishmonger. Three months previously she had had pitchy evacuations, evidently containing blood; but she had been free from pain. Three weeks before admission she had profuse purging; the evacuations contained blood, and scarcely any solid *fæces*; she suffered much tenesmus and general pain in the abdomen; the pain was occasionally aggravated, but was especially situated at the right iliac fossa. *Scybala* were occasionally passed. There was febrile excitement, and before death vomiting came on.

She took opium alone, then lead, ipecacuanha, and copper; mercurial inunction was used; leeches and blisters, &c., were applied to the abdomen. Wine and suet and milk were administered.

Inspection.—The body was well nourished, and there was a considerable amount of fat in the abdominal parietes. On opening the peritoneum, it was found to be exceedingly dry; the transverse colon was adherent to neighbouring viscera by soft adhesions; the omentum extended to the pelvis, and on raising it soft adhesions were found between it and the intestine, which were also much injected at their points of contact with each other. In the left iliac region, on drawing aside the sigmoid flexure, soft adhesions gave way, and a small circular perforation was found; no extravasation had taken

place. In the right iliac fossa, the cæcum was more firmly adherent; and close to the union of the vermiform appendix a long defined opening was observed, but closed by adhesions.

There were also several perforations in the ascending and transverse colon similarly closed; in other parts the peritoneum only was left. Perforations had also taken place in the rectum, but no extravasation had followed from any part. The vermiform appendix was healthy. The cæcum and ascending colon were distended and thickened.

On opening the whole length of the large intestine, the following appearances presented themselves:—Portions of mucous membrane which had escaped ulceration were softened, and were of a greenish or red colour; large transverse ulcers were found at other parts, their margins were defined; in some the peritoneum formed the base, and in nearly a dozen places the peritoneum also was destroyed. Hard, dry scybala adhered in some parts. About six inches above the cæcum the intestine appeared somewhat contracted, and large pouches were formed both above and below. The small intestines were pale, and no disease was observed in them; they contained fluid fæces. The mucous membrane of the stomach was thickened and softened. The liver was pale and soft. The gall-bladder was much contracted and adherent to the colon; it contained a small quantity of white, thick mucus, and crystals of cholesterine; the duct was blocked up by a gall-stone, which was about half an inch in circumference. The spleen was larger than natural, and it was softened. The kidney presented an irregular contraction on its surface.

The lungs were emphysematous; one or two consolidated lobules were situated at the apex. The heart was flabby; but, with the exception of slight atheroma, the valves were healthy.

In this case the most severe inflammation of the colon had been set up, the coats of the intestine had sloughed, and numerous perforations had resulted. The disease had lasted for three months; but a short time before admission it became much aggravated. It could scarcely be expected that the administration of small doses of astringents could check such extensive degeneration; and it was evident that the patient died, not from exhaustion, but from the severity of the disease, and its extension to the peritoneum.

CASE CLXIX.—*Chronic Bronchitic Phthisis. Cirrhotic and Lardaceous Liver. Contracted Abscess of Liver. Chronic Dysentery, and Chronic Peritonitis.*—Thomas R—, æt. 31, a soldier, or rather pensioner, was admitted under my care Oct. 24th, 1856. He had been in the West Indies as a soldier, and had been exceedingly intemperate in his habits, spending all his money in rum, &c. He stated, however, that till three years previously he enjoyed good health, but had had syphilis six or seven times, and had been salivated five times; and that when a child he had had ague.

Two and a half years before, while serving in Bermuda, he was exposed to cold at night; the following morning he had severe cold and cough; but

did not report himself as ill for six months, having then gradually become much worse. At that time he spat blood, and had night sweats, and had great pain in the precordial region. He remained in hospital for nine months, and left very little relieved. He subsequently went to the Crimea, but was at once invalided, and sent to Scutari.

Sixteen days before his admission into Guy's his ancles became swollen, and dropsy rapidly increased.

He was a tall, emaciated man, with an exceedingly anxious, haggard expression; the nails were clubbed; the respiration was difficult and hurried; and he was almost in a dying condition. He complained of pain in the chest and abdomen; the respiration was twenty-four per minute, and he expectorated much thick, greenish, and rusty-coloured mucus.

The chest could only be examined anteriorly. On the *left* side it was generally dull; the apex was flattened, and gurgling was very distinct. The right was dull at the base, and sibilant râles were audible. The abdomen was very much distended, and hot; the superficial veins were enlarged; the liver could not be distinctly felt, but fluctuation was very perceptible; the bowels were relaxed; the evacuations were thin and watery; the appetite was bad; urine high-coloured, non-albuminous, sp. gr. 1·012.

Opium gr. j was given, and repeated at night, and a poultice was applied to the abdomen. On the following day, the Compound Logwood Mixture of Guy's, and some port wine were prescribed.

The heat and pain of the abdomen became somewhat less, and the patient rather more comfortable for a few hours. He gradually sank, and died November 2nd. Before death the scrotum became distended, erythematous, and almost purple in colour.

Inspection, Nov. 3rd.—The body was much emaciated. The abdomen was distended, but the enlargement of the superficial veins had disappeared. *Chest*.—Pleura universally adherent on the left side, and much thickened; at the apex it was semi-cartilaginous. On the right side there were adhesions at the apex; but below there was some effusion, with a few flakes of fibrin. *Left lung*.—At the apex were several small vomices freely communicating with the bronchi, and surrounded with dense red lung; nearly the whole lung presented fleshy tissue, with irregularly dilated bronchi, which in some places extended nearly to the periphery; dense white fibrous tissue, and some iron-grey pneumonia were found in other parts. There was evidently old pulmonic disease, and more recent acute disease with it; quite at the base was a small portion of crepitant lung. On the right side there was old pneumonic condensation; but throughout the lung there was granular deposit in small clusters, resembling tubercles, with some condensation around them. The bronchi were much injected, and the mucus membrane thickened.

Heart.—In the pericardium there was an excess of fluid, and some flakes of fibrin. The valves were healthy.

Abdomen.—The peritoneum contained several gallons of fluid; the intestines were moderately distended. The peritoneum was opaque, slightly granular, and very delicate bands were found between the intestine, like sticky albuminous serum. The liver was much contracted, nodulated, and its surface was opaque. On section, an irregular cheesy mass was found,

about three inches in length, extending from the surface of the liver into its substance, and surrounded by slight fibrinous investment; the surface was contracted. This mass appeared to consist of two or three collections appended the one to the other. There were numerous other small cheesy masses of smaller size, situated throughout the liver, in the course of the portal branches. They were all apparently the result of inflammatory action, or dried abscesses. The rest of the liver was semi-transparent, and in many parts was lardaceous. The gall-bladder was contracted. The spleen was enlarged and lardaceous. The kidneys were healthy.

Colon.—The whole of the colon presented irregularly healed ulcers, and was granular and thickened; and scarcely any healthy mucous membrane was observable. Small circular, smooth spaces, evidently healed ulcers, studded the whole surface. The coats of the intestine were thickened. The ileum and stomach were healthy; so also were the kidneys. The omentum was adherent near the inguinal canal on the left side.

The original malady appeared to have been dysentery, contracted in the West Indies, and it was followed by abscesses in the liver; these abscesses dried, and probably constituted the cheesy masses found after death, affording a remarkable instance of abscess of the liver, and perhaps of pyæmia not necessarily fatal. The intemperate and dissolute habits set up other diseased action, as that in the lungs; the inflammatory deposit there had broken down, and had led to the formation of a vomica; repeated attacks of inflammation took place, and the disease encroached more and more upon the lung tissue, and at last in a more severe attack, ascites came on and led to fatal termination. A few granular collections were the only representatives of tubercles, which were, probably, deposited a short time before death, when nutrition was so much impaired. The lardaceous state of the liver was interesting in its connexion with syphilis, and struma. The colon was filled with cicatrices; and they had led to partial obstruction, as shown by the hypertrophy of the intestinal coats.

CASE CLXX.—*Chronic Dysentery. Hepatic Abscess. Pyæmia. Abscess in the Brain and Lung.*—Thomas D—, æt. 25, was admitted February 14th, and died March 19th, 1855. He was a sailor, and had been for two years in the East Indies. At Burmah he had ague and dysentery, and was ill for several weeks; and for two months he had had pain in the side.

On admission, he was sallow, and generally cachectic. There was pain in the right side; the chest was dull; and it was supposed from the history that he had abscess in the liver.

On February 21st, when sitting by the fire, he fell down in a fit, and was

convulsed; for several days he continued in a semi-conscious condition. On the 28th, he could speak and give his name; he continued apparently to improve till the 14th, when he again fell into a semi-conscious state. On the 16th he was unable to sit up and take his breakfast; but shortly afterwards became quite insensible; and had stertorous breathing, which continued till death. It was observed, throughout, that the right leg was weak, and at last was paralysed; the right pupil was smaller than the left; but a few hours before death it became widely dilated.

Inspection twenty-four hours after death. *Brain*.—The surface of the hemisphere was dry; and at the base were slight adhesions between the surfaces of the arachnoid. In the posterior lobe of the left hemisphere was an abscess about the size of a hen's egg, containing thick, tenacious pus; it nearly reached the surface, and was surrounded with softened brain substance; at the anterior part of the abscess was a clot of blood, also surrounded by softened tissue. The abscess had broken into the left lateral ventricle at its posterior corner; the left ventricle was filled with pus; the right with about $\frac{3}{4}$ of clear serum; the fourth ventricle was healthy.

In the *chest* were old and recent adhesions at the bases of both pleural cavities. The bronchi were slightly inflamed, and contained muco-purulent secretion. The base of the left lung contained a small abscess; the base of the right was in a state of incipient pneumonia.

Liver.—In the right lobe, at the upper surface, were two chronic abscesses, capable of holding about $\frac{3}{4}$ of pus; the pus was thick and green; the walls of the abscess were very thick, bounded by a smooth cyst, and firm tissue about one eighth of an inch in thickness; on the circumference of the abscess a compressed vein was observed.

In the *colon* the mucous membrane was thickened; several well-marked cicatrices were found in the ascending colon; the mucous membrane was puckered, and in some parts was of a slate colour; the muscular coat was slightly hypertrophied.

It appears probable that the dysentery, which had been contracted in Burmah, had led to abscess in the liver, and that this remained passive for many months, producing hectic, with pain in the side, &c., and at last, from some fresh exciting cause, new action was set up, acute pyæmia produced, and abscess in the brain the consequence. Dr. Hughes diagnosticated this course of morbid changes, which was completely confirmed, on inspection after death.

CASE CLXXI.—Dysentery. Abscess of the Liver. Perforation of the Diaphragm. Empyema.—John J—, æt. 32, admitted into Guy's Hospital, July 28th, 1858, under the care of Dr. Wilks. Till February, 1857, he had been perfectly well, and went to the coast of Guinea; during two months he remained on that coast, and had dysentery, producing much purging, with tenesmus, and a discharge of a small quantity of blood. The dysenteric

symptoms continued for two months, and quite incapacitated him from work. The symptoms came on suddenly, and were accompanied with severe pain; he took spirits for relief, but the following day pain again returned, with a febrile condition, and with severe headache, &c., he was then completely laid aside; and for a month was unable to take food. On June 14th, he arrived in London, and had been under treatment till application at the hospital.

He was a man of fair complexion, slightly sallow, and emaciated. He complained of pain in the right side, had occasional rigors; but no cough; the bowels acted regularly; the tongue was slightly furred. The abdomen was collapsed; but there was pain in the right side of the chest, which was uniformly dull; tubular breathing was audible, and on the left side puerile respiration; and it was evident that fluid effusion had taken place into the right pleura. The pulse was irritable and compressible, and the appetite was altogether gone; he had very little sleep. Comp. infusion of gentian, with iodide of potassium, and syrup of poppies, were given three times a day. One week before death, his distress greatly increased, and paracentesis thoracis was performed, and a pint of pus was evacuated; he died, August 20th.

On inspection, the right pleura was found to be filled with several pints of grey purulent fluid; the compressed lung being at the posterior and upper part of the chest; a large opening in the diaphragm on the right side allowed free communication between the pleura and an abscess situated between the liver and diaphragm, it was this abscess which had been opened by the paracentesis. The left lung and the heart were healthy. In the abdomen there were local adhesions between the intestines in various parts; at the lower part of the ileum, and in the ascending colon, there was some recent ulceration, and in some other parts a small quantity of granular lymph was found covering the mucous membranes. In the cæcum were two large ulcers of older date, with raised thickened edges, and the mucous membrane destroyed; towards the rectum the mucous membrane was of a slate colour.

The large abscess above the liver was between the liver and the diaphragm, and the capsule of the liver formed part of the walls of the abscess; there were several abscesses in the substance of the liver itself, and the whole of the gland was filled with small deposits of pus. Many of the hepatic veins were filled with clot and pus, the latter having apparently entered the veins secondarily from the tissue; the liver weighed 9½lbs. The spleen and kidneys were healthy.

This was a case of great interest. Acute dysentery, which the patient had contracted on the coast of Guinea, was followed by hepatic abscess. This latter abscess extended to the surface of the liver, and led to perforation of the diaphragm; the symptoms of pleuritic effusion were then produced, and led ultimately to a fatal termination. Dr. Wilks had diagnosed the purulent collection, and determined to draw off the fluid; the trochar passed above instead of beneath the diaphragm; but since the communication through the diaphragm was perfectly free, the

whole of the contents of the abscess and of the pleura could be emptied at the same time. There was evidence of old disease in the mucous membrane of the colon.

CASE CLXXII.—*Cicatrization and Contraction of the Rectum, and Sigmoid Flexure after Dysentery.*—Evan T—, æt. 33, was admitted into Guy's Hospital November 16th, and died November 22nd, 1855. He had received, from a weight falling upon him, fracture of the pelvis, which produced laceration of the urethra, &c.; he became prostrate, and died from the consequences of the accident in four days.

He was a sailor, and had been living at Deptford; he had previously had dysentery, and had been subject for some time to winter cough. His habits of life had been intemperate.

Inspection twelve hours after death. The head was not examined. In the chest were some patches of ecchymosis on the pleura; and in the lungs several lobules were grey and granular; this state was the pneumonia of pyæmia. There was extravasation of blood into the peritoneal cavity; and the peritoneum was inflamed. The membranous portion of the urethra was lacerated, and the cellular tissue around the bladder was sloughing.

In the rectum and sigmoid flexure, the mucous membrane was grey, and thickened; so also was the submucous cellular tissue; in several parts there was irregular corrugation, apparently from cicatrization. At the commencement of the rectum the calibre of the intestine was diminished; in some parts the mucous membrane had been destroyed, but the cicatrix was smooth, though destitute of gland tissue. In the prominent portions elongated glands were very distinct, resembling follicles of Lieberkühn.

This case is recorded as a marked instance of the appearance presented by dysentery which had been cured. The cicatrices in the sigmoid flexure and rectum showed the severity and extent of the previous ulceration, and by the contraction of the new tissue served, in some measure, to constrict the intestine.

CASE CLXXIII.—*Chronic Ulceration of the Intestine. Dysentery. Cicatrization. Contraction. Perforation. Abscess near the crest of the Ilium* (From the Museum Records).—Mr. W—, was a young gentleman of tolerably temperate habits, but who had had syphilis many times; several years before his death he had a dysenteric affection, on the subsidence of which his bowels were habitually constipated; this state was attributed to stricture of the rectum, which was felt at no great distance from the anus. A bougie was passed with the effect of considerably dilating the stricture. He subsequently went to America, but he did not prosper. The death of his wife and misfortune were followed by declining health. An abscess formed above the crest of the ilium, towards the posterior part on the left side, and there was continual pain at that part; after the application of leeches several sinuses formed, diarrhoea came on, and he wasted rapidly.

On inspection, except pleuritic adhesions, the thoracic viscera were healthy. In the left iliac region the integuments were separated from the tendon of the external oblique, by the sinuous ulceration. In that region, the intestines were glued together; the peritoneum and adjacent cellular membrane were much thickened; the rest of the peritoneum was healthy. In the sigmoid flexure there were numerous traces of old ulceration, of a lightish green colour, the surface was uneven, and the structure of the intestine at the part was thickened and condensed; the calibre of the intestine was also much contracted. There were three or four small perforations in the intestine at this part; the rectum was healthy, except immediately above the anus, where there was considerable thickening with induration. This evidently depended on an old ulcer, occupying about half the intestine; and the part was of a leaden colour. The liver was much enlarged and fatty. The gall-bladder contained some ropy mucus. The kidneys and the rest of the intestines were healthy.

This case is a very interesting one, for although the dysentery was relieved, the cicatrization and subsequent contraction were followed by constipation; ulceration was set up above the points of contraction, and ultimately the intestine was perforated. The sinuses opened near the crest of the ilium; fæces do not appear to have been discharged; but the case might easily have been mistaken for suppuration from diseased bone.

The constriction in these instances arises from fibro-elastic tissue, which becomes more dense than the original muscular coat; it closely resembles that found after the destruction of the skin in burns, and has a similar disposition to contract.

II. Acute inflammation of the colon sometimes takes place in common with diseases of other organs; and these cases present a marked difference from those previously detailed.

The thoracic viscera are affected with acute disease; the bronchial tubes and lungs are inflamed, and so, in fact, are almost all the mucous membranes. The symptoms of disease of the chest are more marked than those of the abdomen; dyspnoea, cough, febrile excitement are present with the physical signs of thoracic disease. The countenance is anxious and flushed, the skin hot and dry, or clammy, the pulse becomes gradually more depressed, the tongue brown and dry, and the patient prostrate. These symptoms are accompanied by dysenteric diarrhoea, which indicates a diseased state of the colon.

In some of these cases, the exciting cause of the inflammation of the lungs and bronchi is also the cause of like disease of the

whole of the contents of the abscess and of the pleura could be emptied at the same time. There was evidence of old disease in the mucous membrane of the colon.

CASE CLXXII.—*Cicatrization and Contraction of the Rectum, and Sigmoid Flexure after Dysentery.*—Evan T—, æt. 33, was admitted into Guy's Hospital November 16th, and died November 22nd, 1855. He had received, from a weight falling upon him, fracture of the pelvis, which produced laceration of the urethra, &c.; he became prostrate, and died from the consequences of the accident in four days.

He was a sailor, and had been living at Deptford; he had previously had dysentery, and had been subject for some time to winter cough. His habits of life had been intemperate.

Inspection twelve hours after death. The head was not examined. In the chest were some patches of ecchymosis on the pleura; and in the lungs several lobules were grey and granular; this state was the pneumonia of pyæmia. There was extravasation of blood into the peritoneal cavity; and the peritoneum was inflamed. The membranous portion of the urethra was lacerated, and the cellular tissue around the bladder was sloughing.

In the rectum and sigmoid flexure, the mucous membrane was grey, and thickened; so also was the submucous cellular tissue; in several parts there was irregular corrugation, apparently from cicatrization. At the commencement of the rectum the calibre of the intestine was diminished; in some parts the mucous membrane had been destroyed, but the cicatrix was smooth, though destitute of gland tissue. In the prominent portions elongated glands were very distinct, resembling follicles of Lieberkühn.

This case is recorded as a marked instance of the appearance presented by dysentery which had been cured. The cicatrices in the sigmoid flexure and rectum showed the severity and extent of the previous ulceration, and by the contraction of the new tissue served, in some measure, to constrict the intestine.

CASE CLXXIII.—*Chronic Ulceration of the Intestine. Dysentery. Cicatrization. Contraction. Perforation. Abscess near the crest of the Ilium* (From the Museum Records).—Mr. W—, was a young gentleman of tolerably temperate habits, but who had had syphilis many times; several years before his death he had a dysenteric affection, on the subsidence of which his bowels were habitually constipated; this state was attributed to stricture of the rectum, which was felt at no great distance from the anus. A bougie was passed with the effect of considerably dilating the stricture. He subsequently went to America, but he did not prosper. The death of his wife and misfortune were followed by declining health. An abscess formed above the crest of the ilium, towards the posterior part on the left side, and there was continual pain at that part; after the application of leeches several sinuses formed, diarrhoea came on, and he wasted rapidly.

On inspection, except pleuritic adhesions, the thoracic viscera were healthy. In the left iliac region the integuments were separated from the tendon of the external oblique, by the sinuous ulceration. In that region, the intestines were glued together; the peritoneum and adjacent cellular membrane were much thickened; the rest of the peritoneum was healthy. In the sigmoid flexure there were numerous traces of old ulceration, of a lightish green colour, the surface was uneven, and the structure of the intestine at the part was thickened and condensed; the calibre of the intestine was also much contracted. There were three or four small perforations in the intestine at this part; the rectum was healthy, except immediately above the anus, where there was considerable thickening with induration. This evidently depended on an old ulcer, occupying about half the intestine; and the part was of a leaden colour. The liver was much enlarged and fatty. The gall-bladder contained some ropy mucus. The kidneys and the rest of the intestines were healthy.

This case is a very interesting one, for although the dysentery was relieved, the cicatrization and subsequent contraction were followed by constipation; ulceration was set up above the points of contraction, and ultimately the intestine was perforated. The sinuses opened near the crest of the ilium; fæces do not appear to have been discharged; but the case might easily have been mistaken for suppuration from diseased bone.

The constriction in these instances arises from fibro-elastic tissue, which becomes more dense than the original muscular coat; it closely resembles that found after the destruction of the skin in burns, and has a similar disposition to contract.

II. Acute inflammation of the colon sometimes takes place in common with diseases of other organs; and these cases present a marked difference from those previously detailed.

The thoracic viscera are affected with acute disease; the bronchial tubes and lungs are inflamed, and so, in fact, are almost all the mucous membranes. The symptoms of disease of the chest are more marked than those of the abdomen; dyspnoea, cough, febrile excitement are present with the physical signs of thoracic disease. The countenance is anxious and flushed, the skin hot and dry, or clammy, the pulse becomes gradually more depressed, the tongue brown and dry, and the patient prostrate. These symptoms are accompanied by dysenteric diarrhoea, which indicates a diseased state of the colon.

In some of these cases, the exciting cause of the inflammation of the lungs and bronchi is also the cause of like disease of the

the cerebral oppression. Infusion of serpentary, with ℥xv of aromatic spirit of ammonia, and ℥xx of the compound tincture of camphor were given every four hours, and every night Dover's powder, gr. v, with grey powder, gr. ij.

July 19th. There was the rusty expectoration of acute pneumonia; the cough was frequent, and the delirium continued as before, so also the diarrhœa. Chalk mixture was given.

July 20th. The diarrhœa was less, but the tongue covered with a thick brown fur.

July 21st. The symptoms of pneumonia continued. Dover's powder, gr. v, with grey powder gr. j, were given three times a day, and the serpentary mixture was continued.

July 23rd. The right apex was more free; the expectoration yellow and rusty; the tongue less furred; but the patient was exceedingly prostrate. Wine, ℥viij was ordered.

July 24th. Greater exhaustion; low muttering delirium. Dover's powder, gr. x; grey powder, gr. j, given at night; to continue the mixture.

The prostration increased, and the patient gradually sank; he died August 3rd. There were no maculæ on or after admission; the diarrhœa did not continue after the first few days, nor was any blood discharged with the motions.

Inspection, August 6. The body was of moderate stature, and spare, the face was wasted; decomposition had moderately advanced. *Chest*.—On the right side there was effusion of fibrin, and about a pint of bloody purulent serum in the pleural sac. The lower lobe of the right lung was compressed, the upper rather fleshy; the left pleura also was quite free from adhesions; the lower lobe was in a state of pulmonary apoplexy, the upper part of the same lobe was very much congested and slightly consolidated; it sank in water. The bronchi, trachea, and larynx were healthy, but stained by post-mortem transudation; the bronchial glands did not appear to be diseased. The *pericardium* contained several ounces of bloody serum; the heart was flabby, but was otherwise healthy. The *abdomen* was moderately distended; the peritoneum was smooth and healthy; the stomach presented advanced gastric solution, the mucous membrane at the cardiac extremity being exceedingly thin; the small intestine contained yellow, bilious fluid fæces; in the last foot of the ileum the mucous membrane was grey, Peyer's patches were slightly raised, and in a few parts presented irregular ulceration; they had not, however, the raised swollen appearance common in typhoid fever. The cæcum contained several small ulcers, and the whole mucous membrane was of an iron-grey colour; the edges of most of these ulcers were smooth and contracting; the mucous membrane was thickened. In the transverse and descending colon, and in the rectum the whole of the mucous membrane had a swollen, œdematous and almost villous appearance; this was most marked towards the rectum. Studding the swollen membrane were white patches of adherent lymph, in some parts merely constituting a thin, delicate, but adherent membrane, in others, forming a large flocculent mass about three quarters of an inch in length, firmly adherent to the membrane; there were

numerous small ulcers scattered over these portions of the intestine; some with smooth, others with irregular and congested margins; some contained a small portion of false membrane, like a slough; from others large masses of false membrane could be detached. On tearing off portions of this membrane, an injected granular surface or superficial ulceration was observed. The submucous cellular tissue was white, thickened, and œdematous; the muscular coat was contracted and distinct, and nearly one eighth of an inch in thickness; the mesenteric glands were enlarged. On carefully examining portions of the false membrane, it was found to consist of granular cells closely matted together with very little blastema; the cells were large and full of granules, some contained a faintly marked nucleus; but scarcely any columnar epithelium was observed. On examining the mucous membrane itself, small excavations were found to contain similar cells. (Plate III, fig. 3.) The liver and spleen were healthy.

This diseased condition of the colon had led to the effusion of a blastema, and to the production of cells, instead of the ordinary columnar epithelium; the follicles and solitary glands had become inflamed, and a subsequent condition of this action produced superficial ulceration. The disease was of an inflammatory character, and not only the follicles and glands, but the whole surface of the mucous membrane was affected; the effused membrane appeared more cellular than is sometimes observed.

The symptoms closely resembled those of typhoid fever or typhoid pneumonia. The irritation of the large intestine, as indicated by the diarrhœa, appeared to be checked by the Dover's powder which was administered. The administration of alcoholic stimulants in a case of this kind is a question of great difficulty; the general symptoms—failing pulse, subsultus, &c.—appeared to require the free administration of stimulants. This case strongly favoured the idea of a constitutional origin of the disease, which resembled, if it were not identical with typhoid fever.

III. Another class of cases are those in which chronic disease has slowly advanced, for months or for years; acute inflammation of the alimentary canal is then set up, and in a short time leads to fatal termination. Cases of this kind are not of unfrequent occurrence, in which a patient is already broken down, as by incipient phthisis; there is already a disposition to the ulcerative disease of the small intestine, so common in phthisis, when, probably from fresh exciting cause, acute inflammatory

disease is set up, and diarrhœa, which can scarcely be checked by any means, is rapidly fatal. This inflammation, and the effusion of false membrane, may be confined to the rectum or descending colon, or be found in the cæcum and ileum. (See Strumous Disease of Intestine, Cases CXXIII, CXXIV, and CXXV.

In Case XI of perforating ulcer of the trachea and œsophagus, the rectum was acutely inflamed, but this appeared to have been produced by the direct irritation of the nutrient injections, by which life was entirely sustained for six weeks.

CASE CLXXVII.—*Strumous Peritonitis. Diphtherite of Rectum. Chronic Inflammation of the Large Intestine. Tubercular Lung. Simple Ovarian Cyst. Vascular Polypus of the Urethra.*—Esther W—, æt. 37, was admitted October 25th, 1854. She had menstruated regularly till eight months previously; and two months afterwards she perceived enlargement of the abdomen; this increased gradually with pain; and the swelling appeared to proceed from the right side.

November 20th. Six quarts of purulent fluid were drawn off by tapping; after a few days, sickness and diarrhœa came on, which continued more or less until death.

The body was extremely wasted; the eyes were sunken; the abdomen projected as a large round fluctuating tumour. Both lungs were closely adherent to the chest; they contained large masses of tuberculo-pneumonic deposit; and scattered groups of tubercle were found in both lungs. In the upper lobes the tubercles were associated with indurated puckered lung and cretaceous matter. The heart was small but healthy.

Abdomen.—The parietes were adherent to a cyst, which contained about a gallon of fluid. The intestines were adherent to one another, and presented here and there white tubercular masses; the greater part of the small intestine was behind the cyst, but about two feet of jejunum were closely adherent to its upper and anterior surface. Throughout the large intestine were raised rough and hardened patches of mucous membrane; and in the centre of these patches the mucous membrane was destroyed. The margins of these ulcers presented imperfect gland structure, with a considerable quantity of fibrous tissue; and in the centre also was fibrous tissue, but no gland structure. These ulcers covered a considerable portion of the large intestine; and the whole of the submucous tissue contained white fibrous tissue. In the intervening portions of membrane were several raised minute tubercles, about the size of an ordinary pin's head, and one-eighth of an inch in diameter; these were red in colour, and appeared to consist of congested fibrous tissue beneath the mucous membrane, and were not true tubercle.

The rectum was intensely congested, and covered with a whitish layer of secretion, which was firmly adherent to the mucous membrane. This diphtheritic membrane consisted of granules, of mucous cells, of columnar epithelium, some imperfect granule cells, and slightly fibrillated secretion.

Some of the mesenteric glands contained strumous matter. The liver was slightly fatty, 3lbs. in weight; there was a large calculus of cholesterine in the gall-bladder. The spleen corpuscles were large.

The ureters were much distended; the kidneys were diseased from obstruction; and their pelves distended; their cortical structure was full of white deposit, and there were some points of suppuration. The left kidney was large, and contained small abscesses. The bladder was distended; at the orifice of the urethra was a vascular growth. The ovarian cyst was simple; the fallopian tube extended upon it, and was filled with strumous deposit; the ovaries were obliterated; the ovarian cyst contained purulent serum and fibrin. The uterus and its cervix were much elongated.

The fatal termination in this severe case was much hastened by the ulcerated state of the colon; there was strumous disease of the abdomen and of the lungs; and it is probable, that the exhaustion consequent on the suppuration of the ovarian cyst led to the rapid development of strumous disease. It is important to guard against each fresh accession of diseased action, because the changes are of a rapidly degenerative character, and react upon the original disease.

*Discharge of mucus from the rectum.**—The discharge of mucus in cases of catarrhal diarrhoea has already been referred to, but this flux sometimes follows as a sequence of acute dysentery; and in other instances it is set up and perpetuated by the presence of local disease of the rectum, as by hæmorrhoids, polypoid growths, &c. In irritation of the colon, such as occurs in diarrhoea, mucus is often passed with the ordinary fæcal discharges; and in dysentery, the evacuations consist occasionally of simple mucus, with blood; it is to more chronic discharges that we here refer. The discharge may be very frequent, occurring not only with defecation, but at other periods; and it is especially present when there is local disease of the rectum, but is not limited to these states. Secondly, the mucus may be in shreds; but this kind of discharge we have not found except in acute diseases; and, thirdly, it may constitute casts, more or less complete, of a portion of the intestine. These are found to be several inches or a foot in length, and consist of mucus-cells, in a tenacious albuminous basis; but it is only in those cases in which shreds of mucus are discharged

* See some interesting remarks on this subject, by Dr. A. Clark, in 'Lancet,' December, 1859.

during acute disease of the intestine that any fibrillation of the product is observed; the patient not unfrequently regards the mucus cast in chronic disease as an intestinal worm.

The patients who suffer from this chronic discharge of mucus are generally in a cachetic state, pale, and more or less emaciated; the bowels are irregular, sometimes confined, at other times affected with diarrhœa; and there are occasionally intervals of several weeks or months between these mucous discharges of flakes and casts. During their passage a sense of distress and faintness, and even actual syncope may be produced, with severe colicky pain; the pulse is compressible, the tongue may be clean or furred, and the appetite uncertain; the mind is generally irritable or dejected, and sometimes almost melancholic.

In those instances which are not produced by disease of the rectum or of the sigmoid flexure, as hæmorrhoids, polypus, &c., nor are the sequence of acute dysentery, we have generally found, either that long-continued congestion of the vena portæ, or irritation of the urino-genital organs has produced or at least perpetuated the disorder. In young women, painful menstruation and ovarian disease may thus be the cause; in men, prostatic disease, calculus, &c.

The disease is a remedial one, and the prognosis may be favorable when the irritating causes can be removed, when a patient will submit to well-regulated dietetic regimen and careful habits of life, and when, for a sufficient time, he will persist in the use of proper medicinal measures.

Where disease of the rectum is present, and local and general means are unavailing, the help of the surgeon may be required to remove an irritating growth or hæmorrhoid. If the latter be present, the bowels should be regularly acted upon by the confection of senna and of black pepper; the gall ointment may be locally applied or an astringent wash used; and at the same time the diet must be carefully regulated and hepatic congestion avoided. It is, however, to the chronic form of mucous discharge that we here especially refer, and in this, careful attention to the diet and to the state of the skin is especially needed; a nourishing but unstimulating diet should be taken, and, if possible, stimulants altogether avoided. Although mercurial

medicines, in aperient or in very small doses, sometimes afford considerable temporary relief, we have often found the distress greatly increased by them; and we have seen greater benefit accrue from the use of nitro-hydrochloric acid with henbane and vegetable infusions, as of the calumba or cascarilla. If the bowels be loose, the krameria, with ipecacuanha and tincture of catechu, may be used, or the quinine with Dover's powder. When aperients are necessary, a small dose of colocynth may be given with Dover's powder, or of castor oil with tincture of rhubarb, but aloes and the more powerful purgatives are injurious. Astringent injections are sometimes of service, but are unavailing if portal congestion or pelvic irritation continue. In dysmenorrhœa and ovarian disease, absolute rest, at least for a time, is necessary, and the avoidance of tight corsets is indispensable. Dr. Clark recommends astringents, as alum, the pernitrate and the sesquichloride of iron, &c.

CHAPTER XI.

ON TYPHOID DISEASE OF THE INTESTINE.

IN typhoid fever diseased action of a special kind takes place in the glands of the intestine. This state passes through very definite conditions, and has been described by Rokitansky as the typhoid process. Dr. Jenner and others have pointed out the essential difference in the signs of typhoid and typhus fever, but it is only in the former disease that we find this abnormal condition of the intestine. The glands especially affected are those at the lower part of the ileum, namely, Peyer's or aggregate glands; the solitary also become involved, and the mesenteric are enlarged, congested, and swollen. The questions naturally arise, In what does this state consist? is it a necessary sign of fever? what are the indications of the typhoid state of the glands? and what is the course which the disease pursues?

The intestinal disease consists in the effusion of an abnormal product into the substance of the glands just mentioned, and the exudation effused from the capillary vessels is composed of a blastema, which undergoes but little development, and consists of an immense aggregation of granules, and of some large cells containing nuclei.

Soon after the commencement of the fever the glands are swollen and enlarged, and the mucous membrane becomes more vascular than usual; as the fever advances the glands are raised sometimes two or three lines above the surface of the membrane; about the fourteenth day of fever the product either becomes absorbed, or ulceration takes place, or the gland sloughs; a few days later the slough is found to have separated,

and an irregular ulcer occupies its position ; the muscular coat is exposed, and the margin of the ulcer is ragged and congested. If the patient do well this ulceration, of greater or less extent, gradually heals, a cicatrix is formed, and the health is slowly restored ; the convalescence extends over several weeks, and is interrupted, it may be, by relapses consequent on this condition of the intestine. The glands nearest to the ileo-colic valve are those which are most severely affected ; and sometimes the whole valve itself is converted into a slough, and the disease extends to the glands in the cæcum.

The mesenteric glands slowly assume their normal condition ; but, in some instances, the hyperæmia thus induced tends to the effusion of unorganizable strumous product.

In the examinations of the intestines after death from fever, we find the process of glandular disease in various stages in the same subject ; the glands may be merely swollen and raised, or the sloughing process may have commenced in small patches, or the whole glands may be converted into sloughs, which are partially detached, and stained by fæces ; the glands nearest to the cæcum may be in this latter state, whilst others further removed from that part are, in earlier stages of the same process, either beginning to slough or merely swollen and raised. In some cases, when death has taken place several months after fever, we have found cicatrices, without ulceration ; the disease had been cured. A very interesting case of this kind occurred at Guy's Hospital under the care of Dr. Gull. A young man was admitted, having the ordinary symptoms of typhoid fever, with the indications of ulceration of the intestine ; he appeared to convalesce favorably, but about three months afterwards he was seized with typhus, and died in a few days. On inspection, there was no injection of the mucous membrane of the ileum, but only cicatrices.

As a consequence of this diseased action the whole of the mucous membrane and even the deeper tissues become inflamed ; the intestine, from its enfeebled muscular power, readily yields to any distending force ; the peritoneum is sometimes injected, and presents in many cases a delicate, fibrinous exudation upon it ; again, the mucous coat often becomes ulcerated, so also the muscular, till at last only the semi-transparent peritoneum is

left; this also in many instances sloughs, and a minute opening takes place in the peritoneum, leading to rapidly fatal peritonitis; and even when the peritoneum is not actually perforated, transudation is frequently found to take place, so as to produce acute disease of the serous membrane.

Strumous disease, although not a necessary sequence, frequently follows fever; thus low organized product may be effused into the mesenteric glands; tubercles may be deposited in the substance of the mucous membrane, and ulceration follow; strumous disease of the peritoneum may take place; and, lastly, phthisical disease of the lungs and other structures may supervene.

The indications of diseased intestine in fever are to be distinguished from those of the fever itself, namely, the dry and hot skin, the excited and depressed pulse, the parched lips, and injected, furred, or dry tongue, the excited or oppressed brain, and the presence of rose-coloured maculæ, &c.; thus the abdomen is full and rounded, and on pressure in the region of the cæcum a gurgling sound is produced, with more or less pain; diarrhœa is generally but not always present; for sometimes the bowels are confined, even when the ulceration is severe; the evacuations are of loose consistency and of an ochre colour, and often contain blood; the pulse is compressible, the tongue is red and injected, and it becomes dry or cracked; there is often also a circumscribed flush on one or other cheek; the brain, too, is more depressed than in other cases. If perforation takes place there is sudden severe pain in the abdomen with collapse, and death then generally ensues in five to ten hours.

The symptoms of typhoid fever are sometimes so slight, even with existing ulceration of the intestine, that the patient is able to walk, and does not appear much enfeebled. I well remember a case of this kind, attending, and continuing to attend, as an out-patient at Guy's Hospital for three weeks, under the care of one of my colleagues. About the twenty-third day, a short time after admission into the hospital, when I first saw him, perforation of the intestine had taken place into the peritoneal cavity, and death quickly followed; and in other cases, when the severity of the fever has passed, and the patient has apparently begun to convalesce, having regained power and mental

energy, after some indiscretion of diet, or an attempt to move from the bed, perforation takes place, and the bright, beaming hope of returning health is lost in the terrible foreshadowing of speedy death. These are painful cases, trying to the physician, who has encouraged the hopes of the patient and his friends, and still more to those who are thus deprived of kindred. Perforation takes place from the third to the sixth week, and until this period is passed the greatest care should be used in allowing changes of food, or increased muscular movements; for among the fatal cases of typhoid fever a large majority die from this cause.

The hæmorrhage from the intestine in fever may be exceedingly severe; it may lead to a fatal result, or be the cause of lingering illness; or be of a critical character, and issue in speedy recovery. In Case C of typhus, fatal hæmorrhage took place from the stomach.

The general symptoms and treatment of fever, and the question whether there be any essential difference between typhus and typhoid fever, are not within the sphere of this work; the able manner in which they are discussed by Drs. Stewart, Jenner, Wilks, Peacock, &c., and earlier by Bretonneau, Louis, Broussais, Bouillard, Chomel, Christison, &c., render it unnecessary, and I must refer to their treatises.

Treatment.—The question may, and has been raised, how far diarrhœa is beneficial, and whether we ought at once to check it? We may be assured that purgatives are injurious, especially those of an active or drastic character. The disease of the intestine has been often aggravated by the injudicious administration, in the early state, of jalap and scammony, of senna, and the like; so also mercurials tend not only to increase the subsequent depression, but to aggravate the ulcerative action. It is often beneficial to act on the bowels and on the liver by a mild mercurial purge, where there is disorder of the abdominal viscera, but all irritation of a prolonged kind must be avoided. Where the diarrhœa is continued, it is well to check it by enemata of starch, or by the administration of chalk with opium, or vegetable astringents; but, in reference to the use of opium, it is well to be on our guard, lest the cerebral oppression be increased and a state of coma be induced. When hæmorrhage takes place,

the acetate of lead is sometimes of service, or vegetable astringents, as kino, catechu, logwood, &c.

We must strongly urge extreme caution in the return to solid and highly nutritious food; and equally important is it that, during the severity of the fever, and for many subsequent days, no muscular exertion should be attempted, but absolute rest in the recumbent position maintained. The attention to these means would have saved many valuable lives; and few diseased conditions require such constant watchfulness in the maintenance of rest, and the frequent administration of mild nutritious aliments.

Ammonia and serpentary may be given to stimulate the heart's action; and in some, quinine proves of service, especially where there is any sign of miasmatic influence. The judicious administration of wine in fever is one of the most difficult questions in practice; many are benefited by it, whilst others appear to do well without it; a failing circulation, especially in advanced life, as shown by a compressible character of the pulse and a feeble action of the heart, require its free use; other indications for the employment of stimulants are dryness of the tongue, prostration of strength, a dusky condition of the skin, the presence of petechiæ, &c. In many cases, if too long postponed, patients die, however freely wine may afterwards be given; and, on the contrary, I have witnessed in others the reduction and entire cessation of the administration of wine and ardent spirits followed by moisture of the tongue and the abatement of all the febrile symptoms.

CASE CLXXVIII.—*Fever. Peritonitis.*—Henry H—, æt. 23, died on the twenty-third day of fever; and on the day before death he had symptoms of peritonitis; he had previously had syphilis, and was in a cachetic state.

On inspection, there was acute peritonitis; but no perforation could be found; there was very extensive ulceration of the ileum and cæcum, affecting both the solitary and Peyer's glands; in one or two places the muscular coat was also destroyed. The liver, spleen, and kidneys, were healthy.

In this instance, acute peritonitis was set up, although complete perforation had not taken place; in some parts, however, the serous membrane only was left entire; and there can be

little doubt that transudation had resulted in the acute serous inflammation.

CASE CLXXIX.—*Fever. Perforation of Intestine in the Seventh Week.*—Joseph C—, æt. 17, was admitted, August 18th, 1858, and died September 1st. He had been residing at Woolwich, and six weeks before admission had suffered from fever, with much diarrhœa. The day before he was brought to Guy's he was said to have had a discharge of blood from the bowels. On admission, he was pale, emaciated, and almost pulseless; scarcely any complaint of pain was made, but he had an anxious and distressed countenance; the abdomen was moderately distended; he rallied slightly, but sank a week afterwards.

Inspection on 2nd September. *Chest.*—The pleura was healthy; the lungs were pale, and distended; the *heart* was healthy. *Abdomen.*—The intestines were distended; the peritoneum contained about two pints of offensive pus, which was collected in the depending parts; there was scarcely any adhesive lymph, and no adhesions; but the general peritoneal surface was slightly opaque. About eight inches from the cæcum there was a sloughing opening into the ileum, capable of admitting the tip of the little finger; and a second opening of rather smaller size was situated nearer to the cæcum. On the mucous surface of the ileum at its lower part, were the remains of ulcers; many of them cicatrizing, their edges were smooth and grey; the floor of the ulcers were formed in several instances by smooth submuscular areolar tissue on the peritoneal surface; these remains of ulcers, covered nearly the whole of this part of the intestine. There were also several ulcers in the cæcum; the rest of the intestine was healthy. The *liver* was discoloured on its surface, but it was otherwise healthy; the *spleen* was slightly enlarged; the *kidneys* were healthy.

The comparative absence of pain in this case was explained by the great prostration from which the patient suffered; the great care required after typhoid ulceration of intestine is shown in this instance; for six weeks had elapsed, and many of the ulcers had healed before fatal rupture of the intestine and peritonitis took place. It is probable that injudicious diet or increased muscular exertion had led to the perforation of the ileum.

CASE CLXXX.—*Perforation of the Ileum. Typhoid Fever. Cyst connected with the Œsophagus.*—John C—, æt. 28, was admitted in a dying state into Guy's Hospital. He stated that he was well till the previous evening, when he was seized with acute pain in the abdomen; and on admission he had the symptoms of peritonitis. He died in a few hours.

On inspection, the body was found to be well nourished. The abdomen was distended; the whole surface of the intestines was covered with fæcal matter,

and the end of the ileum was perforated; the lower part of the ileum was ulcerated, as in typhoid fever, and one of these ulcers had given way; there was deposit in Peyer's glands, and the mesenteric glands were enlarged and softened. The liver was healthy, so also the kidneys. The spleen was large, soft, and of a dark colour. There was a cyst in the upper part of the œsophagus at its posterior part, about the size of a pigeon's egg; it consisted of a distension of the mucous membrane, and was situated beneath the muscular coat; *i. e.*, the muscular coat did not cover it; it was filled with thick mucus.

The patient seemed to have been so far convalescent from a mild attack of fever as to consider himself well, when fatal perforation of the ileum took place. The appearance of the intestine was precisely that found in typhoid fever, nor was there any indication that the ulceration was of a strumous character.

CHAPTER XII.

ON COLIC.

By the term colic we mean a severe twisting pain in the abdomen, about the region of the umbilicus, without inflammatory action, generally with constipation, but sometimes with looseness of the bowels and vomiting. Internal strangulation of the intestine and intussusception are considered as more aggravated and severe forms of colic; but the former is often in its early stages altogether free from pain, until distension and violent peristaltic action set up inflammation, which involves all the coats of the intestines; unless, then, we consider some forms of colic to be free from pain, in some of its stages, we cannot regard fatal obstruction, arising from displacement and internal strangulation, as a form of the disease.

Dr. Copland divides colic arising from functional disorder of the bowels into four classes :—

1. Flatulent, nervous, or spasmodic colic.
2. Colic from an injurious character of the food.
3. Colic from morbid secretion, or retained excretions.
4. Colic from lead.

This division appears to be a just and useful one; for they indicate widely different conditions, which require different treatment.

In *flatulent* colic the intestines become distended with flatus; severe twisting pain comes on round the region of the umbilicus; if the pain be severe, the patient becomes cold, and a clammy sweat breaks out; the pain extends to the back, and the severity of it is relieved by its changing position, but especially by eructation, or by the discharge of flatus from the rectum; during the

severity of the pain, the pulse becomes depressed, feeble and irregular.

In *nervous and spasmodic colic* there is less distension of the abdomen, which may be slightly tympanitic; the pain extends to each side and to the back, and is sometimes situated across the chest. The removal of pain from one part to another, and especially the discharge of flatus, as in the flatulent colic, afford relief. The abdomen is, however, tolerant of pressure, and pain is occasionally relieved by this means. There is, also, an anxious expression of countenance, with coldness and clamminess of the surface, and with depression of the pulse, if the pain be severe. The collapse has even been mistaken for ruptured intestine, so complete may be the prostration; in colic, however, the collapse entirely disappears in a few hours; in perforation, it soon terminates in death. The tongue is not generally affected, and it may be clean or furred; the evacuations from the bowels also may be as in health, but in most cases they will be found to deviate from their normal condition. In the nervous colic of hysteria, the urine is abundant and limpid. In the spasmodic colic of gout the urine contains an excess of lithic acid, and may be turbid and scanty.

Causes.—Flatulent colic is observed in nervous and hysterical subjects, and is produced by the rapid evolution of gases from the contents of the alimentary canal, whilst in some cases it appears to arise from change in the secretions of the mucous membrane itself. In the intemperate, and in gouty patients, the chylipoietic viscera are often in a congested, and morbidly excited state; and in these cases a very trifling cause will produce intense colic.

In patients reduced by exhausting diseases, by loss of blood, by the too long continuance of farinaceous and fluid aliments, as also by over lactation, we find that colic of this kind is readily induced.

The exciting causes are alarm and affright, exposure to cold and wet, especially of the lower extremities, food not in itself indigestible, but taken when the powers of digestion are diminished, either from an enfeebled condition, or from the state of the nervous system.

Diagnosis.—There are several conditions with which this

functional colic may be mistaken, and which are important to remember.

Perforation of intestine is generally known by the intensity of the collapse; it is exceedingly unusual for collapse at all approaching in severity to that produced by ruptured intestines to arise from functional colic, but that is sometimes the case. During the passage of a calculus from the gall-bladder or from the kidney, intense pain is produced; but the position of the pain, the character of the vomiting, and in the latter, the pain and retraction of the testicle, and blood in the urine, enable us to distinguish these diseases from ordinary colic.

In disease of the spine, severe and sometimes intense pain is produced in the abdominal parietes, but this pain has less of the twisting pain of colic, and may be traced in the course of the spinal nerves.

Aneurism of the abdominal vessels and abdominal tumours are sometimes the cause of intense suffering, but the constancy of the pain, and the presence of tumour with other signs afford easy diagnostic marks of their true character.

In peritonitis there is exquisite tenderness of the abdomen, whilst in colic the pain rarely amounts to more than a diffused soreness, and pressure can often be borne.

In strumous and chronic peritonitis flatulent distension of the abdomen is associated with soreness or tenderness, less severe than in ordinary peritonitis; and these conditions may, in the *early stages*, be mistaken for simple colic; this is important, because by an over active plan of treatment disease may be accelerated; and afterwards, when the intestines are matted together, and attacks of peritonitis are set up, the pain and tenderness come on in severe paroxysms. It is only in the early conditions of this disease, and especially in young people of nervous and excitable temperament, that there is liability to such mistake.

The distinguishing marks between colic and hernia need not be dwelt upon; for the presence of an external tumour, with constipation and stercoraceous vomiting, can only be mistaken for simple colic by great carelessness; and in every case of colic, before severe vomiting has come on, it is well *always* to examine the patient for hernia. In intussusception also,

intense recurrent pain in the abdomen of a twisting character is sometimes associated with diarrhoea, and I have seen the disease mistaken for colic produced by irritating substances; this idea was strengthened by the occurrence of the attacks of pain on the days when the patient was visited by his friends. Flatulent distension of the stomach is sometimes a severe and even *fatal* disease, but the distension and tympanitis are great, and the pain is constant. The intense pain arising after poisons, as mineral acids, &c., is associated with violent vomiting; but the suffering is especially experienced in the mouth and gullet, and has other characteristic symptoms.

Our *prognosis* in colic of this kind is generally favorable, but it must be much more guarded when we have had evidence of previously existing disease, and where the collapse is great, and again in some cases of gout.

Treatment.—The ordinary treatment in colic, often before the patient is seen by a practitioner, is to administer some hot brandy and water; and if the disease be of the simple kind, which we have described, the patient may be thereby relieved; but in peritonitis, in hernia, in perforated intestine, nothing can be worse, for it takes from the patient the chance of recovery. Opium, gr. i—ij, and laudanum, are the most useful remedies in severe functional colic, and may be given either alone or with ether; chloroform in doses of ℥v—xv, with or without camphor, gr. j—v; chloric ether also, and the salts of morphia, may be advantageously administered. Warmth should be applied to the abdomen by means of hot water, hot flannels, poppy fomentation, &c., and sometimes a mustard poultice and a hot flannel sprinkled with turpentine, may be employed to relieve severe pain.

When the bowels are inactive, and when no indications of acute disease, hernia, or internal obstruction exist, a warm saline purge may be given, but some practitioners prefer gr. v—x of calomel, with gr. j or ij of opium, or colocynth with henbane, or castor oil with tincture of rhubarb and of opium. Again, enemata are sometimes of much service in emptying the colon, and thus may entirely relieve the disease; castor oil and oil of turpentine, colocynth and rue, may each be in this manner employed.

If the attacks be less severe, but repeated, and if the patient

be exhausted and anæmic, steel and quinine may be combined with henbane and with mild aperients. At the same time many of the vegetable bitters—calumba, cascarilla, gentian—may be prescribed with the aromatic spirit of ammonia, with henbane, with the carbonated alkalies of soda, potash, or magnesia, &c. In hysterical subjects the compound galbanum and aloes and myrrh pill may be used, and some practitioners employ valerian, musk, castor, stambul, and the essential oils, to relieve the painful and flatulent distension of the abdomen.

The diet should be of a form easily digestible, but sufficiently *varied*, and neither bulky nor entirely of a fluid kind.

When a gorged state of the portal system exists, from an excess of aliment and of stimulants, it is well to abstain from alcohol in every form; but this abstinence is the more difficult to attain, because the colic is itself often relieved by fresh doses of ardent spirits. So also in gout, an excess of animal food and of stimulants aggravates the disease, but the patient may be so enfeebled as to require the continuance of stimulants in a well-regulated manner; and in instances of colic with exhaustion, from over lactation, loss of blood, great mental alarm, &c., alcoholic stimulants are of great value.

2. Colic arising from the *injurious character of the food*. This disease has many symptoms in common with the colic just described; but its cause is different, and also its mode of relief.

Severe pain comes on in the region of the scrobiculus cordis and umbilicus, sometimes with flatulent distension, two or three hours after eating. Vomiting occasionally supervenes, and it may be, if the food is of an injurious character, either in itself or from the idiosyncrasy of the patient, that diarrhœa is set up. The tongue is whitish and furred, or enlarged, and red papillæ are observed through the fur, or it is injected at the tip and edges. The pain is followed by a soreness of the abdomen, which may persist for several hours or days. The pulse is compressible, and the respiration is less free than normal. This condition may pass into that of enteritis, or of diarrhœa; or, after vomiting, and disturbed unhealthy evacuations from the bowels, the patient may be restored to health.

This form of colic is often associated with disturbance of the cerebral function, and severe pain in the head, dimness of sight,

irritability of temper come on ; or it may set up disturbance of the skin, producing urticaria and roseola, and in children, strophulus and other lichenous eruption.

If the injurious character of diet be persisted in, the colic may cease, but other conditions consequent on general impaired nutrition may be set up.

The exciting causes are salads, cold drinks, acid and fermenting wines, raw fruit, especially stone fruit, mussels, and so-called shell-fish, the imperfect mastication of food, either from the inefficient means or that time is not allowed for the partaking of it ; this is especially the case in those who are actively engaged in business, and amongst those who from choice or necessity postpone the meal till the frame is almost exhausted. Severe colic of this kind is sometimes produced by mushrooms, especially where other forms than the edible agaricus are taken, and dangerous and alarming symptoms may follow.*

In other cases the diet may be perfectly right in itself, but improperly administered ; thus the most severe colic may be produced by giving cold milk to young children. Sudden prostration of strength, pain, sunken eye, vomiting, and afterwards diarrhoea are produced, and the motions indicate the undissolved state of the food taken.

In the *diagnosis* of these cases, equal care is necessary as in flatulent colic, for the same conditions may mislead in each ; it is well also to remember that hernia, perforation of the intestine, peritonitis, intussusception, and enteritis may produce many of the symptoms of this form of colic. Ordinary care will in most cases enable us to detect external hernia ; the pain of peritonitis and that from perforated intestine is more severe in kind, and different in character from that of ordinary colic. Enteritis and intussusception may follow as the consequence of intestinal irritation of the kind just described, but these symptoms are of greater duration than those of simple colic ; and when due to the administration of irritating poisons, the examination of the vomited matter is an essential element in forming a correct opinion as to the nature of the case.

Treatment.—If vomiting have come on, and irritating matters

* Taylor on 'Poisons.' Christison on 'Poisons.'

have already been freely ejected, the symptoms of disease often spontaneously subside, or soothing demulcents may be given; but if pain and nausea continue, an emetic is of service. Action from the bowels should be ensured by saline aperient medicines, as the carbonate of magnesia, the sulphate of potash, the tartrate of soda, or by a free mercurial purge; and demulcents, as arrow-root, milk, rice, mutton or veal broth, &c., should constitute the only diet; the administration of saline medicines, with antispasmodics, sedatives, and anodynes may be necessary for a short time, and opium may be required to check the irritated action which has been set up, but opium should not be given whilst irritating substances continue to disturb the intestine.

3. Colic from retained secretions and morbid excretions.

When severe pain comes on with diarrhoea and with dark bilious evacuations, a state of disease is produced which is closely allied to the bilious diarrhoea and English cholera, which we have previously noticed. The severe pain in the region of the umbilicus may be associated with violent vomiting and purging, without being caused by any impropriety of diet. The patient sometimes becomes prostrate, the motions fluid, the surface cold, the pulse compressible, and in a very short time he is brought to extreme collapse, resembling Asiatic cholera. This in its most severe form constitutes the English cholera that is found to prevail each autumn in our own country. But there are less degrees of this condition; the vomiting, pain, and purging may be more moderate, the tongue furred and injected, whilst the prostration is less. Again, in other instances, severe pain in the abdomen of the character of colic is present, without any purging or vomiting, but with a sallow complexion, with furred tongue, pain in the head, oppression of the mind, and impaired physical energies.

In some cases the prostration is so severe that the patient succumbs; but more generally, I may say in most cases, the symptoms gradually subside, and are followed by speedy recovery.

Causes.—In the autumnal season there is much greater liability to this disease, on account of the sudden variations in temperature to which persons are often exposed. The exhalations from decaying animal and vegetable produce, the effluvia

from drains, also induce this form of colic. In miasmatic districts, and in damp localities, there is still greater liability to this state, and so great may be the predisposition, that a very slight excitement is sufficient to set up the disease.

The most frequent cause, however, is disturbance in the function of the liver, whether from intemperance, indiscretion in diet, mental disquietude, &c. ; and whatever leads to congestion of the portal system tends to induce colic upon slight exciting causes. Attacks of this kind are often designated "spasms."

In infants it is exceedingly common to have colic from retained and from morbid secretions ; pain is produced, as shown by the drawing up of the lower extremities, the cry is almost incessant, there are green or watery evacuations, containing portions of coagulated milk or undigested food, the countenance is anxious and distressed, and the sleep is disturbed. If this condition continue, it extends so as to affect the mucous membrane of the stomach, and is then associated with violent vomiting, so as to constitute gastro-enteritis ; rapid prostration may ensue, and death quickly follow, or more slow muco-enteritis or intussusception may supervene ; and in some older children a less serious, but a troublesome disease, prolapsus ani, is occasionally produced.

In the diagnosis, the remarks made in reference to the other descriptions of colic are equally applicable ; and in the severer forms the disease approaches in character to that of cholera.

Our prognosis must be a guarded one, for although most cases recover, still in some, especially in infants, an untoward result follows, and the patient becomes perfectly prostrate and dies.

Treatment.—It must be remembered that the effect of the vomiting and purging in these cases is to remove the offending matters from the alimentary canal, so that many cases, if left to themselves, recover. In milder cases the pain, the vomiting, and purging are entirely removed by the administration of arrowroot, or by the injection into the rectum of thin starch. If offending substances and secretions are retained, castor oil, with tincture of rhubarb, and with opium, afford great relief, repeated as need be ; and it is in this condition, antecedent to the aggravated forms of Asiatic cholera, that we may expect and

derive benefit from the plan of treatment recommended by Dr. Johnson. Some administer magnesia with good effect, calcined or carbonate, with a little conium or henbane; and grey powder, with Dover's powder, in gr. v doses, or calomel with opium, may be given several times during the day, so as to remove the abnormal contents of the intestine, and to check the pain of the colic. If, however, the pain and diarrhœa continue, it is well to give absorbent alkaline medicines, with astringents, as chalk with catechu and opium, or kino, krameria, logwood, tormentilla, &c., and to repeat the starch injections, or injections of oak bark, &c.

In the subsequent prostration, mineral acids, the sulphuric, nitric, hydrochloric acids, with vegetable tonics, are of great service in restoring tone to the mucous membrane. The sulphuric acid has been much used in this form of diarrhœa, and we have already alluded to its use. The secretions from the mucous membrane of the small and large intestine are of an alkaline character, and when the membrane is irritated are poured out in greater quantity, forming an unusually thick covering to the membrane; in this state the mineral acids correct their character by their astringent effect on the capillaries, checking the further secretion of watery mucus, and they assist the removal of that already formed. The solution of potash, and the alkalis generally, have a soothing influence upon the mucous membrane of the alimentary canal, and I think are of greater service than acids in the early stage of colic and diarrhœa from offending secretions.

If there be persistent pain, the application of leeches to the abdominal parieties, or to the anus, and warm applications to the surface, afford relief.

Food should be very sparingly administered, and only of the amylaceous and most bland form, as arrowroot, rice, tapioca, &c., veal or chicken broth; if the strength fail, we must add brandy, or port wine, &c.

4. *Lead colic.*—Till attention was drawn to the subject of lead poisoning, the colic arising in the wine and cider districts was attributed entirely to the character of the fluids drunk; this has still been shown to be in a great measure the case, and since the observations of Sir George Baker, the effects

of poisoning by lead in its several forms have been completely verified, and we are now able to distinguish the effects of lead poisoning with, perhaps, more ease than almost any other form of disease.

The patient exposed to the influence of lead becomes of a sallow and anæmic aspect, his muscular development is diminished, and his mental capabilities are somewhat enfeebled; if colic come on, he experiences severe pain in the abdomen, at first moderate, but afterwards becoming intense and of a twisting and grinding character about the umbilicus; the abdomen is contracted, and the patient experiences relief by firmly compressing the abdomen with his hands, or even across a chair; the bowels are obstinately constipated, the abdomen is neither tender nor hot, but hard and contracted; nor is there generally any vomiting, but the patient writhes with the severity of the pain; the tongue may be clean or furred, the pulse is feeble, but not increased in frequency, and the urine is pale. After some hours the severity of the pain subsides, but may again return during the next night, or after taking food. The severe colic is sometimes accompanied by cerebral disturbance, but this is a rare occurrence, although severe cephalalgia or epilepsy may precede or follow colic, as another of the effects of the lead poisoning, as also paralysis of the extensor muscles of the forearm; or the colic may be associated with severe cramps and pains in the extremities; the constipation sometimes gives place to diarrhœa, but still the pain continues, or rather severe soreness, occasionally aggravated into intense suffering. On examining the gums we find along the edge a grey line, composed of minute particles of sulphuret of lead, from the mutual decomposition of the lead permeating the capillaries and the sulphocyanide of the saliva. This is itself a very distinctive sign of lead poisoning, and when the pain is unaccompanied by tenderness is sufficient to guide us to a correct diagnosis. It rarely occurs that lead colic alone terminates fatally, unless associated with other diseases or other conditions of lead poisoning. In a case to which I have referred in diseases of the stomach, lead colic was associated with chronic ulceration of the stomach, which led to perforation and fatal result. We sometimes find the paralysis of the hands or wrists, and epilepsy,

coincident with the colic ; it is unusual to have paralysis of the ankles, but such a case I have seen.

The proximate cause of lead colic is not known, whether it arises from irregular peristaltic action, or paralysis of one part, and spasmodic contraction of another. In those cases which I have examined, and in others recorded, no abnormal appearance was found in the intestine. The manner in which the lead enters the system is, in some cases, very obscure, but generally it is sufficiently manifest. Drinking fluids from leaden vessels which are not covered with any protective carbonate, &c., and acid drinks, as cider, &c., from leaden vessels, are the common modes of its introduction ; but lead colic is most frequently observed in plumbers, painters, type-founders, &c., men who are constantly employed in handling lead, and who breathe an atmosphere contaminated with minute particles of it. It appears probable that in the mixture and using of paints containing lead there is still greater liability to its absorption, the volatile oil containing minute particles of the metal, and thus its ready inhalation is effected. In many instances the want of proper cleanliness in washing the hands before taking food, and in changing the clothes, very much aggravates the liability to poisoning by lead. It is sometimes, however, difficult to ascertain how lead has entered the system. Dr. Addison mentions a publican who was thus poisoned by drinking in the morning, as his first draught, the ale which had remained in the leaden pipe during the night. Several instances have been known where lead was found in the snuff which the patient had been in the habit of taking. It has sometimes been produced by the medicinal use of acetate of lead ; but Dr. Thompson has shown that there is less liability to this effect being produced when the lead is combined with opium, or given with dilute acetic acid.

The diagnosis of lead colic is sufficiently clear when ordinary caution is used, the lead line along the gums, with pain relieved by pressure, and the contracted abdomen, distinguish the disease ; but, as before mentioned, it may be associated with chronic ulcer of the stomach, with hernia, &c., which obscure the diagnosis, and may lead to a fatal result.

In lead colic alone we may give, especially in the earlier attacks, a favorable prognosis.

Treatment.—The indications of treatment appear to be sufficiently plain in this disease—to relieve the pain, to act on the bowels, and to remove lead from the system. For the relief of the pain, opium or chloroform is the best remedy, and may be administered freely; to act on the bowels, croton oil with opium, or calomel with opium, or castor oil and laudanum, or the sulphate of magnesia with compound infusion of roses and henbane may be used; or we may administer injections of castor oil or colocynth; warmth should be applied at the same time to the abdomen.

In relation to the subsequent treatment, we should not be content with the subsidence of the colic, as long as the patient retains his sallow and anæmic aspect, and has a lead line along the gums. Iodide of potassium has been used, and it has been found that the urine contained lead during its administration; this I have often attempted, but unsuccessfully, to verify. Considerable benefit has been found in dropped hand from rubbing iodine ointment into the paralysed parts,* and still more by the use of electricity and galvanism locally applied. An insulated water bath has been recommended, the patient in the bath being connected with one pole, the sides of the bath with the other. The lead is stated to be removed from the body of the patient, and deposited upon the walls of the bath; but I have not seen electricity applied in this manner; the only opportunity in which I have known galvanism used in colic, to excite the bowels to action, was in the case associated with gastric ulcer; the existence of gastric ulcer was not known, and fatal peritonitis followed. Warm baths, perfect cleanliness, bracing air, and preparations of steel, after the removal of lead, are of great service; but a considerable time is required for the complete removal of the poison from the system.

The prophylactic treatment is an exceedingly important consideration to those employed in the use of lead. The importance of perfect cleanliness, of changing the clothes, of not partaking of the meals in the workshop, are now generally acknowledged, although they are not acted upon as their cogency demands.

A drink containing dilute sulphuric acid is mentioned by Dr. Watson, as having been used by Mr. Benson, in lead works, with very great advantage.

* *Medical Times and Gazette,* May 1857.

CHAPTER XIII.

ON CONSTIPATION.

WASTE and repair are necessarily connected with the performance of every function of the human body ; and the various excretory organs are the channels by which the materials of waste are separated as substances no longer of any benefit, and the retention of which becomes increasingly detrimental to the whole economy.

The large intestine may be looked upon as a very important excretory organ ; and the removal of its contents is as necessary for the continuance of human life, as the separation of carbonic acid from the lungs in ordinary respiration.

The colon is well adapted for the purposes of excretion, and by its arrangement serves as a reservoir ; and it thus allows an occasional, rather than a continuous, separation of its contents.

But in this periodical movement of the intestinal canal there is great difference ; and the variation within the bounds of health is much greater than is usually supposed ; with some, and perhaps by far the larger number, an action of the bowels takes place once every day, or it may be two or three times, although either may be the state of health ; on the contrary, with others, it may be that every second or third day is the normal condition ; and the usual period may be even extended to every fourth or seventh day. This variation must be borne in mind, otherwise, in the attempt to produce what is considered beneficial, an abnormal condition may be set up, and comfort and health lost in striving to bind all to the same universal law.

Much, however, may be acquired by habit ; regularity may be attained ; or inattention and want of care may induce a condition which will almost baffle subsequent exertions to eradicate. Premising that the healthy action of one is disease with another, we may define constipation to be a less frequent action from the bowels than is the healthy condition of each individual. Ordinary constipation arises from the insufficient contraction of the muscular coat of the intestine ; the canal becomes more and more distended, and with each increase in the circumference of the tube greater power is required to force onward its contents. I have sometimes observed a colon so enlarged by distension and loss of power, with obstinate constipation, that it has measured as much as twelve to fifteen inches in circumference ; the power required to propel the contents must have been enormous. And it appears probable that in this extreme distension, a state closely allied to paralysis of the muscular parietes is the result ; sometimes, however, this gradual distension is the effect, rather than the cause of paralysis.

A second effect of constipation is that the lateral pouches of the colon formed by the circular and longitudinal bands of muscular fibre, become more and more distended, and being thus filled out, their contents are removed from the central current, and become impacted, while the bowels act with some degree of regularity ; these impacted *faeces* may frequently be felt as tumours through the abdominal walls, alarming the patient, but disappearing under judicious treatment.

Pouches of the colon sometimes become of considerable size, and generally the circular fibres of the canal surround them ; but not unfrequently the circular fibres yield, and the mucous layer projects, covered only by the peritoneum, thus forming a mere elongated sac, filled with mucus, or more frequently with *faeces*. The orifices of these small sacs are bounded by the hypertrophied circular and longitudinal fibres, and their contents remain almost shut off from the intestinal canal. These pouches are the result of constipation, the muscular fibres become hypertrophied, but their efforts to propel onward the contents of the canal lead to these minute hernial protrusions.

I have most frequently observed pouches in connexion with the sigmoid flexure ; but they, probably, occur at any part where

the longitudinal fibres form a triple band rather than an uniform layer. In one case they were situated about every half-inch, forming a double row on each side of the colon. No muscular fibres could be detected in several of them, beyond the immediate vicinity of the mouth of the sac, but merely mucous membrane, submucous cellular tissue, with fat and peritoneum. These pouches do not appear to produce any symptom, nor do they lead to dangerous results.

CASE CLXXXI.—A remarkable case of this kind I observed in a patient, aged sixty-two, who died from cancerous disease of the liver and lungs, with bronchitis and emphysema. The sigmoid flexure and rectum were contracted, and presented numerous pouches, some of which were half an inch in length; they were arranged in two rows about one inch apart; these pouches consisted of mucous membrane and peritoneum; the circular muscular fibres were placed between the pouches, and the longitudinal fibres on either side, and both were hypertrophied. The pouches were filled with mucus and fæces. There was neither ulceration nor evidence of cicatrix, but it appeared that the constipated bowels to which the patient had been subject had led to unequal pressure and saccular distension, or herniæ of the mucous membrane. Appearances of this kind, though in less degree, are by no means uncommon in the colon, especially towards its termination.

Continued distension with solid contents alters the position of the colon; this is especially observed in the transverse colon, and in the sigmoid flexure; the convexity of the former becomes greatly increased, and the double curve of the latter is rendered more evident. The attachment of the great omentum, and the ready separability of its layers, are especially designed to allow of free distension of the transverse colon, but a continued pressure increases the curve, till at last it may form a large sigmoid flexure, reaching nearly to the brim of the pelvis. Increased curvature of the transverse colon is of common occurrence, but sometimes its malposition seems to be congenital; thus we have seen the descending colon passing downwards in close contact with the ascending, and then terminating in a transverse colon, which

was situated at the brim of the pelvis, thus connecting the descending colon with the sigmoid flexure.

The most important result arising from continued constipation is the retention within the blood, or the reabsorption into it of materials which are essentially excrementitious. The excrementitious portion of the bile is not removed, and the function of the liver is imperfectly performed; the blood of the whole portal system is rendered more or less impure; the complexion becomes changed, sallow, and muddy; the brain does not act with its wonted energy, and there is a manifest diminution in the elasticity of the whole mind and body.

The functions of other viscera become disordered, and the enlarged and distended colon mechanically interferes with the healthy action of adjoining viscera. The cæcum and ascending colon may press injuriously upon the ilio-hypogastric and genito-crural nerves, leading to severe neuralgic pain over the crest of the ileum or groin; and pain of this kind may be mistaken for rheumatism, lumbago, and sciatica, whilst it entirely disappears when mechanical pressure on the nerves has been removed. This pressure is, however, more frequently exerted on the left side by the sigmoid flexure; the veins of the lower extremity and the testicle or ovary becoming pressed upon, and œdema of the feet and varicose veins result; by the distended transverse colon the stomach is interfered with, and its movements to a certain extent crippled.

In reference to the *causes* of constipation, the *first* to be mentioned is original peculiarity of habit, or idiosyncrasy; that such peculiarity does exist cannot, I think, be doubted, although it must not be considered as disease in the same light as constipation arising from organic change.

A *second* cause is an abnormal condition of the abdominal walls. The contraction of the parietal muscles is an important aid in defecation, and their tonic contraction assists the peristaltic action of the intestines. The constipated condition of the bowels in diseased and fractured spine arises from this cause, namely, paralysis of the parietal muscles; but diminution of contractile power also arises from degeneration of the muscles themselves, and from excessive development of fat; and sometimes the contraction of these muscles is checked by pain, either

of a neuralgic character, or from local inflammation, as boils, fascial abscess, carbuncle, &c. Inactivity, or sedentary life, tends to produce constipation in the same manner. How different the condition when many hours are spent, day after day, in nearly the same position, from that of active muscular exertion ! Contrast the mechanic, where the whole frame is in constant movement, with the overworked sempstress ; the clerk, sitting for hours over the desk, with one engaged in active out-door occupation ; the professional or literary man, almost deprived of walking exercise, to another in the full enjoyment of it. The muscular exertion of walking, horse-riding, various athletic exercises, or other means by which the muscles of the abdominal walls are brought into play, are essentially necessary for sustaining good health.

A *third* cause of constipation arises from the alteration of the secretions poured into the large intestine. These secretions, or rather excretions, arise from the mucous membrane of the large intestine itself, from the small intestine, from the liver, and from the pancreas, and they undergo various changes ; thus a congested condition of the liver and of the portal system of veins, induces modification of the whole chylopoietic viscera, for the vena portæ receives its branches from the large and small intestine, stomach, &c. ; hence also a state of congestion of the liver not only checks the formation of bile, but it interferes with normal secretions from other parts, often diminishing them in quantity, and altering them in quality ; in this manner we have constipation from hepatic disturbance, and from the imtemperate use of alcoholic liquors ; thus also in jaundice, constipation is generally the result, the motions become clayey, white, and exceedingly offensive.

Diseases of the lungs and heart, which interfere with the free circulation of the blood, render the right side of the heart engorged ; and as a necessary consequence of this distension, the liver and the whole portal system are congested, the secretion from the mucous membrane becomes scanty, and constipation is the result. This constipation increases the original disease of the heart ; and the remark is often made by those who are the subjects of chronic disease of the lungs and heart, as chronic bronchitis, emphysema, asthma, and valvular disease of the

heart, that as soon as the bowels become confined, they experience increased discomfort.

A state which may be called chronic catarrh of the mucous membrane is sometimes induced from this congestion of the portal system, and constipation very frequently follows; but another cause of this altered secretion arises almost from an opposite cause, namely, a diminished supply of blood to the mucous membrane. The secretion is scanty, but from a different reason; in the former case, secretion is checked by engorgement of the vessels with blood; in the latter, by a diminished supply.

The various excretory organs are closely connected the one with the other. The excretions from the lungs, the skin, the kidneys, the alimentary canal, are intimately associated. Their nicely adjusted balance continues during health, but if one becomes greatly in excess, the others consequently, and almost in that proportion, suffer; thus excessive secretion from the skin diminishes secretion from other parts. The box of rhubarb pills is often carried by the pedestrian—and why? The muscular exercise and action of the abdominal muscles should induce increased action; and such would in many cases happen if the exercise were moderate; but if persisted in so as to induce free perspiration, with rapid molecular changes in the muscles, blood is actually withdrawn from the alimentary canal to the skin and muscles; the internal secretions become diminished, and constipation results. A similar condition is observed when excessive action of the kidneys carries off the aqueous portion of the blood too freely. The kidneys act less when the skin energetically performs its function; and on the contrary, when the warm air of summer is suddenly changed to a cold, chilly temperature, the action of the skin is checked, and increased renal secretion is induced. We have already alluded to this in our remarks on diarrhoea and dysentery; for the sudden interference with the action of the skin often induces those diseases; hence the autumnal diarrhoea, and the severe dysenteries of hot climates. Cerebral congestion, over-anxiety of mind, extreme mental occupation, act also in this manner, as well as more directly upon the nervous condition of the alimentary canal. There is increased circulation of blood in the brain, and less in

the abdomen ; for great excitement of the cerebrum is associated with diminished activity in the nerve of organic life.

Constipation is also induced by *general* anæmia, and loss of blood ; and very frequently in spanæmia or poverty of blood, as in the chlorosis of young women. The condition of the blood is here the primary cause of other secondary changes. There is inactivity or irregular muscular exertion, and the secretions are imperfect both in their character and quantity.

A *fourth* cause of constipation is a diseased state of the coats of the intestine itself. I have already alluded to the secretion from the mucous membrane, and especially refer here to the condition of the muscular layer, and to the nervous supply of the alimentary canal.

The muscular layer, in a state of health, contracts from slight direct stimulus upon the contents of the canal, but this contractile power is variously modified ; sometimes it is excessive, leading to the immediate expulsion of the contents, but more frequently it is inactive, leading to constipation. This inactivity may arise from the muscular coat having been unwisely excited to action by improper means—as by the injudicious use of purgatives, either from their habitual continuance or from their powerful character ; and the muscular coat is left in such a state that it will not contract from the normal stimulus, whilst the diminution of contractile power is increased by the constipation with which it is associated. The intestine becomes distended, the calibre increases, and the muscular fibre, which could easily propel the contents of a cylinder one to two inches in diameter, is unable to do so when the cylinder is increased to three or four inches in diameter, and the canal sixteen times its normal size. A state of actual paralysis of the muscular fibre of the intestine may be thus induced ; and in the same manner as the urinary bladder, if enormously distended, it is unable to expel its contents. Repeated doses of blue pill and black draught, of violent purgative medicines, of mercurials, &c., render the whole coat of the intestine in a relaxed and enfeebled condition ; the mucous membrane is debilitated, the muscular fibre is inactive, and partially paralysed ; not that I mean for a moment, that such remedies are not frequently attended with marked relief to existing morbid conditions ; but the continued use of them

leads to chronic disease, which is perpetuated, as well as induced by the remedy itself, although in some instances this is borne with apparent impunity.

Dr. Billing related to me an instance of a lady, who for thirty years took a grain of calomel every night; and a colleague of his own at the London Hospital for more than thirty years had taken the same quantity daily after dinner.

It is, I believe, universally acknowledged that the long continued habit of taking snuff irritates the fauces and epiglottis, producing cough, &c. Nor is dyspepsia the only further ill effect of this habit; the irritating particles extend through the whole length of the alimentary canal. Several inveterate snuff-takers have intimated to me the irritable state of the bowels; in whom it appeared that the mucous membrane was unnaturally stimulated and irritable. The oft-repeated stimulus leads to an enfeebled condition of the mucous membrane, to a loss of contractile power, as well as of healthy secretion and of nervous stimulus; as regards the stomach, dyspepsia is the result; in the intestine, diarrhoea or constipation; in some cases the rectum is principally affected, and it either retains the fæces so as to form an impacted mass, which it is unable to propel; or, if the excreta be fluid, the same weakness allows the contents to pass rapidly to the sphincter, which is itself so enfeebled as to be unable to restrain an involuntary discharge. Snuff may actually be seen among these excreta.

Drinking excessively of cold water induces an enfeebled, relaxed condition of the mucous membrane of the alimentary canal.

Cicatrices of the mucous membrane after ulceration, as in dysentery, lead to contraction and diminution of the canal, and act mechanically by obstructing the canal, and they interfere with regular peristaltic action. Tumours, or any growths pressing upon either small or large intestine, may induce constipation in a similar manner; but we defer entering into the causes of insuperable constipation, arising from cicatrices, till we speak of ileus. With these cases also we shall consider other more serious causes of constipation, namely, cancerous and fibroid growths, tumours connected with the intestine, or pressing upon it, and the various forms of internal strangulation and intussusception, &c.

In speaking of constipation arising from diminished secretion, we have alluded to cerebral disease, and to determination of blood to the head, from over anxiety and mental work. Various causes often co-operate in these instances; a sedentary life and diminished muscular exertion, is associated with changes in the secretions and with diminution of contractile power of the intestine; and, although the muscular layer of the intestine acts independently of nervous influence, there is strong reason to believe that it is modified by it, though in a less degree than the secretions of the mucous membrane. In many diseases of the brain, the abdomen becomes collapsed, as if the healthy tone of the parts was lost.

In diseases or injuries of the spinal cord, this relation with the alimentary canal is still more evident; the bowels are constipated, and action is often induced with difficulty; not only from paralysis of the parietal muscles, but from diminution of the contractile power of the intestinal muscular layer, as well as from change in the secretions of the mucous membrane. The paralysis is painfully shown in these cases by the want of control over the sphincter muscle; for the motions escape involuntarily.

In advanced life the feeble contraction of the parietes, the diminished excitability of the intestinal muscular coat, and the necessarily less active life, often produce constipation, which is increased by the nervous alarm of the patient.

Constipation is also a sign of inflammation of the peritoneal investment of the intestines; the muscular coat becomes involved, and ceases to contract with energy. This is a wise and beautiful provision, to which we have already referred.

Constipation also is induced when defecation is painful, as in inflamed hæmorrhoids, in ulceration of the rectum, and in diseases of adjoining parts. So severe is the pain in some cases, that action of the bowels is prevented by the sufferer, who is unwilling to undergo, or rather is desirous to postpone to the latest period, that which produces such intense suffering. It is a merciful provision that in health such necessary actions are free from pain.

It sometimes happens that a spasmodic constriction of the alimentary canal, especially the rectum, induces constipation;

in most cases, however, it will be found that there is associated with this spasmodic contraction some direct cause of irritation at the part, as minute fissure or ulceration of the mucous membrane, disease of the bladder and uterus, &c.

A *fifth* cause of constipation may be the state of the *contents* of the large intestine.

The *faeces* having become hard and impacted, remain like a foreign body, and are only removed with considerable difficulty. The character of the food may have been such as to induce this impaction; for many cases are recorded of substances which have been taken habitually, as brown coarse bread, leaving the undigested parts to become agglutinated; so also with calcined magnesia, taken medicinally day after day; and amongst lunatics stones and pebbles which have been swallowed may thus become impacted.

It is in the lower part of the large intestine that *faeces* generally become thus hardened; although it sometimes takes place to a less degree in the *cæcum*, and in the ascending and transverse colon.

Sixthly.—Mechanical obstructions have been cursorily alluded to in reference to tumours, as affecting the coats of the intestine; and it is of very common occurrence in pregnancy and ovarian growths, to find that direct pressure is exerted upon one or other part of the colon, so as to interfere with the regular and free action of the bowels.

Symptoms.—Constipation manifests its effects on the brain by inducing torpor of the mind with diminished energy and activity; the sleep is disturbed, and not refreshing, the mind easily agitated, and often melancholic. There is also a general *malaise*, which renders the patient unwilling to undergo ordinary exertion and fatigue; pain in the head, sometimes at the forehead, at other times in the occipital region, is often present; and when diseased arteries of the brain, or other predisposing causes of disturbed cerebral circulation exist, there is not unfrequently vertigo, with disturbed vision, haziness, sparks before the eyes, *muscæ volitantes*, ringing noise in the ears (*tinnitus aurium*); and occasionally there is momentary loss of consciousness.

When disease of the heart exists marked symptoms of dis-

turbance in the circulatory organs are sometimes produced; the most frequent, perhaps, is irregularity of the pulse, and uncomfortable palpitation of the heart; and the pulse is generally compressible. As to the respiratory organs, dyspnœa is not unfrequently induced by the impediment to free action of the diaphragm. Pain, especially across the sternum, is often ascribed to the chest, whilst it really arises from distended colon.

The abdomen is full, and sometimes masses of a round and hard character can be felt in the course of the colon, simulating morbid growths, which, when perceived, cause alarm to the patient; this state of partial impaction may exist, although there is daily action from the bowels, a central channel being left; the tongue is flaccid and indented by the teeth, showing an atonic state of the muscular fibre.

Various neuralgic pains are often induced, from direct pressure upon the nerves, sometimes in the right hypochondriac regions; frequently over the crest of the ilium in the course of the ileo-hypogastric nerve, and in the course of the genito-crural, to the groin and the testicle.

Aching pain in the loins and in the lower extremities arises from interference with the free return of blood; and beside this symptom, a varicose condition of the veins is induced or aggravated; and consequent œdema is produced. A similar condition of the hæmorrhoidal veins is also the result of habitual constipation; and all the discomfort attendant on hæmorrhoids follows. Irritation of the adjoining pelvic organs is sometimes excited, as irritability of the bladder.

It has been stated, that distended transverse colon may exert pressure on the duodenum, so as to lead to symptoms of dyspepsia, such an effect is exceedingly doubtful, but when adhesions have taken place between the first portion of the duodenum and the colon, great distension of the latter then exerts pressure; generally, however, these symptoms are due to the imperfect separation of excreta, and to congestion of the portal system.

Diagnosis.—The diagnosis of constipation may be considered as generally sufficiently clear, but the various secondary symptoms may lead to serious misapprehensions. As to impacted fæces

in the course of the colon, they have very often been mistaken for tumours; but their local character, mobility, and general symptoms serve to distinguish them. This discrimination is more easy in the ascending or transverse colon; but in the descending colon, and especially in the sigmoid flexure, the diagnosis is more difficult. Cancerous obstruction at the sigmoid flexure is very insidious, and gradual constipation is its principal symptom; but local pain, and a small, firm, hard tumour at that part are very diagnostic of an obstruction of this kind. Impacted fæces, however, in the rectum and sigmoid flexure, sometimes become so firm and immovable, that the symptoms may closely resemble organic disease; weeks may be passed without evacuation, and gradually severe symptoms result, as vomiting, and occasionally extreme pain. A careful examination will, in most cases, render the diagnosis easy, and the patient perseverance in injections and mild aperient remedies will be effective; nor do we find in simple impaction of fæces that the stomach becomes so irritable as in organic strangulation.

A case is recorded by Mr. Staniland,* of a patient, æt. 73, who had habitual constipation, so that, during the last five years of her life the bowels were only acted upon once in every two months; after being confined for four months and eight days they were very freely acted upon; seven months then elapsed without any pain or evacuation. Some weeks before death she had a fall, which produced very severe pain in the region of the cæcum, and led to local inflammation, gangrene, and fæcal extravasation into the peritoneum. The intestines were found enormously distended with fæces, the transverse colon was nine inches in diameter, and the sigmoid flexure ten and a-half; the rectum six inches. A remarkable instance of constipation of nearly four months' duration, after fever, is recorded by Mr. Gay, in the 'Pathological Transactions' of 1854. The patient, æt. 6, recovered.

The *treatment* of constipation is a subject of great interest, because it is one which so frequently tests the skill of the practitioner. A knowledge of the habits and diet is essential

* 'Medical Gazette,' p. 246. 1832-33.

to us in devising means of cure ; thus regular exercise, when the life has been sedentary, and especially walking or horse exercise, is of paramount importance. It is true that the beneficial effect of pure air may be otherwise obtained, but not all its good effects ; for carriage exercise is not alone sufficient. To be brought to town in many of the crowded conveyances which hurry to the city day by day, to be wearied by standing, or quietly sitting at the desk, and when exhausted conveyed home in a close omnibus, or railway carriage, is sufficient to induce discomforts of a hundred kinds in London, without the additional causes of the anxieties of life ; and in a less degree the same thing is everywhere observed.

An actual distaste or aversion to walking may be easily acquired, and in circumstances where wealth, position, and the value of time render it unneedful and undesirable, the beneficial effect of walking exercise is easily forgotten.

It is very desirable to promote constant regularity in the action of the bowels ; with many persons, an early movement before or after breakfast, removes discomfort for the rest of the day ; with others, though less desirable, the time immediately before going to bed is chosen.

The character of the food is an important consideration ; sometimes injury is done by taking more than the frame requires, and the stomach can digest ; or by a too great sameness in the diet, for variety is required ; not that at each meal numerous forms of food should be taken and satiety induced by the niceties of the culinary art ; but, an admixture of animal and vegetable food is necessary, and a change in them is requisite.

Vegetable food contains a large quantity of indigestible material and of alkaline salts which stimulate the alimentary canal, so that when there is a tendency to constipation, this means alone may be sufficient to remove it ; thus, brown bread acts by the irritating character of the indigestible parts of the grain. Again, gentle palpation of the abdomen, kneading the parietes with the palms of the hands, has sometimes induced action. The bracing tonic effect of a shower-bath, or in less degree of cold sponging, when it is not contra-indicated, may obviate constipation. These means produce their effect by the increased action

of the abdominal muscles ; but, another agent acts in a similar manner, namely, *electricity*. A galvanic current transmitted through the abdominal walls induces a very speedy action, or rather emptying of the colon ; it has been sometimes recommended in the constipation of painter's colic. I have used it with manifest advantage in paralysis. A case of partial paraplegia, in which injections did not act satisfactorily, and drastic purgatives were undesirable, had a galvanic current passed through the abdomen every morning ; in a few hours a free evacuation was produced without any discomfort. This agent, which has been employed to excite contraction of the uterus, may be frequently used with benefit in constipation.

Medicines, directly purgative, may be divided into several classes :—those which are—

1. *Laxatives*—Manna, figs, prunes, raisins, fruits, brown bread, cold water.

2. *Aperients*—Castor oil, almond oil, cod-liver oil.

3. *Saline purgatives*, as magnesia, sulphate of soda, and of potash, saline waters, bitartrate of potash, &c.

4. *Mild purgatives*—Senna, rhubarb, aloes, mercurial medicines.

5. *Drastic purgatives*—Jalap, colocynth, gamboge, scammony, turpentine, croton oil, elaterium.

Inspissated bile has been used as an aperient, from the idea that the excrementitious portion of bile is naturally purgative in its action ; but although ten or fifteen grains may act as an aperient, and assist in unloading the intestine, it is an offensive and less satisfactory remedy than others which we possess. These remedies act on different portions of the intestine and in various ways ; thus mercurial purgatives stimulate all the secretions, both those of the liver and of the mucous membrane ; senna, and saline purgatives act on the small intestine, and render the evacuations more fluid ; aloes, and the drastic purgatives act on the colon ; rhubarb has an astringent effect, and sometimes irritates and offends the stomach. Some stimulate the intestine to increased peristaltic action and excite griping pain. The action of the salines is partly due to exosmotic current from the capillaries of the

intestine, thus leading to the effusion of fluid in a greater extent into the canal than is absorbed from it.

The rapidity of the action of aperients is also very diverse. The salines act quickly, especially if given with a considerable quantity of diluent fluid. Aloes is slow in its action, and requires several hours to produce any effect. Drastic purgatives are often followed by much trying irritation in the rectum, and by tenesmus.

Strychnia, or nux vomica, is a valuable remedy in constipation; it excites the muscular coat to contraction, at the same time that a tonic effect is produced on the mucous membrane. It is well to combine with it purgatives and sedatives, as aloes, and henbane, &c. Preparations of steel often act as purgatives in the same manner.

Podophyllin, from the *Podophyllum peltatum*, the May apple or mandrake, has been long used in the United States; in small doses grs. ss—j, it acts as a mild purgative, producing an evacuation from the bowels slowly, but efficiently and without pain. I have observed it act thus favorably in chronic ulcer of the stomach; in large doses it produces violent vomiting and purging; thus in an instance in which a nurse, contrary to directions, gave seven to ten grains, severe colic, vomiting and dysenteric diarrhoea followed, but subsided in a few days.

The use of glysters is too frequently neglected in ordinary constipation; but, their beneficial effect is now more generally acknowledged; some act simply by irritating and distending the intestine, thus exciting it to contract; as warm water and gruel; purgative substances may be added, as soap, castor-oil, colocynth, turpentine, and rue; the last two are especially used when constipation is associated with flatulent distension of the intestines.

The excessive use of glysters even of the mildest kind, as water and gruel, and especially when they are administered in large quantities, induces distension of the rectum, and an un-readiness to act without the wonted stimulus.

I cannot leave the subject of the use of purgatives in ordinary constipation, without speaking of the injurious effect of their indiscriminate and injudicious use; to some the use of a dinner pill or an aperient at night, is constant, year after year; in

others a slight discomfort leads to the use of the blue pill or black draught, or to still more active agents. Temporary relief is afforded by powerful purgatives, but the delicate mucous membrane of the intestinal tract is weakened, a state of chronic catarrh is induced, and the very condition sought to be removed is aggravated tenfold. In an enfeebled person, violent purgative medicine has in very many cases induced excessive prostration, and even fatal results; and, in them, it is easy to excite a state of irritation which it is almost impossible to subdue.

The administration of vegetable tonics, with mild purgative medicines, and with ammonia, is often of great utility; a valuable preparation of this kind is the compound gentian mixture, which contains senna, gentian, orange and lemon peel, ginger, and tincture of cardamoms. The combination of aloes and myrrh is a preparation of a somewhat similar kind, the tonic effect of the myrrh is associated with the purgative of the aloes.

Purgative medicines sometimes act more beneficially in combination; as slight mercurials, when the secretions of the liver are imperfect, with aloes, rhubarb, and colocynth.

The addition of an anodyne or carminative, as hyoscyamus, Dover's powder, the essential oils, &c., with more active remedies, is beneficial in removing their irritating character, and in preventing the griping pain sometimes induced by them when given alone; thus the compound gamboge pill, and compound colocynth pill with henbane, act as efficient but tolerably mild purgatives, emptying the large intestine; or the purgative may be sheathed by mucilaginous and oleaginous substances, as rhubarb with linseed oil.

In infants, constipation is sometimes an exceedingly troublesome affection; the repetition of castor oil is trying, and even injurious; an old-fashioned remedy is that of exciting the intestine to contract by introducing a very small glyster pipe into the rectum, or a portion of soap cut into a conical shape; magnesia may be given in a tasteless form, the calcined, or citrate, &c., or sometimes a small quantity of gruel will excite the bowel to slight action; in any case, however, irritating medicine must be avoided. It is difficult to over-estimate the injurious effect in

children of repeated doses of calomel, of jalap, &c. ; muco-enteritis is induced, and sometimes fatal results follow ; scammony with milk is a convenient remedy in some cases, but must be used with caution.

In the *aged*, enfeebled either by a life of activity or by declining strength, the intestines lose their normal power of wonted contraction ; to use drastic purgatives is out of the question, and a constant change of milder aperients is necessary. The mildest laxatives may suffice, as a draught of cold water, prunes, figs, roasted apples, brown bread, manna, the confection of senna, or the compound rhubarb pill, alone or with henbane, so the compound colocynth pill, and scammony pill with henbane and Dover's powder, or a few grains of dried rhubarb with capsicum and soap, may be given with each principal meal ; and to these, in some instances, very minute doses of strychnia are added, with considerable benefit.

When the muscular coat of the rectum loses its contractile energy the contents sometimes become so impacted and hardened as subsequently to withstand the most powerful efforts at expulsion ; purgatives, and even copious injections, are insufficient to soften the hard contents, and mechanical assistance has in not a few cases been required. Hard masses may be retained in the colon and rectum even although there be periodical action, even of a fluid kind ; for, laterally, fluid may pass, especially after purgative remedies, whilst scybala are still retained. In advanced life, in spinal disease, in constipation after powerful purgatives, this state is occasionally present, but it has in rare cases been witnessed in very early life. Warm copious injections, whilst enteric irritation is avoided, will suffice to relieve most of these impactions.

Concretions may form in the intestines from deposit upon extraneous substances, from impacted biliary calculi, but more frequently they consist of hardened fæces, or of the undigested portions of food and medicine, as from oatmeal, brown bread, &c. Dr. Harley, in an interesting communication on this subject to the Pathological Society, in 1859, records an instance of concretion consisting of starch, taken to relieve dysentery ; another of benzoin, which had been taken to improve the voice, and had formed a concretion as large as a bean. Again, large

quantities of magnesia have been found as a mass in the colon, and portions of string become impacted in like manner.

Foreign bodies of considerable size sometimes pass through the whole intestinal canal without producing any injurious symptoms, as coins accidentally swallowed, stones taken by maniacs; in other cases they are retained at the sphincter and require mechanical assistance in their removal, as fish-bones, &c., placed across the intestine.

In a remarkable instance in which a sailor swallowed clasp-knives, several were discharged from the bowels, and one was found fixed transversely in the rectum; the case is recorded by Dr. Marcet, in the 'Medico-Chirurgical Transactions,' and the thickened stomach, with the fragments of the blades found on examination after death, are preserved in the Museum of Guy's Hospital. The patient was an American sailor, aged 23, who, in June, 1799, swallowed four clasp-knives; three were discharged from the bowels. In March, 1805, he swallowed fourteen knives in two days; in December, 1805, he swallowed fifteen to twenty more; making thirty-five swallowed at different times. His health became impaired; he vomited the handle of one, and passed portions of the blades of others; and in March, 1809, he died in a state of extreme exhaustion. The œsophagus and stomach were dilated and thickened, and in the latter were numerous blades of knives partially dissolved. In the abdomen there was a general discoloration of the intestines; one blade was found perforating the colon opposite the kidney, but without extravasation of fæces; another blade was transversely fixed in the rectum.

CHAPTER XIV.

INTERNAL STRANGULATION.—INTUSSUSCEPTION.— CARCINOMA OF INTESTINE.

VARIED conditions, leading to insuperable constipation, have frequently been indiscriminately associated together, under the term ileus ; and whilst we are willing to acknowledge, that very great difficulty is connected with the correct diagnosis of these cases, we believe that when a full history of the symptoms can be obtained, careful examination will enable us to divide them into several classes, and to make an approximative diagnosis, not only as to the character, but the position of the obstruction. Each minute circumstance is important in assisting the correct diagnosis of these cases, the accurate detail of previous symptoms, the mode of attack, the position of pain, the vomiting, the relative severity and period of commencement of these symptoms, the state of the abdomen, the general appearance of the patient, the quantity of the urine, &c. Dr. Barlow has drawn attention to several of these conditions, and has shown the importance of ascertaining the period of the commencement of the vomiting,* and the condition of the renal secretion.

The causes of insuperable constipation are numerous :

1. Internal strangulation arises from bands of adhesion, which may be the result of inflammatory action ; or from simple adhesion without constricting bands, as of the ileum to the uterus.
2. From congenital intestinal pouch becoming adherent to other parts, or from perforations in the mesentery, &c.
3. From the appendix cæci assuming a fixed and adherent position.

* 'Guy's Reports,' 1844. Clinical cases. Practice of Medicine.

4. From the twisting of the intestine, of which Rokitansky gives three forms—I, Upon its own axis; II, Upon the mesentery; and III, Upon other coils of intestine. Other causes are—

5. The development of tumours in the mesentery leading to constriction.

6. Intussusception.

7. Cancerous disease of the intestine.

8. Contraction of cicatrices, as after dysentery; fever.

Beside these, we must also enumerate:—

9. Enteritis and peritonitis.

10. Impaction of *fæces* and of foreign bodies, as gall-stones, &c.

11. Obscure forms of hernia, as into the obturator foramen, &c.

12. Prolapsus ani, and inflamed hæmorrhoids.

13. Abdominal and pelvic tumours.

Abercrombie* describes cases of ileus in which no cause of strangulation nor obstruction was detected after death, and he believed them to arise from distension, or “simple derangement of action” of the intestine; thus he states, “that distension appears to constitute a morbid condition which may be fatal without passing into any farther state of disease;” and again, that “ileus does not appear to be necessarily connected with obstruction in any part of the canal; for we have seen it fatal without obstruction, and we have seen everything like obstruction entirely removed without relieving the symptoms.” He mentions other instances in which adhesions had formed without sensible diminution of the area of the intestine, and which were followed by the symptoms of insuperable obstruction; in the former we believe that either enteritis was present, or the bowel was twisted; in the latter that spasmodic contraction rendered a partial impediment complete; and the author just mentioned writes, “I admit, however, that there may be irregular contractions of portions of the intestine, analogous to that to which the term spasm is usually applied, and that these may form the first step in that chain of derange-

* Abercrombie on ‘Diseases of the Stomach and Intestine.’ Third edition.

ments of the harmonious action of the canal which leads to an attack of ileus."

Internal strangulation.—The general symptoms of this condition are pain sometimes very slight, but occasionally very severe, gradually increasing distension of the abdomen, constipation, generally of an insuperable character, vomiting at first bilious, afterwards stercoraceous; and after a longer or shorter period peritonitis, prostration, and death.

Pain.—In many cases of internal strangulation there is a sudden catch in the bowels, as of some displacement, and the patient can place the hand on the exact part, which generally indicates the seat of disease; although, if fatal, we may find that distension and other causes have led to considerable alteration of position. When a portion of intestine has slipped under a band of adhesion, or into a hole of omentum or mesentery, this character of pain is observed, but when there has been a twist of the intestine the pain is more gradually developed, and for many days may be entirely absent. The most obscure cases are those of internal strangulation, in which there has been chronic partial constriction, and from indiscretions in diet, or other causes, slight enteric irritation has led to spasmodic constriction at the part; in these cases the pain closely resembles ordinary colic. Tenderness of the abdomen may be absent for many days; in some instances the peritonitis does not come on till nearly the close of life, from a state of continued and extreme distension of the intestine, and from ulceration of the mucous membrane extending to the serous coat; but where there has been sudden strangulation, the serous membrane is more quickly implicated, and the symptoms bear a closer resemblance to those of ordinary external strangulated hernia. If the strangulation be in the small intestine, either near the cæcum, or in the jejunum, the pain will generally be found to be in the region of the umbilicus; where the colon is involved the position of the pain is in the course of that part of the intestine, and often marks its precise seat; thus, in diseases of the sigmoid flexure, the pain will generally be found in the left iliac fossa or in the left groin. Its character in intussusception is often severe, paroxysmal, and resembles that of ordinary colic; in many instances a tumour

can be felt in the abdomen, arising from the involution of the intestine, or adhesion between its coils.

Tympanitis.—Unless the obstruction be very high in the alimentary canal, as in the case recorded with disease of the duodenum and of obstruction twenty inches from the pylorus, the abdomen gradually becomes distended, and tympanitic on percussion. The enlarged coils of intestine may be observed through the stretched parietes, and the peristaltic movements are often clearly perceptible, especially in obstruction of the colon. If the ileum, or the commencement of the ascending colon, be constricted, the distension is central in its character, and is less evident; but if the descending colon, sigmoid flexure, or rectum, the portions of the large intestine above the seat of disease become greatly distended; they may be observed in the peculiar outline of the abdomen, and the tympanitic resonance extends to the *loins*; where, however, the obstruction arises from portions of twisted large intestine, as of the cæcum or sigmoid flexure, we find that there is some deviation from the general character just mentioned; an enormously distended cæcum may be twisted over to the left hypochondrium, and constitute a prominence in that region.

Vomiting.—The character of the vomiting, and the period at which it has commenced, especially when irritating and powerful purgative medicines have not been administered, are important guides to our diagnosis. If the obstruction be sudden, and be situated in the small intestine, the vomiting comes on very quickly, in from half an hour to two or three hours; if it be high in the jejunum, the vomited matters are of a bilious character, but if near to the cæcum they may assume a fæcal odour, and be completely stercoraceous. In one instance, in which the obstruction arose from a band of adhesion high up in the jejunum, the vomiting was so sudden as to resemble that produced by cerebral disease; and this view of the case was favoured by the partial insensibility of the patient. In the case recorded of twisted cæcum; where the obstruction was near the termination of the ileum, so fully fæcal was the character of the vomited fluid that it was for a time supposed that a communication existed between the stomach and the transverse colon. When the large intestine is the seat of disease, as in cancer of

the sigmoid flexure, and of the rectum, &c., several days sometimes elapse before vomiting supervenes; the time is, however, much accelerated if powerful drastics are given, as in Case CII. In the latter state, also, the vomiting is more easily checked by the administration of remedies, as of ice and opium, &c. As to the immediate cause of stercoraceous vomiting, Dr. Brinton, in his valuable remarks in the 'Encyclopædia of Anatomy,' has clearly shown that the peristaltic action is not in itself reversed, but that the contents of the bowel are propelled onwards in their normal manner till the obstruction is reached, when the fluid assumes a central retrograde direction, thus producing a double current, a parietal or onward, and a central or reverse current; this retrograde movement continues till the vomited matters are of the same character as those found at the seat of stricture.

Hiccough is also more severe and more speedily produced in the strangulation of the small than of the large intestine. It must be borne in mind, that the vomiting and hiccough are sometimes extreme in peritonitis, where the serous membrane of the stomach is involved.

Urine.—Dr. Barlow has drawn especial attention to the amount of urine excreted, as a sign of the seat of obstruction; that where the obstruction is high in the canal, as in the jejunum or ileum, absorption is partially checked, the renal vein receives a diminished supply of blood, and a small quantity of urine is excreted; if, on the contrary, the rectum or sigmoid flexure be occluded, nearly the whole of the capillaries of the alimentary canal are free to absorb fluid, and thus the blood contains more watery constituents, and the urine is abundant. This is a symptom deserving our attention, but it is not a certain one; several cases among those illustrative of disease of the sigmoid flexure had scanty urine among their earlier signs, and we shall find that the amount of urine may be measured by the quantity of fluid vomited; that if in obstructed colon powerful drastics have been administered, and speedy vomiting induced, or peritonitis quickly set up, the urine will be found to be small in quantity. However, it is as true that the urine is abundant when the obstruction is low down in the canal, as that the vomiting is late in its occurrence. The fluid character of the

contents generally observed in the distended intestine above the seat of stricture is to be remarked, and is an indication that no remedies are needed in these cases to render them more watery, but that the spasmodic state of the diseased bowel, in addition to the mechanical impediment, often prevents a drop of fluid or any gas from passing the stricture.

State of the rectum.—Dr. Barlow has here also added his diagnostic acumen to the elucidation of the symptoms presented. He has shown that in obstruction suddenly produced, the rectum retains its natural power of contraction, and will be found to be empty; if the disease be of gradual formation, that it is more patulous and readily yields to injections. To a certain extent this is the case, but it is not a symptom upon which we can rely. The intestine below the obstruction is generally contracted, and sometimes after the occurrence of the strangulation or other occlusion, a fæcal evacuation may take place, or be removed after injection from below the seat of stricture, thus giving a delusive hope of recovery, or misleading in diagnosis. Mr. Charles H. Moore has proposed the injection of fluid into the colon, the extent of the dulness on percussion in the loins being carefully noticed, as a sign of the position of the obstruction; and that in this way fluid may be forced into the ascending colon, and indicate that the disease is above that part.

The discharge of blood, or of offensive mucus, has been mentioned by Mr. Gorham* as a very frequent sign of intussusception; and it may be here remarked that it is important in all cases of this kind to make a careful manual examination of the rectum, as well as of all the parts in which hernia may occur. By this simple means impacted fæces, inflamed hæmorrhoids, cancer of the rectum, prolapsus ani, suppuration in the pelvis, each of which may lead to symptoms of insuperable constipation, may be diagnosed.

Previous attacks of peritonitis, constipation, or glandular disease, afford some indication of the probable cause of the malady; inflammation is very likely to have left bands of adhesion, and attacks of constipation of less severe character are often found to have occurred in cancerous disease. Abercrombie

* 'Guy's Reports.'

mentions several instances of this kind, and the cases here recorded furnish illustrations of the same fact.

The progress of the symptoms, and the period at which a fatal result ensues, is subject to great variation. In sudden strangulation of the small intestine, we sometimes observe death following in five to seven days; whilst in other cases, especially of the colon, several weeks may elapse, and the patient remain free from pain and distress till about forty-eight hours before death. The obstruction and distension of the intestine lead to enteritis, and the mucous membrane becomes ulcerated. The inflammatory action extends to the peritoneum, so that it is very rare to find a case of fatal obstruction without peritonitis; sometimes merely a dry and congested state of the serous membrane, in others lymph is effused, and in many there is perforation. The perforation of the intestine is often observed at the seat of the constriction, and is most marked at its upper limit; but in diseased rectum and sigmoid flexure, it will be frequently found that perforation has taken place, not only at the constriction, but at the cæcum. The ulceration of the mucous membrane in these instances is also peculiar; it is somewhat similar to that presented by the skin which has been overstretched, and affected with erythematous inflammation and superficial ulceration. It is, in instances where the obstruction is primarily from the mucous membrane, as in cancerous growth, that peritonitis is most slowly developed. Where all the coats of the intestine are involved, as in many cases of internal strangulation, the vessels of the mesentery become also obstructed, œdema is produced, and in a short time gangrene follows.

The part of the small intestine most frequently strangulated either by loops, bands, or adhesions, is the lower portion of the ileum; in the colon the sigmoid flexure is that part which we find most commonly diseased, and very frequently the disease is of a cancerous character; sometimes, as in one instance under the care of Dr. Rees, in Guy's Hospital, the constriction is two-fold, a band constricted the upper part of the jejunum, and a second band the ileum.

Rotation of part of the intestine upon the mesentery is also more usually observed in the small intestine; in such cases a loop appears to have twisted, and by the traction of the mesen-

tery, and the pressure of superincumbent parts, it becomes strangulated.

In the colon the intestine sometimes turns over upon its own axis, as in the instance of the cæcum which we have related, and as in the cases of obstruction in the sigmoid flexure, recorded by Dr. Barlow in the Guy's Reports, in the year 1844. I have never witnessed any instance of the third class referred to by Rokitansky, in which one loop of intestine becomes twisted upon another as an axis.

Intussusception is that condition in which one portion of the intestine passes into another, as the finger of a glove drawn within itself. In this state it would not necessarily become entirely obstructed, were it not that the congestion, effusion, and inflammation produced close the canal completely. The section presents us with three layers of intestine; two mucous surfaces and two serous being opposed to each other, and in the centre are placed the ordinary mucous surfaces. There is sometimes a second involution of the intestine from below, passing in an opposite direction; or the only involution may be from below upwards. The mesentery attached to the involuted portion is also drawn in, and by its traction the central portion of intestine becomes somewhat curved laterally, and the opening of the most depending part is observed to be linear. The vessels of the portion of intestine thus incarcerated become engorged, and render the obstruction complete; the whole of the folds involved become swollen and deeply congested, and blood is extravasated into the substance of the mucous membrane, as well as into the mesentery; in a short time both the serous and the mucous surfaces become inflamed, and an effusion of lymph takes place; the opposed serous surfaces become adherent, and so to a less degree the mucous surfaces; bloody serum and mucus are effused into the canal, and this discharged per rectum is a diagnostic sign of intussusception. If life be prolonged, and the intussusception continue, the serous surfaces at the opening or upper part are rendered adherent; the contained intestine becomes gangrenous, and is often detached as a slough. In this way many inches of intestine may be discharged per rectum; in one instance as much as forty-four inches of large intestine were evacuated; in another, which terminated favorably, and the

specimen of which is in the museum at Guy's, the whole of the cæcum and ascending colon were thus passed. If the adhesions be disturbed or broken down after the slough has separated, faecal abscess may be the result. In some instances the intussusception is restored ; more frequently, more and more intestine is forced in, symptoms of internal strangulation supervene, and death results from perforation into the peritoneum, or from peritonitis set up by the direct extension of disease from the strangulated part. It must not, however, be supposed that the passage is always occluded ; such is sometimes not the case, and even diarrhoea may be produced. It would seem that the intestine may be thus incarcerated within another fold, without being strangulated. The case recorded by Dr. Hughes, in the Guy's Reports, was of this kind ; so also those of Mr. Phillips in the 'Medical Gazette ;' and still more remarkably a specimen exhibited by Mr. Hutchinson at the Pathological Society, in which the symptoms extended over seven months.

The usual position of intussusception is in the small intestine, but a portion of ileum not unfrequently passes into the cæcum, and lastly the intussusception may consist only of one portion of colon into another ; the cæcum and ascending colon may become so involuted as even to reach the rectum. It would appear that in intussusception into the colon, constipation is less constant as a sign of disease.

According to the following tables from Dr. Brinton's Croonian Lectures, recorded in the 'Lancet,' 1859, it would seem that ileo-cæcal intussusception is the most frequent variety ; this may be the case, if we exclude the numerous instances of intussusception which take place during the article of death :

Intestinal Obstructions (excluding Hernia).

Frequency, 1 in 280 deaths (from 12,000 promiscuous necropsies).

Varieties, relative frequency per cent. (from 600 necropsies of obstruction).

Intussusception	43
External (bands, &c.)	32
Parietal (strictures, &c.)	17
Torsions	8
	<hr/>
	100

Intussusception, varieties of, per cent.—

Ileo-cæcal	56
Ileac	28
Jejunal	4
Colic	12
						<hr/> 100

The number of intussuscepted portions also varies much, being sometimes single, but in young persons, and especially infants, it is exceedingly common to find numerous parts so diseased, six to twelve, or even more. Some of these, however, are probably produced immediately before death; there is absence of all symptom of strangulation, and in the intestine itself neither congestion, effusion, nor ulceration exists; they are most frequently observed in inflammatory disease of the brain, and hydrocephalus.

The *symptoms* of intussusception are those of colic with constipation; sudden local pain is produced in the bowels, followed by vomiting, constipation, prostration, haggard expression of countenance, failing pulse, distension of the abdomen, stercora-ceous vomiting, peritonitis, and death. It is exceedingly difficult to distinguish this condition from ileus arising from internal strangulation and local enteritis; but after a time there may be discharge of blood and mucus from the involuted portion, which materially assists in forming a correct diagnosis. And in intussusception, at the seat of pain a firm mass may often be felt at one or other part, which is not the case in ordinary internal strangulation. In intussusception of the small intestine the tumour may be central, whilst in ileo-colic and colic involution the mass will be in the course and position of the colon. The sudden onset of the pain, and its subsidence, becoming aggravated in paroxysms, is an indication of this form of obstruction. It has been before mentioned that diarrhoea sometimes supervenes, especially where the large intestine is affected; such is occasionally noticed where the disease is of a chronic character. In seeking to arrive at a correct diagnosis, it is well always to examine the rectum:

The *cause* of this abnormal involution appears to be sudden and spasmodic contraction of a portion of intestine, impelled onwards into a part which is less contracted or altogether flaccid.

It occurs at all periods of life, but is much more frequent in youth and infancy.

The *prognosis*, although very unfavorable when we have well-marked indications of the existence of intussusception, is not without hope, and we have seen almost hopeless cases recover. It is probable that in some the intestine is restored to its normal state; in others, the strangulated bowel sloughs off, the canal becomes free, and the divided intestine unites.

Cancerous Disease of the Intestine.—Cancer of the stomach is a disease of frequent occurrence; but cancer is more rare in other portions of the alimentary canal. We have already described instances of it as affecting the duodenum and the cæcum, and other portions of the small intestine and of the colon are occasionally thus diseased; still the rectum and the sigmoid flexure of the colon are the parts of the intestine most frequently affected by it, and it is to the latter that we direct especial attention as a cause of obstruction.

The termination of the sigmoid flexure appears to be particularly prone to this form of disease, and many of those which are described as at the first third of the rectum, are at this part, and have become pushed down into the pelvis by the obstruction produced. It is a peculiar form of disease that we find thus developed; not the extensive deposit with glandular infiltration, though this is sometimes the case, but it is a modification of scirrhus. There is a growth from the mucous membrane, the muscular fibre is infiltrated and contracted, and the calibre of the intestine is diminished. The glands are frequently not at all affected, and in this respect it closely resembles epithelial cancer. The constriction and growth from the intestine are sometimes circular, sometimes one side is much more affected than the other. On examining the condensed part we find fibrous tissue, and some elongated nuclei; but the growth from the mucous membrane presents more of the elements of cancer. These, however, are not like the ordinary epithelial cancer elements, but many of the cells resemble large columnar epithelium, with a large nucleus, as if the growth had a modified form on account of its being situated on a columnar epithelial surface, an instance of the differentiation of abnormal growth.

Medullary and colloid cancer sometimes affect this part, but

their course is often different from that of scirrhus disease; the ulcerative process extends through the coats of the intestine more rapidly, and instead of intestinal obstruction we have fæcal abscess, either in the iliac fossa, or within the abdominal cavity itself. In the rectum similar forms of disease are presented, and occur in its several parts: and scirrhus, leads to contraction, thickening of the external tissues, and obstruction of the intestinal canal. Medullary cancer, on the contrary, causes ulceration and communication with the other pelvic viscera, with the vagina, bladder, or uterus, so that all the viscera become matted together into one mass. The rectum also frequently becomes involved, by the extension of disease from the uterus and vagina, leading to terrible manifestations of disease and suffering. Epithelial growths of a cancerous character arise from the mucous membrane of the rectum, as well as of the sigmoid flexure, and they lead slowly to obstruction or to exhausting diarrhœa.

Where the coats of the intestine are thus diseased, the intestine above the stricture becomes gradually distended, the mucous coat thickened, the muscular hypertrophied, so as to be in some cases a quarter of an inch in thickness. The extent of these changes varies much, and in chronic and slowly progressive disease they are more manifest. The distension of the intestine also produces inflammation and ulceration of the mucous membrane above the stricture, and it leads in many cases to perforation; this condition of ulcerative erosion is sometimes very extensive, at a considerable distance from the seat of obstruction; thus we find perforation of the cæcum taking place in obstruction of the sigmoid flexure.

Cancerous diseases of the ascending or transverse colon take place more rarely, and appear to be produced by some local exciting cause, as the cicatrix of an ulcer, or by a blow; still they are far from infrequent. In a case of colloid cancer of the stomach we observed a similar state of the ascending colon, but in a less advanced condition.

Mr. Birkett, in the 'Pathological Transactions,' has recorded a remarkable case of vascular villous growth from the colon near the liver; the growth was covered with epithelium, and its cancerous character was very doubtful. It was taken from

a man, aged fifty-eight, who a year before his death had had pain in the abdomen and diarrhoea. Two months before admission into Guy's, he had had constipation and pain, and when brought to the hospital he had symptoms resembling strangulated hernia, with constipation of one week's duration; he had had scrotal hernia, and the sac remained. The cæcum could be seen distended, and so also the ascending colon, as far as the liver, where there was pain on pressure; the descending colon could not be felt. Mr. Birkett explored the hernial tumour, but no intestine was within it. The propriety of opening the ascending colon was discussed; but the patient died four days after admission. (See Prep. in Guy's Museum, 1854⁶⁵.) In another instance, Case CCX, a blow on the hypochondrium was followed by a cancerous growth, which led to fæcal abscess and to perforation of the jejunum.

Carcinoma of the stomach sometimes extends to the transverse colon; such was the case in one of the instances we have recorded of disease of the stomach; but although there was fæcal eructation, no stercoraceous vomiting occurred. Drs. Gairdner and Murchison have shown the important diagnostic indications of this symptom in communication between the stomach and intestine.*

The ileum and jejunum are very rarely affected with primary scirrhus and medullary cancer; they are sometimes involved in cancer of the mesenteric glands; but we have never observed insuperable obstruction thus produced. In Case CCXIII the symptoms of disease were very obscure.

Symptoms.—In scirrhus disease of the sigmoid flexure, if we possess a history of all the symptoms, the nature of the malady may often be correctly shown. There is slight pain, fixed in character, and remaining for a variable period, in the left iliac fossa, with constipation, or an irregular condition of the bowels: after one or two attacks of this kind, with several months or years between them, the constriction becomes narrowed to such an extent that a very slight increase renders it complete. The bowels again are confined, the patient feels uncomfortable from their loaded condition; the abdomen is gradually distended, vomiting comes on, and the symptoms of

* 'Edinburgh Monthly Journal.'

insuperable obstruction follow. The vomiting occurs much later than in obstruction of the small intestine, unless powerful drastic purgatives have been administered; the secretion of the urine continues free, and the patient may appear in comfortable health, except that the bowels have not been acted upon. After ten or twelve days, however, if no evacuation ensue, the colon becomes much enlarged, its distended coils can be seen through the parietes, and there is tympanitis in both lumbar regions; the urgent peristaltic movements may be detected through the parietes; at last ulceration takes place above the seat of stricture, and leads to fatal peritonitis and extravasation; or peritonitis arises from the enormous distension, and the more general inflammation of the coats of the intestine. Sometimes, after symptoms of threatening peritonitis, with judicious treatment the bowels are acted upon, and the patient is for a short time spared; even diarrhœa will occasionally supervene; the patient then continues much enfeebled, and after a few months sinks exhausted, or another attack of constipation terminates fatally.

The same symptoms of insuperable obstruction sometimes arise in medullary and colloid cancer; but, as before stated, they less frequently terminate in complete occlusion. The intestines become united together, ulceration extends through the coats, local peritonitis and fecal abscess are the result, with severe pain and hectic fever; or the cancerous ulceration may extend into the iliac fossa, and suppuration may burrow down beneath Poupart's ligament, as in disease of the cæcum.

There is much less pain in cancerous disease of the sigmoid flexure than of the rectum, because the parts are more free, there is less pressure on the nerves, and the adjoining structures are less involved. If the *rectum* be affected, the constipation and difficulty of defecation is more constant; the pain produced is often intense, especially where the lower third is affected. The feces become flattened; this may be the case when the sigmoid flexure is the part diseased; but is less liable to occur, for the feces can be retained for a sufficient period in the rectum to re-assume their ordinary character. In the later stages of cancerous disease of the rectum, and in the fibro-cellular thickening and contraction of its coats, diarrhœa instead of constipation may

occur ; I have seen several such instances, and Nelaton refers to this liquid condition of fæces and their free separation as not unfrequent in syphilitic disease. Tactile examination may detect disease at the lower part of the rectum, but not at its upper third. The extension of disease to the bladder, vagina, and uterus, leads to most distressing complications, and special symptoms are produced. In obstruction of the alimentary canal, as we have before said, the rectum should always be examined.

The symptoms of cancerous disease of the ascending or transverse colon are of the same kind as we have just described, but it is more easy to detect a hardness or tumour produced by the growth in the intestine. In many instances there is severe pain at the seat of the obstruction at an early stage, and this pain I have in several instances noticed as being produced as soon as fluids were taken ; diarrhoea is sometimes present, or it alternates with constipation, Case CLVIII. Again, we do not find that the transverse colon becomes distended and tympanitic across the abdomen ; nor that there is the same resonance in the left lumbar region. These indications, however, must be used with great caution, because the distension of the small intestine may lead to enormous coils, and be mistaken for an enlarged colon. The suggestion of Mr. Moore may be tried, namely, the injection of water into the colon and the examination of the amount of dulness produced. The intestine, however, in some cases, becomes so contracted below the seat of stricture, as not readily to yield to the injection of water, and we might be led to very incorrect diagnosis by this means, if we trusted to it alone.

In many patients who are affected with cancerous obstruction of the sigmoid flexure, there is but little emaciation or appearance of cancerous cachexia. They may be well nourished, and apparently in health ; generally, however, there has been some indication of disease, as shown by troublesome constipation, an occasional fixed pain, and sometimes by a discharge of mucus from the rectum.

These forms of scirrhus cancerous disease rarely occur in early life ; at that period it is more likely to be medullary or colloid in character ; but there are exceptional instances in this respect.

The position at which the sigmoid flexure becomes affected is at the brim of the pelvis, where it is more liable to temporary compression. It is also at that part where the triple longitudinal muscular band assumes a continuous character around the intestine.

Diagnosis.—In our remarks on internal strangulation, we have pointed out the diagnostic value of many of the symptoms presented. The varied causes of obstruction must be also borne in mind; comprising not only the forms of internal strangulation, of intussusception, and of cancerous disease, but the presence of tumours, of enteritis or peritonitis, the impaction of fæces or of foreign bodies, tumours connected with the uterus or ovaries, hæmorrhoidal tumours. In *intussusception* there is generally more pain resembling colic, there is the discharge of bloody mucus, and a tumour can frequently be felt at the affected part; and in not a few cases the involved bowel may be detected by examination per rectum. In *internal strangulation* the vomiting is more severe, the onset more sudden than in cancerous disease, and frequently something has been felt to have given way or slipped; there is greater resemblance to the symptoms of ordinary hernia; the small intestine is the part that is generally thus strangulated; and then, whilst the vomiting is more early and severe, the abdomen is less distended, and the course of the colon cannot be so easily traced. In *impacted fæces* alone, unless some foreign body be also present, the symptoms rarely, if ever, become so urgent, and scarcely ever fatal. In speaking of constipation, we have quoted a case from the 'Medical Gazette,' where after seven months of fæcal obstruction, the patient had a fall, and peritonitis was produced; she had had attacks of constipation of two months' duration, for four years. In simple fæcal retention, after a whole month has elapsed, we may find very little discomfort, and the distension not extreme in degree. In *cancerous disease* of the sigmoid flexure, the gradual character of the obstruction, the seat of pain, the distension of the abdomen without tenderness, the abundance of urine, the late period at which vomiting occurs, are the principal signs; and many of these cases closely resemble simple fæcal impaction. In the one we shall probably find that injections per rectum will be effective, and after a time they will be followed by relief; in the other,

the injection will in a short time be returned, only a small quantity can be admitted, and no faecal contents are evacuated.

The early tenderness of the abdomen distinguishes *enteritis*, and *peritonitis* in most cases; and recto-vaginal examination serves to remove other sources of diagnostic difficulty.

Treatment.—If after the administration of mild aperient medicines, or even without their use, it has been ascertained with tolerable certainty, that constipation from one or other of the causes we have described exists, it is exceedingly unwise to employ active treatment. Purgatives of all kinds are better avoided, and the use of drastic measures will tend to aggravate the sufferings, to shorten life, and remove the possible chances of recovery.

The administration of *opium* is now known to be attended with beneficial results, and frequently with partial, if not with permanent relief. By this means the peristaltic action is checked, spasmodic contraction diminished, and the opening which previously would not allow the passage of flatus, will suffer fluid faeces to escape. Solid opium may be given, as in the soap and opium pill. Some combine calomel with the opium, but we prefer opium alone, for the mercurial medicine increases depression; it probably hastens perforation, and extravasation is less likely to be limited by adhesion after its action.

Drastic purgatives, as colocynth, croton oil, scammony, mercurials, &c., stimulate and excite the intestine to greater contraction; vomiting of a stercoraceous character is set up more quickly or is increased, and ulceration or fatal peritonitis is speedily produced. Crude mercury is equally objectionable; and electricity, which is a valuable remedy in simple constipation, is here productive of injury to the patient.

If there be indication of local peritonitis, leeches may be applied freely, and rest in the recumbent posture enjoined.

The diet should be spare, and of a fluid, unirritating, and non-stimulating kind. If, however, we find great prostration, it is well to give brandy or wine, when they can be taken.

Injections are of great value in removing faecal concretions from the rectum, and below the seat of the stricture; and are sometimes followed by the discharge of flatus, affording great

relief to the patient. In this way warm water, soap, castor oil, or turpentine injections may be used ; and several ounces of simple olive or linseed oil thus thrown into the rectum sometimes afford considerable relief. Injections of this kind are most effectually administered by means of Dr. O'Beirne's long tube ; care, however, must be used lest the extremity of the tube turn upon itself. The simple introduction of an enema tube, and its retention for a short time, may excite the lower bowel to contract, and cause the expulsion of flatus, thus relieving the painful distension. In some instances, it is well to use nutrient enemata, which may serve to prolong the exhausted powers of life for a short period.

By the use of these means, when the patient is almost *in extremis*, an evacuation may be passed, and recovery take place. In some, after the continued use of opium, diarrhoea is produced, and may become so severe as to require remedial measures.

Change of position has in some instances appeared to produce benefit, and has been followed by recovery ; but whilst this may sometimes occur, we have witnessed the injurious effect of moving the patient when the peritoneum is intensely congested, and perhaps inflamed from the great distension ; death has in several instances quickly followed.

The application of cold water, or the exposure of the surface of the abdomen to the air, has been sometimes advantageously tried. In one of the cases which I have narrated, the patient said, whilst the abdomen was exposed, that he felt that something had slipped, and in a short time an abundant faecal evacuation was passed, and recovery took place.

In the absence of relief by these means, the question of surgical interference becomes one of anxious consideration ; after death from internal strangulation, the obstruction has been found so simple in character, that with great facility it might have been divided, and perhaps life saved. In several instances, Mr. Hilton has attempted this mode of relief, and with an amount of success which perhaps warrants further trial ; but it must be borne in mind, first, that the peritoneum is already inflamed, or in a state of intense congestion, and that general peritonitis is almost certain to follow ; secondly, that there is great difficulty

in the diagnosis, and that some recover from apparently a dying condition. A very interesting communication on this subject was read at the Hunterian Society, a short time ago, by Mr. Hutchinson, to which we must refer, and to the papers of Mr. Hilton, in the Guy's Reports of 1852. The operation of opening the colon in the loins is not of a very formidable character, and it has in many cases prolonged life; it has been especially performed in cancerous disease of the rectum, to relieve either the obstruction, or the severity of the pain. The diagnosis is here more evident; but in others, where, with some amount of certainty, we have indications of obstruction in the sigmoid flexure, it may be employed, we doubt not, with much success. In the case recorded by Mr. Hilton, in the paper just referred to, the relief was exceedingly marked, and the life of the patient prolonged for several months. In a valuable paper by Mr. Cæsar Hawkins, in the 'Transactions of the Royal Medical and Chirurgical Society,' the result of operation in forty-four cases of stricture of the colon or rectum is recorded; in ten cases death took place within forty-eight hours, in twenty-one within five weeks, and thirteen recovered; of these six died in six months, and nine survived more than one year.

The descending colon may be opened in many cases with facility, without dividing the peritoneum; when the cæcum or ascending colon is diseased, such relief would of course be impossible. In many cases, however, these operations have been deferred so long, that peritonitis has already arisen from the extreme distension, and the skill of the surgeon is then placed under the most disadvantageous circumstances. In the *treatment of intussusception*, every possible means should be employed to quiet the propulsive action of the intestine; but much good may be effected by local means; if the bowel be felt in the rectum, it may sometimes be returned by the introduction of a bougie, or of a candle; and in other cases gentle distension by the injection of warm water, or inflation by air, has apparently produced a return of the bowel.

CASE CLXXXII.—*Internal Strangulation of the Ileum. Band of Adhesion.*—Elizabeth B—, æt. 52, was admitted, March 10th, 1857, into Guy's Hospital. She was a thin, married woman, of sallow complexion, and had had a family. For twenty years she had had occasional pain in the left side,

the bowels had generally been confined, but she had not had any attack like that for which she was admitted.

On March 5th, five days before admission, after breakfast, she experienced a sudden pain in the abdomen; it commenced about the navel, but soon extended over the whole abdomen; vomiting came on an hour afterwards; the bowels had been open slightly the same morning, but had been confined on the previous day. From that time no evacuation took place, the vomiting continued, the abdomen became tympanitic, and moderately distended, and there was slight tenderness. The pain in the abdomen came on in paroxysmal attacks, but was generally absent when she remained quiet; the vomited matters continued bilious, and the urine abundant.

On March 10th, the countenance was expressive of considerable distress, but calm and resigned; the eyes were sunken; the abdomen was slightly prominent in the centre, but not laterally; it was tympanitic, but free from tenderness; the pulse was sharp, the respiration normal, and the urine was abundant. Purgatives had been given, and enemata administered, but she had vomited the former. The vomited matters continued bilious.

She was ordered calomel and opium of each gr. j every six hours, a soap injection to be administered, and a linseed poultice to be applied to the abdomen. As food, beef-tea and ice were allowed.

March 12th. The countenance was more haggard, and the eyes were more sunken; the vomited matters were thick, green, and offensive, but not stercoraceous; the pulse was more compressible; the abdomen was in the same state as far as external appearance, and it was still free from tenderness; there had been no relief from the bowels, and no flatus passed; the urine continued abundant; she had had a restless night, and suffered occasionally from hiccup. To continue the pills.

13th. The symptoms continued the same. To have injections of beef-tea, and to take gr. j of opium without calomel every six hours.

14th. There was no improvement. There was neither pain nor tenderness in the abdomen, nor was it more distended; she was not disturbed by vomiting, but was partially under the influence of opium; she was drowsy, the pupils were contracted; respiration, 16; pulse, 104; rather fuller, and compressible.

16th. She was evidently dying; there had been no change in the symptoms for the better.

17th. At 3.15 in the morning, she died, rather less than twelve days from the time of the strangulation.

Inspection took place about twelve hours after death. On opening the abdomen, by a small incision about four inches in length from the umbilicus, the small intestines were found moderately distended, but on the fingers being passed towards the pelvis the strangulated bowel gave way, and some fecal extravasation took place; when the parietes were fully divided the whole of the peritoneum was found inflamed; it was dry, and deep red lines existed at the points of contact of the intestine; the stomach and transverse colon were moderately distended; the small intestine was still more enlarged. In the pelvis several coils of small intestine were found almost black in colour; there was fecal extravasation, but this probably only took place after

death ; at the site of the right internal abdominal ring was a roughened and injected state of the peritoneum, as if adhesion had existed ; and there was a similar condition also on the right side. On turning aside the small intestine, a firm band of adhesion, round and dense, was found to extend from the region of the cæcum to the margin of the pelvis, at the termination of the sigmoid flexure ; through this loop several coils of ileum had passed, and had become strangulated. The band of adhesion passed from the mesentery of the ileum to the mesentery of the sigmoid flexure, and it appeared to be the free perforated margin of the latter mesentery ; the band was thin, and contained vessels ; and it was doubtful whether it was really a band of inflammatory adhesion, or a part of the sigmoid mesentery, which had become thinned and perforated, and so presented an abnormal and free edge. The strangulation was four feet from the cæcum, and nearly two feet in length. The mesentery of the strangulated part was infiltrated with blood ; its peritoneum was almost black, and in several parts it was sloughing. The mucous membrane at the upper end of the strangulation presented an extensive slough, and the coats were destroyed ; at the lower end the sloughing was rather less extensive and advanced.

The coils of intestine contained within the adhesion, were united by moderately firm lymph.

The appendix cæci was perfectly free ; below the band the small intestine was contracted, so also was the sigmoid flexure, but the transverse colon was moderately distended with flatus. The stomach was not at all dissolved ; but the whole of the mucous membrane was intensely congested with very minute arborescent vessels.

The liver, kidneys, and spleen, were healthy. In the lower lobe of the left lung there were several lobules, in a state of red and gray hepatization ; and the whole of that lobe was in a state of early pneumonic consolidation ; the other lung was healthy. The heart was normal ; its right cavities were filled with blood.

In this case, it is probable that a portion of intestine had existed within the band for some time, for adhesions had evidently existed between that part of the ileum and the parietes near the inguinal canal, and occasional pain had been experienced in the abdomen ; the distension of the incarcerated part and the intrusion of other coils led to strangulation.

It was diagnosed before death, that the obstruction was in the small intestine, from the moderate distension, the short time that elapsed before vomiting came on, and the character of the ejected matters. The quantity of urine did not assist us here ; the vomiting was moderate, because purgatives and irritants were avoided ; the distension also was rather in the central part of the abdomen, and the transverse and descending colon could not be traced, as in obstructed sigmoid flexure. As to the treat-

ment, I believe it was most judicious after admission into Guy's Hospital, and that life was by that means prolonged for several days, and the patient spared intense suffering. The opium quieted peristaltic action, and if violent vomiting had existed, perforation would probably have taken place at an early period. Considerable injury would have resulted from the administration of solid mercury.

In reference to opening the abdomen, if it had been attempted at a very early period, the band might perhaps have been divided, but during the latter days of life, the intestine was in a semi-gangrenous state, and the operation would probably have been hastily terminated by the rupture of the strangulated bowel.

CASE CLXXXIII.—*Colic, Lead ? Simulation of internal Strangulation. Recovery.*—A young man, æt. 22, badly nourished, who had been residing in Rosemary Lane, was admitted August 21st, into Guy's Hospital. He was pale and desponding, and had been suffering severely during eight days. He appeared to earn a scanty livelihood as a porter, and on August 14th, after taking his breakfast in his usual health he lifted about three quarters of a cwt. upon a cart, when he felt a sudden pain below the left hypochondriac region; he, however, went to his work, but was taken back, "doubled up," as he described it; after a few hours, vomiting came on, and both pain and vomiting continued till admission; he had not had any action from the bowels, although repeated doses of medicine had been taken, nor had there been any hiccup. He complained of severe pain across the umbilical region; the abdomen was neither hot, nor tender on pressure; there was some distension laterally, and in the position of the transverse colon, otherwise it was contracted. The tongue was clean and pale; the pulse eighty, and tolerably full in volume. He had passed but little urine, and neither blood nor mucus from the bowels. There was no hernia, but along the gums a dirty line which somewhat resembled lead.

For three months he had been a teetotaller, and he had had occasional pain in the abdomen, but no constipation.

A soap injection was administered, and calomel gr. v, with opium gr. iss, given as a pill. On the 22nd and 23rd there was no relief from the bowels, no medicine was administered. On the third day after admission the bowels acted slightly, castor oil was then given, and was followed by more active remedies. The bowels acted, and he left the hospital in a few days comparatively well.

This case was probably one of lead colic, in which the symptoms came on suddenly after exertion; it somewhat resembled internal strangulation, but the abdomen never became distended;

the importance of not following too active a plan of treatment was also shown, the vomiting became much less after the purgative medicines had been left off; the calomel and opium with enemata were used once; and on the third day the bowels were acted upon.

CASE CLXXXIV.—*Internal Strangulation? Restored.*—William G—, æt. 27, applied among my out-patients at Guy's Hospital, January 30, 1857. He was a spare muscular man. On the 24th, whilst lifting a sack of coals, he stated that he felt something give way in the abdomen, and pain followed immediately below the umbilicus; vomiting came on, and his food was rejected directly after being taken; the bowels had not acted. The abdomen was tense, supple, free from tenderness, but there was a sense of faintness, and he had pain below the umbilicus, the tongue was clean, the pulse quiet. He was ordered by Dr. Hughes, under whose care he was admitted, of opium gr. iss three times a day, and a turpentine enema. The enema brought away a solid fecal evacuation, but the vomiting continued, and he still complained of pain about the umbilicus; he passed between two and three pints of urine. On the end of February the bowels were freely acted on, and on the following day he left the hospital.

The symptoms presented in this case were of a serious character, but it was doubtful whether any internal strangulation or twist existed. The symptoms closely resembled those presented by internal obstruction, but they very rapidly subsided; we fear that there would be a return of these symptoms, but with less amenability to treatment.

CASE CLXXXV.—*Internal Strangulation. A Loop of Small Intestine passed into a hole in the Great Omentum.*—J. D—, æt. 45, was apparently in the enjoyment of good health, till Monday morning, November 29th. On that day he alighted suddenly from a chaise, to fetch his master's carpet-bag; at the same moment he felt sudden pain in the abdomen, low down in the right iliac region; about noon he began to vomit, and the vomiting recurred frequently. He had never suffered from any irregular action of the bowels, and had previously had good health. He was bled, and calomel with purgatives administered, without any effect. Enemata were returned with fecal odour. On the third day the abdomen was moderately distended, but free from tenderness; the pulse was 84; and the tongue was injected and fissured. There had been no action from the bowels, and the vomiting continued.

On the *fourth* day, there was no change in the symptoms; he was placed in a warm bath, and water injected into the rectum. Whilst in the bath he became much worse, collapse came on, and death followed in five hours.

Inspection.—The peritoneum contained thin fecal fluid. The coils of the

large intestine were lying in front of the omentum, which descended into the pelvis. The small intestine was adherent to the anterior abdominal parietes, and air was found to escape from a perforation in the small intestine. A loop of ileum, six inches from the cæcum, together with the mesentery, had passed through an opening in the great omentum, and had led to the fatal strangulation and subsequent perforation. There was no ulceration in the whole of the canal.

In this case we had sudden occurrence of symptoms; the position of the pain indicated the seat of the disease; the vomiting came on a few hours after pain, indicating affection of the small rather than of the large intestine; the case showed that although the abdomen was free from pain and tenderness, the movement required to place the patient in a warm bath, and the injection of warm water into the rectum, were not free from danger; they hastened fatal perforation and peritonitis; how much more easily would such an effect have followed more sudden and violent exertion!

CASE CLXXXVI.—Internal Strangulation of the last eighteen inches of the Small Intestine, fatal after thirty-eight hours.—Henry W—, æt. 19, had been employed as a lead and colour manufacturer in Tooley Street; he had had colic a year previously, but at the time of admission into Guy's Hospital no trace of lead existed on the gums. He was of pale complexion, with light hair, and he enjoyed his usual health till Sunday, July 28th, at 7 p.m., when soon after drinking some beer, he was seized with pain at the lower part of the abdomen, towards the right side. About an hour after this he had moderate action of the bowels. Sickness came on about 9 o'clock, and during the night he vomited whatever he took. An injection, administered before admission, came away with scarcely a tinge of feculent matter. He was brought to Guy's at 10 p.m., July 29th, and placed under Dr. Barlow's care, in an almost pulseless state; pulse 144, the face and extremities were cold, there was frequent eructation, the abdomen was rigid, tympanitic, and very tender on pressure; the tongue was flabby; no urine had passed; and his respiration was entirely thoracic.

At 9 a.m., on the day after admission, the abdomen was tense, slightly hollowed out in the right hypochondriac region, and tender on pressure. He was very restless, turning from side to side in bed, and his legs were occasionally drawn up. He had passed a disturbed night, with the same symptoms as on admission; his pulse could scarcely be felt. Shortly after this he was allowed by the nurse to rise up in bed, became faint, and died in about half an hour, thirty-eight hours after the commencement of the pain.

On examining the abdomen, several pints of bloody, dirty serum, were found in the peritoneal sac. The whole of the small intestine was much

distended; but several coils, corresponding to the last eighteen to twenty-four inches of the ileum, were in a state of approaching gangrene. The latter portion had become strangulated by a diverticulum from the small intestine, about one and a half inches in length, and by a band passing from the mesentery to the cæcal end of this pouch. The large intestine was less contracted than is generally observed in such cases. There was evidence of general peritonitis, lymph being effused between the coils of the intestine. The mucous membrane was continued into the pouch, and much imperfectly masticated cocoa-nut, and the remains of gooseberries, which he had eaten on the morning of the attack, were found in the intestine. The remaining viscera were healthy.

This case is worthy of being recorded as presenting peculiar difficulties in diagnosis; for whilst the urgent vomiting, the state of the abdomen, and the mode of the attack, pointed it out as one of mechanical obstruction, the great depression and rapid termination seemed to refer it to rupture of the bowel. These latter peculiarities, however, probably arose from the extent and completeness of the strangulation leading to speedy gangrene. The hernial sac was represented by the general sac of the peritoneum, which contained a considerable quantity of bloody serum. Although the pain was situated towards the right side, the symptoms were not those usually presented by cæcal mischief. The first indication of disease was pain resembling colic, after taking some malt liquor, and it is probable that the indigestible substances which he had eaten tended to excite distension, and irritation of the mucous membrane. A slight constriction of the canal may become so increased by irregular peristaltic action and over-distension as to become complete; thus a patient may for years suffer from slight attacks of pain, and from irregularity of the bowels, till after some indiscretion of diet he has a recurrence of pain and vomiting with constipation; no tenderness of the abdomen may be present, but rigidity of the abdominal muscles; the vomiting may continue, no further action from the bowels take place, but hiccough, rapid prostration of strength, and death speedily follow. The diagnosis in such a case is beset with difficulty. Abercrombie records several instances of this kind, where there was adhesion without apparent narrowing of the canal, existing probably for years, till, from some unknown cause, complete and fatal obstruction took place; and cases repeatedly present themselves in

which organic disease of great extent has existed for a considerable period, and symptoms of obstruction are manifested only a short time before death.

CASE CLXXXVII.—Internal Strangulation. Death on the fifth day.—Edward J. T.—, æt. 27, was admitted under the care of Dr. Rees into Guy's Hospital, October 11th, 1861, and died on the 14th. He was a spare man, with a rather sallow and anxious expression, and his eyes were sunken. The skin was cool; he stated that he was quite well till Wednesday, the 9th, when, having returned to his work after dinner, sudden pain came on in the abdomen, close to the umbilicus; he sent for a dose of castor-oil, which at once produced vomiting; the vomiting continued till his admission on the Friday. The rejected matters had the appearance of fluid fæces, but the fæcal smell was not very manifest. The tongue was clean, the pulse compressible; there was no evidence of external hernia; the abdomen was rather small, and there was no distension of the colon; neither was there any tenderness, nor tumour, nor hiccough. Calomel and opium were given, and, after an injection, flatus was discharged; after a second injection flatus and some fæcal matter were passed; the pain ceased; during the twenty-four hours a pint of urine was passed. On the 14th collapse came on, and in a few hours death took place.

On inspection, the lungs were found in a healthy state; the heart was small, the right side was flaccid, the left was firmly contracted. On opening the abdomen a band of adhesion was at once seen constricting a large portion of the small intestine. The general peritoneum was clear, smooth, and shining; the constriction was formed by a hole near to the centre of the omentum, the edges of which were thickened; and the ends of the constricting fold were adherent to the brim of the pelvis. The stomach and about six feet of the jejunum were distended; the rest of the jejunum and the ileum nearly to the cæcum had passed through the opening. The constricted part was contracted; its peritoneum was partially inflamed, and in one part softened, but there was no perforation. The colon was small and empty. The kidneys and the liver were healthy.

The symptoms of strangulation in this case were suddenly developed; and it was evident that the small intestine was the affected part, from the early period at which vomiting came on, the absence of distension of the abdomen, and the small quantity of urine which was passed. I have usually regarded the passage of flatus as a favorable symptom, but in this case it had not its usual favorable import; however, the strangulation was at that time either not complete, or the flatus merely came from below the incarcerated intestine. The early period at which death took place was probably due, in part, to the great extent

of the strangulated bowel, which amounted to the whole of the ileum and part of the jejunum; but the cause of the sudden collapse, a few hours before the fatal termination, could not be ascertained, for no perforation existed, and the strangulated portion only had its peritoneal investment inflamed. The omentum had first become adherent, then atrophied and perforated, and on sudden movement or distension the bowel passed through the opening.

CASE CLXXXVIII.—*Internal Strangulation. Old Peritoneal Adhesions. Peritonitis. Suppuration.*—John D—, æt. 33, a labourer, was admitted under Dr. Wilks's care into Guy's Hospital, June 8th, 1860. Till the previous Sunday, June 3rd, he had felt in good health; and on the following day, at 1 a m., he was seized with pain in the abdomen, and vomiting shortly came on; aperient medicines were given, and the vomiting increased in severity; the ejecta became stercoraceous; but no action of the bowels took place. He had a sallow and anxious expression of countenance when he was brought to Guy's; the abdomen was moderately distended, and rather tense; there was slight pain; an injection on the previous day had brought away some fecal matter, but he was not sure that any flatus had passed; the urine was scanty; the tongue was slightly injected and furred; the pulse was compressible. A grain of opium was ordered every four hours; and a warm poultice was applied. On the 9th the countenance was less anxious; but the eyes were sunken; the pulse quiet, the skin normal, the abdomen was not distended, and the muscles were hard and rigid, especially around the umbilicus, where he complained of pain; there had been no action from the bowels, nor had flatus been passed; the urine was moderate in quantity; the vomiting of offensive matter continued. 11th. There had been no action from the bowels, nor had he passed flatus. The stercoraceous vomiting continued, and two pints of urine were discharged. He had slept a little during the night; but the pain came on in paroxysms, and was of a griping character. The abdomen continued rigid, and the muscles contracted; there was slight tenderness towards the right iliac region, but none in the umbilical region; the tongue was red at the tip and edges; the pulse was sharp. Dr. Wilks had directed that he should be placed in a warm bath, and cold water poured upon the abdomen. On the 12th, the patient was in greater distress, and in more severe pain; he had had no sleep, for as soon as he dozed sudden pain darting through him, at once awoke him; the pulse was 100, hard; the tongue was rather dry, the vomiting more severe, and stercoraceous; the abdomen was not distended, and still rigid; the pain was especially situated in the umbilical region and about the cæcum. Opium gr. j was continued every four hours; and a long tube was passed and water injected into the bowel. On the 13th he was in still greater pain. Six pints of water had been thrown up three times; and the dulness consequent on the injection extended apparently as far as the cæcum; some scybala were brought away, and the vomiting became less.

On the 16th, he had not vomited for three days; on the previous day a six-pint injection was again thrown up, and some scybala passed; flatus was discharged; the pulse was quiet, the skin cool; the urine was abundant; he had suffered during the morning from severe twisting pain in the region of the umbilicus, and the pain continued; but his general expression was much improved—more cheerful and less haggard; he was able also to take beef-tea. To continue the opium.

During the afternoon of June 16th, on the thirteenth day of obstruction, the bowels acted. On the 17th, the bowels acted twice. On the 18th, he was cheerful, but suffered occasional pain in the abdomen. In a few days he was considered sufficiently well to leave the hospital.

On October 17th, 1860, he was admitted into St. George's Hospital, under Dr. Fuller's care, and I am indebted to the kindness of Dr. Dickinson for the following report:—The abdominal pain had never entirely left him since he had been in Guy's Hospital. "When admitted (October 17th) he was much emaciated and sallow. The abdomen was somewhat tympanitic, and slightly painful, but not tender. The bowels were regular; the motions pale. The urine contained phosphates in excess; but was in other respects natural. The sounds of the heart were normal. The pulse was weak, 88; the tongue slightly coated. He was allowed good diet, and had a draught, at intervals, containing soda, taraxacum, and gentian. 19th. The motions remained pale; and the bowels were generally torpid, requiring a laxative. He had pain in the abdomen, which he described as cramp, which was worse when the bowels were confined. The tympanitic swelling varied much. 23rd. The appetite remained pretty good. On the night of the 24th, he was kept awake by pain; and in the morning his tongue was glazed, and the appetite gone. The abdomen was much distended. He was now ordered (25th) to take every night a pill which he had before taken at times (opii gr. $\frac{1}{4}$ hydr. chlor. gr. iij). Hot fomentations did not afford any relief. 30th. The pulse was now rapid, the tongue red and dry. As the other symptoms continued, turpentine fomentations were applied, and the pill given twice a day, as well as effervescing saline medicine, and wine. Under these measures, he appeared by the 6th November to be somewhat better. He was still very emaciated, and had an anxious expression of countenance, but he was stronger. The pill was now given every night only, and a draught containing taraxacum, chloric ether, and ammonia. On the 8th, he was more prostrate, and the pulse quick and irregular. Vomiting came on and continued all day, leaving him at last faint and sinking, in which condition he died, in full possession of his faculties.

Inspection was made sixteen hours after death. The body was much emaciated; the abdomen very tympanitic. The brain was healthy. There were extensive old pleural adhesions on both sides. The lungs and heart were healthy. All the opposed surfaces of the peritoneum were closely united by old adhesions. In front the parietal peritoneum was thus united to the great omentum, which was of great extent and much thickened by old inflammation. It was closely adherent below to the walls of the abdomen near the pelvis, so that on dissecting off the abdominal walls, nothing else was seen. On cutting through the omentum, a large collection of thin fetid

pus was found, bathing the intestine; there could not have been less than a quart of this fluid. The small intestines were much convoluted, and in some places greatly distended. They were vascular, and their surfaces smeared in many places with recent lymph. Beside these, there were old adhesions between neighbouring coils. The small intestines were carefully traced down from the stomach (which was itself collapsed, but healthy), but no obstruction, nor trace of obstruction, could be found either in it or in the large intestine. The coats of the bowel were in many places much thickened, so as to resemble tripe. No morbid condition of their mucous membrane was anywhere found. The mesenteric glands were healthy, and near the ileo-cæcal valve was a large chalky mass, which appeared to have originated in a diseased gland. The liver and kidneys were healthy."

When this patient was under the care of Dr. Wilks he was suffering from the symptoms of internal strangulation of the small intestine, as shown by colic, with constipation and stercoraceous vomiting, &c.; and from the subsequent inspection, which revealed the presence of old peritoneal adhesions, it would seem probable that the obstruction arose from this cause, namely, the pressure of adhesions, perhaps rendered complete by enteric inflammation, as in Case CXVIII, in which slight thickening from a diseased mesenteric gland was sufficient, after indigestible food, to determine obstruction. The thickening of several portions of intestine indicated the presence of chronic impediment in the transit of the fecal contents. The cause of death appeared to be exhaustion from recurrent peritoneal attacks, terminating in suppuration. The advantage of an opiate plan of treatment, with enemata, was shown in the subsidence of the severe symptoms of the first attack.

CASE CLXXXIX.—*Lead Colic. Internal Strangulation of the Intestine from old disease of a Mesenteric Gland, and subsequent Fibroid Contractions.* (Reported by Mr. George Eastes).—Charles S —, æt. 29, was admitted into Guy's Hospital, under Dr. Rees' care, November 8th, and died on the 20th. He had been a painter by trade, but had left that occupation to become a chimney-sweep, seven or eight months before his last illness, on account of the attacks of colic from which he suffered; but even when a child he frequently suffered from pain in the abdomen. The first attack of this sort occurred three years previously, but that attack had been preceded by many threatenings; the next was twelve months later; in six months more, a third time the colic came on; and from that time repeated attacks were suffered from, the intervals of freedom from pain lessening in duration; so that during five months prior to admission, he was quite laid aside, on account of the abdominal pain and distress being almost constant. During

the latter part of his illness he had been much annoyed by vomiting; this symptom had harassed him continually, and it had generally come on about two hours after food, of whatever kind the food might be; the bowels were meantime costive, so as only to be opened by injection; the pain gave him no rest, by night nor by day, and it was accompanied by gurgling and spasm of the abdominal walls; the peristaltic movement of the intestine became visible. The patient described the passage and quantity of urine as free. He did not suffer from headache, nor from pain in the joints; but he had a well-marked lead line on his gums. His complexion was dark, and he had a large quantity of black hair; his countenance was anxious; the conjunctivæ were dingy; the tongue was red at the tip and edges; the pulse 68. The abdomen was not tender during the intervals of quiet; but as soon as the pain came on, the rectus muscle was drawn up into distinct knots, and then the abdomen became so tender that he could not bear even the bedclothes to rest upon him. He was not always sick, but generally vomited for a day or two at intervals of two or three months. The appetite was pretty good, but for four months he had taken only bread and arrowroot; meat was almost certain to produce vomiting, soon after taking it. His bowels were much constipated, acting only every three or four days; and the evacuations were hard and scybalous. There was slight difficulty in micturition, and the urine contained lithates.

On November the 8th he was ordered sulphate of magnesia ʒss, dilute sulphuric acid ℥xv, syrup of orange ʒss, and ʒj of water every six hours. On the 9th, the bowels had been once opened, but he had vomited much since admission, and had suffered from intense pain in the abdomen. Calomel and opium, of each a gr., were given every four hours, and a simple enema administered night and morning. On the 10th and 12th the vomiting continued; and opium gr. j was given every six hours, and chloroform liniment applied to the abdomen. 14th. The vomiting ceased for two days; a castor-oil injection was ordered, and the pills repeated every four hours. 15th. Vomiting again returned; the pain came on at intervals of half-an-hour; previously it came on every five minutes. He had had a little sleep. The enemata produced action from the bowels. During the next few days the symptoms remained as before, and twenty leeches were applied to relieve the intense abdominal pain, but without beneficial result. After a night of agonizing pain, he expired on November 20th, at 3 a.m.

Inspection was made thirty-six hours after death. The thoracic viscera were healthy. *Abdomen*, there was general and acute peritonitis; nearly the whole of the ileum and jejunum were almost of a black colour, and greatly distended, and in several parts the peritoneal surface had ulcerated, exposing the muscular coat, apparently from distension. The small intestine was drawn backward towards the spine from contraction of the mesentery; and on examining the ileum, near to the cæcum a hard mesenteric gland projected into the intestine, and had led to fibroid thickening and contraction; at this part the muscular coat of the ileum was much hypertrophied, and projected, like a nodule into the intestine; it closely resembled thickening of the pylorus from fibroid degeneration; the hypertrophy of the muscular coat extended some distance up the ileum, and contrasted remark-

ably with the thin muscular coat below the stricture; the mucous coat was also thickened, and slightly ulcerated; the peritoneal coat presented the ulceration before mentioned; the colon was contracted. The other viscera were healthy.

The constriction in this instance apparently originated in disease of one of the mesenteric glands during early life, fibroid thickening took place at that part of the mesentery, and led to partial and afterwards to fatal occlusion and peritonitis. The intense colic from which the patient suffered, was produced by the irregular peristaltic and spasmodic efforts to overcome this obstruction; attacks of colic appear to have come on in early life, before he had adopted the trade of a painter; afterwards it is probable that the lead aggravated the colic. The circumstance of chronic poisoning by lead, the blue line along the gums, tended to obscure correct diagnosis; the intensity of the colic, however, the intolerance of pressure, and the supervention of peritonitis, were all opposed to simple poisoning by the absorption of lead.

CASE CXC.—Mechanical Obstruction terminating favorably after seventy-eight hours.—For many of the particulars of the following case I am indebted to my friend Dr. Gull.

J. S—, æt. 33, a coal porter on a wharf, rose on the morning of the 26th June, 1850, in his usual health, and before going to his work, went, as his habit was, to stool, and had a good evacuation from the bowels. About half-an-hour afterwards, whilst stooping to fill a sack, he was suddenly seized with a sharp pain across the abdomen in the hypogastric region, accompanied by a sense of constriction. He was obliged to leave his work and to go home; in a short time he began to vomit, and after the attack was unable to pass anything downwards. He was treated by Mr. Mitchell, of Deptford, but without effect, and on the evening of the following day, forty hours from the accession of the symptoms, he was sent to the hospital with a note, saying that no hernia could be found, but that an internal obstruction was suspected. The assistance of Mr. Cock was obtained, who examined all the outlets, but could detect no protrusion. On admission he had the usual symptoms of strangulated hernia, urgent vomiting, anxious countenance, pulse rather frequent, the temperature of the surface was depressed, the abdomen rigid, and rather tumid, and slightly tender on pressure, urine small in quantity, and high coloured. He was ordered a grain of opium every four hours, and to abstain from relieving his thirst. The report of the third day at noon was, that he had passed a restless night, vomiting continually. He was seen early in the morning, everything was interdicted, even to cold water, and he was then better, the paroxysms of pain in the

abdomen being less urgent. As he was under the influence of opium the dose was diminished to half a grain, and a copious enema of salt and water was thrown by a long flexible tube into the rectum; it passed up readily, but without bringing away any feculent matter. In the evening he was restless, his countenance was anxious, the vomiting and other symptoms continuing as before.

Fourth day, eight o'clock a.m.—He vomited during the night in considerable quantity; the abdomen was tense, and coils of distended intestine could be partially traced, the peristaltic action rendering them prominent, with increase of pain in the abdomen, of which he complained bitterly. His countenance was still expressive of great anxiety, and the features were shrunk. He had passed about half a pint of urine, clear and well coloured; the pulse was accelerated, and diminished in power. He was ordered to go on with the opium. During the morning his abdomen was exposed for some time, whilst a sketch was made of its peculiar form, and the position and direction of the prominent convolutions, in order to determine more accurately the precise seat of obstruction; when suddenly, about noon, he expressed himself relieved, saying, that "something had given way within him," and this feeling was quickly followed by a copious flow of liquid feces inundating the bed. From this time he steadily recovered, the vomiting and hiccough at once subsided, and the face acquired a cheerful expression.

Certainly no cases present a less promising prognosis than those of mechanical obstruction of the intestines, nor has the enterprise of modern surgery yet succeeded in diminishing their mortality. The case here recorded presented points of no common interest; that it was one of mechanical obstruction there can be but little doubt, and if so, we had an instance of its spontaneous solution, and it answers in the affirmative the question, can we hope for a successful result in mechanical obstruction without surgical interference? From what we have seen in hospital practice, there is reason to believe that irregular peristaltic action following upon indigestible food, is not an uncommon cause of internal misplacement; but in the case here recorded, it came on after a night's fast, and before any meal had been taken in the morning. He rose as well as usual; the bowels acted according to his daily habit; he went to his work in good health, but whilst in a stooping position the pain came on. It need not be mentioned, that there was neither history nor trace of lead in the system, nor indeed were the symptoms such as arise from that mineral. The only remedy trusted to in the treatment was opium, but the happy termina-

tion of the case whilst the abdomen was exposed to the cold air, renders it probable that moderation of temperature had somewhat to do with the result. The application of cold has been suggested in such cases, and has much in theory to recommend it, and might be expected, in conjunction with opium, to effect all that mere treatment can effect. For, suppose a portion of intestine to have insinuated itself under any accidental band in the abdomen, by what means can we so well hope to liberate it, as by reducing its volume, and by allaying the vomiting? I would also suggest whether opium suppositories would not sometimes more efficiently promote the latter object than opium in the stomach. Of the opiate plan of treating intestinal obstruction too much cannot be said. It has both reason and experience on its side; and yet in the reports daily given of such cases, purgatives form generally the early part of the treatment; and are persevered in until the stomach will bear them no longer, serving only to exhaust the patient and increase the symptoms.

CASE CXCI.—*Internal Strangulation and Constipation. Subsidence of Symptoms. Death from Phthisis.*—(From the Museum Records.)—William H—, a man of middle age, was admitted into Guy's in 1829. There was obstinate constipation, vomiting of a stercoraceous character; but no hernia could be detected. The symptoms gradually subsided, but the patient died from phthisis several months afterwards.

On inspection, there were vomices in the lungs. The intestines were irregularly contracted. The appendix cæci was bound by adhesion to the brim of the pelvis, and several bridges of adhesions extended to portions of small intestine at this part; one of them was very long, and had apparently led to constriction, and to the previous symptoms of strangulation. No ulceration of the intestine existed.

Tumours sometimes become developed in the mesentery, and act as either the predisposing, or as direct causes of mechanical obstruction. Among the records of the inspections at Guy's, is that of a boy, aged seventeen, who, after a blow on the abdomen, two years previously, had gradual distension of the abdomen, fluctuation, vomiting, and constipation. The jejunum was found to be enormously distended. One portion of the mesentery near the commencement of the ileum contained numerous tubercles, supposed to be cancerous, and the contraction around these had led to obstruction; other tubercles were situated in the pelvis.

CASES OF INTUSSUSCEPTION.

CASE CXCII.—*Colic. Lumbrici. Diarrhœa. Intussusception of the Ileum and Ascending Colon into the Descending Colon.*—This case is fully reported by Dr. Hughes in the Guy's Reports of 1856.

Daniel D—, æt. 14, was admitted into Guy's, February 27, 1856, under Dr. Hughes' care. He had resided near the Tower, and had assisted his father as a tailor. His previous health had been very good till seven weeks before admission, when he was exposed to severe cold, and the following morning he was seized with acute pain in his abdomen, which continued for several hours; the pain returned on the following day, and similar paroxysms took place till admission, but at uncertain periods. The attacks generally came on towards evening, and sometimes twice in the day. He was free from pain from the 21st to the 25th, when he took some castor oil, and from that time he suffered from tenesmus, vomiting after meals, and loss of appetite. He described the pain as a twisting and tearing of his intestines principally about the umbilicus, and he detected "lumps" in the abdomen, which disappeared on the subsidence of the paroxysm, during which he lay with his legs curled up and his hands on the abdomen; the duration of the pain varied, and was frequently relieved by passing flatus. In the intervals he felt well. The appetite was capricious, and sometimes excessive. The bowels were open twice a day, the motions semi-fluid or scybalous. On admission he was much emaciated; the expression of countenance was one of distress; there was a white fur on the tongue, and the pulse was weak and compressible. Shortly after admission he voided a lumbricus teres with some mucus. Calomel gr. v and opium gr. ½ were given, and were followed by a senna draught. Poppy fomentations were applied, and milk diet ordered.

Repeated paroxysms of severe pain came on during the next fortnight; but in the intervals he was able to go about the ward. Calomel and opium, purgatives and enemata were ordered; diarrhœa then supervened with tenesmus. On March 15th he was suffering from severe pain, the tongue was coated with a white fur, and the pulse was quick and compressible. The abdomen was distended; coils of intestine were visible, and there was tenderness. He vomited a considerable quantity of green bilious fluid, and the alvine evacuations were of a dysenteric character, consisting of bloody mucus without fecal matter. Three leeches were applied to the anus; and starch and opium injection used; a draught consisting of linseed oil and tincture of rhubarb was given; and small doses of opium with mucilage and mint julep were repeated. There was, however, no relief to the symptoms, and the vomiting became more severe; another lumbricus was also ejected. One coil of intestine in the left iliac region became very manifest. Opium was given, with some relief to the vomiting and pain, but on the 23rd the symptoms of peritonitis became suddenly aggravated, and he died on the following day, twenty-seven days after admission, and eleven weeks after the commencement of the attack.

Inspection.—The body was badly nourished. The lungs and heart were healthy. The abdomen was considerably distended. On opening the peritoneum the descending colon was found to be enormously enlarged and full; so also the sigmoid flexure, which made a great curve nearly to the right side of the abdomen. The transverse colon could be traced in a similar state to the right side of the median line; it was thrown into transverse folds, and the ileum was found to be pushed into it. The cæcum and ascending colon were entirely intruded. The ileum which was left was much distended; a great part of the jejunum, however, was collapsed, and situated behind the transverse colon and stomach, in the position described, as the sac of the lesser omentum. It occupied this position either from the congenital looseness of the colon, or from its meso-colon having been drawn aside by the intussusception; the foramen of Winslow was normal. The general peritoneum was intensely injected, and was covered with lymph, and there was general acute peritonitis; the small intestine, however, which was situated behind the stomach, was not inflamed.

The stomach contained semi-feculent fluid; but the duodenum was normal. Several lumbrici were found in the jejunum; the ileum presented towards the commencement of the intussusception an ulcer about half an inch in diameter, which much congested at its margin; the intestine was full of yellow fluid fæces. On tracing the intestine onwards the lower part of the ileum, the cæcum, and ascending colon were found pushed into the descending colon. It could be felt within the large bowel, and reached into the rectum, within a few inches of the anus. Opening the sigmoid flexure and rectum, the termination of the intussuscepted portion was observed, almost black, but surrounded by semi-fluid fæces; the apex of the invaginated portion was very tense, its opening, which would admit the little finger, was marked by a fissure towards one side on account of the contraction of the mesentery. Turning aside the bowel it was found to be convex and twisted from the dragging of the mesentery, and at the concave side was a large irregular ulcer at the most tense portion. (Plate IV, fig. III.) In the sigmoid flexure, which was considerably distended, was a small opening into the peritoneal cavity, which had set up general peritonitis; at the other extremity of the intussuscepted portion the finger could be easily passed round the bowel, although there was commencing adhesion from the effusion of lymph. The liver, spleen, kidneys, &c., were healthy.

This case was one of peculiar interest, on account of the obscurity of the character of the disease; the colic appeared to be due to the lumbrici, but the severity of the symptoms, the intense pain, the purging of bloody mucus, the almost incessant vomiting, and the distended coils of intestine, indicated a more serious abdominal lesion. The disease lasted eleven weeks, and it is probable that the intussusception continued during that period, at first perhaps slight in extent, but gradually increasing to a greater degree. The canal did not become entirely oc-

cluded till near the fatal termination; and it is possible that the intussusception may have become *partially* restored with the relief of the symptoms, and at each fresh paroxysm the intestine may have been pushed further onwards. The cause of death was peritonitis, consequent on rupture of the sigmoid flexure; and the exciting cause of the intussusception was probably the irregular peristaltic action consequent on the lumbrici associated with unusual and perhaps congenital freedom of the cæcal mesentery. As far as can be judged by a post-mortem consideration of treatment, opium was the most desirable remedy, with rest, bland nutriment, and the avoidance of any purgative medicines; but with such an extensive intrusion of intestine no remedy would probably have been effective. The gangrenous condition of the enclosed bowel had a reparative tendency, which in many cases has resulted in recovery to comparative health.

I have observed instances in which symptoms very similar to those manifested in this case have gradually subsided, and the patient recovered; such was Case CLXXXIII.

CASE CXCIH.—Intussusception. Recovery. Cæcum and the whole of the Ascending Colon passed per Rectum. (See Prep. in Guy's, 1875.)—W. P—, æt. 6, a patient of Mr. C. King's, City Road, in 1852. The previous health of the child had been good, till he was attacked with œdema and discoloration of both legs; these symptoms soon subsided, but constant vomiting came on, with constipation and pain, and with tenderness of the abdomen, particularly in the right iliac region; these urgent symptoms remained for four days, when convulsions and insensibility ensued. He remained in this condition for twelve hours, apparently dying; on the two following days he was a little better, the vomiting ceased, but constipation continued; during the next four days there was no change. Eleven days after the seizure, and five days after the cessation of the vomiting, he had an evacuation from the bowels, and the cæcum with the vermiform process and the ascending colon were discharged; when passed, the cylinder of the intestinal slough was complete. In a few days the leg became gangrenous, and was removed; the child did well, and completely recovered.

The symptoms of colic, in this as in the previous case, were very severe, and the recovery of the child very remarkable after the removal of the whole of the cæcum and ascending colon. It must also be noticed that constipation of an insuperable character was not one of the earlier symptoms; there was

evident impairment of the general health of the child, as shown by the œdema and discoloration of the legs, followed by gangrene; and it was suggested that unwholesome food, as ergotized bread, might have produced the disease. In cerebral irritation, also, we find a disposition to irregular peristaltic action of the intestine, and frequently after death from hydrocephalus, numerous portions of invaginated intestine are observed; here, however, the abdominal symptoms preceded the cerebral. The following cases present several points of great interest connected with this subject.

An emaciated man, æt. 28, under the care of Mr. Benjamin Phillips,* had been resident in a miasmatic district. He had suffered occasionally for weeks from an obscure affection of the digestive system; the abdomen was hard and tympanitic; there was frequent nausea, but vomiting rarely took place; the alvine evacuations were sometimes frequent and fluid, at other times they were natural; leeches were applied to the abdomen; the diarrhœa and nausea continued, the evacuations became greenish, and contained blood; and an elongated mass was found occupying the left iliac fossa. The patient had a constant disposition to sleep; and he died seven days after coming under Mr. Phillips's care.

On inspection there was found to be acute peritonitis, and an invagination of the cæcum and ileum into the transverse and descending colon. Several inches of the invaginated intestine were gangrenous, and the serous surfaces of the inclosed bowel were adherent: perforation had taken place.

In another case, reported by the same gentleman, the patient, æt. 31, had suffered for many months; there were sallowness of the skin, emaciation, a tympanitic state of the abdomen, and tenderness in the course of the descending colon and sigmoid flexure. In the left iliac region a tumour could be felt, considered by some to be impacted fæces. On inspection there was general peritonitis, the cæcum and ascending colon were not visible, and a cylindrical tumour was found in the iliac fossa; "two inches of the small intestine had penetrated into the cæcum; this turned upon itself, was introduced into the ascend-

* 'Medical Gazette.'

ing colon, which in turn had passed into the transverse colon, and all these parts thus disposed had reached the left iliac fossa." Several perforations had taken place.

In a case reported by Mr. Jon. Hutchinson, in the 'Pathological Transactions,' the symptoms of colic had existed for several months, and the patient, a young man, had sometimes swung himself on the steps of a ladder, as the only means of relieving the pain. The invaginated portion of intestine was found adherent, and the appearances evidently indicated that it had been so intruded for a considerable period.

CASE CXCV.—*Constipation. Subsequent Perforation. Peritonitis. Intussusception Restored?* (From the Museum Records.)—M. S., æt. 60, ten or twelve days before application had experienced sudden violent pain in the abdomen, with constipation; vomiting came on, but no hernia could be detected; by avoiding medicine the vomiting subsided. A dose of croton oil produced an evacuation, but without relief to the symptoms; the bowels were afterwards moved by castor oil; the symptoms of peritonitis returned, and the patient quickly died. On inspection a portion of small intestine was found, dusky and lurid, and patches of lymph were observed; on moving the intestines feces escaped. The discoloured portion was six to seven inches in length, and the mucous membrane was dark; the mesentery was also slightly discoloured, and greenish at that part. A defined line marked the diseased portion.

The appearances presented in this case were either those of an intussusception restored, which was the opinion of one who had had very great experience in pathological science; or, 2ndly, of internal strangulation; or, 3rdly, of local enteritis, as we have previously mentioned in speaking of that disease; or, 4thly, of a twist of the intestine on the mesentery, which had become partially restored. The last suggestion was, perhaps, the most probable. There was no evidence that either external or internal hernia had existed; and whilst it is very probable that cases of intussusception are restored, we scarcely feel warranted in asserting that invagination had taken place in this instance.

The following case is a remarkable one, as indicating one of the sequences of intussusception. It is from the 'Medical Gazette :—A patient, æt. 65, had constipation, violent pain in the bowels, and vomiting; in four days the pain ceased. It had come on August the 26th; on the 31st there were several offen-

sive dejections, and on September 5th forty-four inches of intestine were evacuated. The patient survived forty days.

On inspection the sigmoid flexure was wanting, and the cæcum and colon, seventeen inches in length, opened into a large faecal abscess, into which the rectum passed.

CASE CXCV.—*Phthisis. Intussusception of the Ileum. Peritonitis.*—James H—, æt. 16, a pale boy, was admitted into Guy's Hospital, May 23rd, 1860. For a year he had suffered from cough and phthisical symptoms. Eight days before admission he complained of pain in the abdomen, which was accompanied with vomiting. The pain commenced in the region of the cæcum, but the tenderness was at first slight; this symptom afterwards increased, and became more general as the indications of peritonitis were developed. There was great restlessness, and persistent bilious vomiting. The bowels acted once after injection, but there was no discharge of blood. Death took place on June 1st.

On inspection, phthisical vomices were found at the apices of both lungs; and the lower lobe of the left lung was in a state of recent hepatization. In the abdomen, both small and large intestines were distended; there were some lines of injection at the margins of contact between the intestinal coils. A foot from the cæcum the small intestine was quite divided to the mesentery; but extravasation had not taken place to any great extent, although no adhesions had formed. It was at once seen that the sloughing had arisen from intussusception of the ileum; and several inches of *separated* intestine were in a sloughy state. The sigmoid flexure and the parts below the intussusception were not collapsed, but were partially distended with flatus.

CASE CXCVI.—*Intussusception of Ileum. Perforation. Peritonitis.*—John S—, æt. 17, admitted into Guy's Hospital, under Dr. Barlow's care, September 17th, 1857, in a moribund state, and died in an hour or two. It was stated that he had suffered from insuperable constipation for a fortnight, with increasing distension of the abdomen, vomiting, &c. The first indication had been two weeks previously, when, on going to stool, he found nothing passed his bowels; and after that time he had only the smallest evacuation after injections. On admission there was peritonitis and faecal vomiting. The body was that of a strong, muscular man; the abdomen was much distended, and tympanitic.

On inspection, the thoracic viscera were found to be healthy. There was acute peritonitis, but only a small quantity of lymph had been effused. The small intestines were much distended, but the large were contracted. The obstruction was at once seen to be an intussusception at the lower part of the ileum; and upon raising this portion, an opening was seen in the gut, and faecal matter was escaping; a small quantity only of faecal matter was at first seen, so that the perforation probably remained nearly closed by being in contact with an adjoining coil, although the whole calibre of the intestine was torn through. The intussusception was found to be at three

feet from the cæcum, and arose from four to five inches of the ileum having passed into a lower portion. The contained part was in a state of slough, and was in shreds, as if it would soon have become detached. At the upper orifice some firm adhesions existed between the serous surfaces, but at the point of constriction these had separated. The opening of the lower part was almost closed, so that a probe could be scarcely introduced. The serous surfaces were closely adherent at the margin, where they passed the one into the other; and upon cutting through the included part, the serous surfaces were seen in like manner to be connected by lymph; and, as a considerable space existed, the lymph uniting them was of some thickness; the interior mucous passage was almost closed, only just admitting a probe. All the small intestines, as well as the stomach, were filled with fluid fecal matter, which in general appearance could not be distinguished from that in the colon.

This patient was dying when admitted, and there were no symptoms to enable us to distinguish this case of intussusception from one of internal strangulation of intestine. It is possible that, had rest been enjoined and proper treatment been adopted from the first, the contained slough might have been discharged without the separation of those adhesions on which the safety of the patient depended.

CASE CXCVII.—*Intussusception of Sigmoid Flexure. External Protrusion. Symptoms of Strangulation. Peritonitis. Death.*—Catherine F—, æt. 25, was admitted into Guy's Hospital, June 18th, 1857, and died on the 28th. She was a single woman, and in the October before her death began to suffer from prolapse of the anus; the prolapse was returned, but again came down, and from mistaken modesty she had neglected her complaint. Three weeks before admission, the bowel came down, and she was unable to return it, and at the same time constipation ensued. Still no advice was sought, and on admission she was found to suffer from strangulation; the bowels had not been moved for three weeks. The intussuscepted bowel lay for several inches outside the anus, but could be easily replaced; although the strangulation was not thereby overcome.

She died on the 28th, ten days after entering the hospital. The body was spare; the abdomen was tympanitic, but not excessively distended. There was acute general peritonitis. All the intestines were slightly distended, and full of fluid feces. Nothing abnormal was found till the pelvis was examined, where, low down behind the uterus, an intussusception was found, and at first it seemed so near to the anus as to be merely a prolapse; when, however, the intestine was removed it was found that the proximity to the anus arose from dragging down of the intestine, for when stretched out the invaginated part did not reach the anus by three inches; measuring from the line of constriction to the anus was nine inches, and the invaginated part measured half this length, making the commencement of the inverted bowel

eighteen inches from the anus, and therefore in the sigmoid flexure. The invaginated part was sloughing, and, as usual, slightly curved on itself by the dragging of its attachment. Other viscera were healthy.

These instances of intussusception present us with symptoms of severe colic, with vomiting and constipation, and often with tumour; 2ndly, they show that the pain and other symptoms are often paroxysmal; 3rdly, that the constipation is not always constant, but on the contrary, that diarrhoea is sometimes present; 4thly, that the discharge of blood and mucus occasionally takes place; 5thly, that the causes of death are perforation and acute peritonitis, or secondary faecal abscess; and 6thly, that the disease is cured by restoration of the parts, and sometimes by sloughing and separation of the invaginated portion.

CASES OF CANCEROUS DISEASE.

CASE CXCVIII.—*Villous or Epithelial Cancer of the Sigmoid Flexure, with Cancerous infiltration of Glands near the Gall-bladder.*—Ralph G—, æt. 44, a stout, plethoric man, who had served for fifteen years in the police force, had been employed at the station house, so that his life was a sedentary one. He had had good health, with the exception of slight attacks of rheumatism, till one year before admission, when after taking less than his usual exercise, his bowels became confined; he had, however, generally a motion every three days.

He was admitted into the hospital, under my care, July 3rd.

On June 20th he passed a solid stool, small in quantity, but without straining or pain; from that time nothing had been passed. He did not feel any uneasiness till the 23rd, when he felt pain and a sense of weight in his abdomen, and he vomited slightly. These symptoms passed off, but afterwards returned. He had hiccough at night, and his sleep had been disturbed; the appetite had failed, his abdomen had swelled, and he had some dyspnœa. Before admission he took various aperients, and had an injection of turpentine, but without effect.

July 3rd. The abdomen was much swollen, measuring forty and a half inches in circumference; it was most prominent in the position of the transverse colon, and tympanitic. This tympanitic resonance could be traced in the course of the colon, nearly to the sigmoid flexure. At that part he had slight pain, and stated that some months before he had had a similar attack. He had not had any discharge of blood, mucus, nor of air, per rectum; there was no pain on manipulating the abdomen, nor any increase of temperature; the pulse was quick and sharp, 98; the respiration was accelerated, the skin was perspiring, the tongue had a white fur upon it. To have a turpentine enema at once, and of the soap and opium pill, gr. v, three times a day.

July 4th. Vomiting took place at 5 a.m., the pulse was strong, 86; the skin cool; he had had no vomiting since the morning. To continue the pills, and to have the rue injection.

July 5th. He passed a considerable evacuation and felt much easier. He afterwards had some sleep, and was able to take some food; the pulse was feeble, 116; tongue more brown.

9 p.m. Calomel, gr. xij, were ordered to be taken, and the rue injection to be repeated.

July 6th. Passed a small quantity of fæces; the tongue was dry, and furred; the pulse 120, and sharp; the skin clammy; respiration 40. A long tube was introduced per rectum, apparently to eighteen inches; a pint and a half of water was injected; the enema caused considerable pain, and was retained for three hours, but failed to bring away any fæces. Calomel, gr. j, opium, gr. j, were given three times a day. Two eggs, and brandy, 3ij, were allowed.

9 p.m. The enema produced severe pain; the patient was restless; skin hot and covered with a clammy sweat; pulse, 140; respiration, 40.

July 7th, 9 a.m. He had passed a restless night, but had rallied somewhat; he did not complain of pain, but merely of sense of tightness and fulness. The whole abdomen was very tympanitic; except in the left iliac region; the pulse was 132, the respiration 36, and the tongue was brown and furred. The patient did not suffer from nausea nor from vomiting; the urine was moderate in quantity and high coloured.

2 p.m. Mr. Birkett could not detect anything on examination per rectum, and did not think the symptoms of insuperable obstruction sufficiently severe to warrant surgical interference. Ordered opium, gr. j, at once, and simple enema.

9 p.m. The patient appeared in the same condition as in the morning, he was prostrate and was covered with clammy perspiration. To repeat the opium, and to have an assafoetida injection, and oil of turpentine and linseed poultice to be applied to the abdomen. Brandy was allowed. The enema was administered with the long tube without causing any pain. The turpentine was applied for half an hour, and afterwards the linseed poultice; it produced slight smarting pain.

The patient gradually sank, and died 2-30 a.m. on the 8th, nineteen days after the commencement of the symptoms of obstruction.

Inspection twelve hours after death. The rigor mortis was well marked. The abdominal parietes contained a considerable layer of integumental fat. The abdomen measured round the umbilicus three and a half feet. On opening the peritoneal cavity, it was found to contain about three pints of opaque serum mixed with shreds of lymph; and the peritoneum was much injected, and was covered with spots of lymph. Both small and large intestines were enormously distended; this was especially marked in the cæcum and colon, as far as the sigmoid flexure, where was the seat of obstruction; the sigmoid flexure was distended and bound to the walls of the abdomen, the intestine then turned inwards towards the promontory of the sacrum where it became suddenly narrow at its union with the rectum. Externally the constricted mass felt hard, and after removal it was found that an

ordinary probe would scarcely pass. The obstruction was nearly an inch in length; on placing it in water, the surface was quite flocculent, resembling villous cancer. The intestine, both above and below, was healthy; above was a large quantity of fluid fæces; below, small scybalous masses. Near the gall-bladder were several glands, infiltrated with cancerous product. Other organs were healthy.

The microscopical examination of the diseased growth presented cells resembling columnar epithelium, but of greater size, and containing large nuclei. The whole of the flocculent surface was composed of cells of this kind, but no large cells, such as are usually found in epithelial cancer, were observed. They appeared rather to be modified columnar epithelium. The muscular coat of the intestine at that part was much contracted. The diagnosis in this case was from the first clear; the gradually increasing constipation, absence of pain, resonance, as far as the sigmoid flexure, with previous slight pain at that part, and the normal quantity of urine, all tended to show that the obstruction was at, or about the sigmoid flexure. It was a matter of regret that, in a case so favorable for surgical assistance, such means were postponed till fatal peritonitis came on; but the apparent mildness of the symptoms, the absence of vomiting, on account of the non-administration of drastic purgatives, led some to the supposition that the disease arose rather from impacted fæces than from an insuperable obstruction. The development of glands infiltrated with cancer near the gall-bladder was an interesting fact with this form of disease, which appeared to be of the character of villous or may be of epithelial cancer, in which there is less tendency to glandular infiltration.

CASE CXCIX.—Cancer of the Sigmoid Flexure.—Sarah O—, æt. 42, was admitted November 18th, 1856, and died the following day, at 8 a.m. In July she had received a fall, and on August 5th experienced pain in the region of the sigmoid flexure of the colon. The pain gradually extended over the whole abdomen; injections were administered which produced evacuations from the bowels, several days before admission.

When brought to Guy's Hospital she was too ill to give any definite statement in reference to herself. The countenance was anxious, the pulse was small and compressible. The abdomen was very much distended, and when exposed the position of the transverse colon was more prominent than other parts, and was tympanitic. The pain and tenderness were general;

vomiting was very distressing; an abundant quantity of urine was passed. Opium was given, and a warm poultice was applied; but she died the following morning.

Inspection was made about six hours after death. The thoracic viscera were quite healthy. The peritoneum was much injected, and the intestines appeared dry, from a delicate stratum of lymph upon them.

The colon was very much distended as far as the sigmoid flexure; the small intestines also were moderately distended. The stomach was healthy. Near the end of the ileum there was considerable congestion and several ulcers; these ulcers, however, were much more extensive in the cæcum. The canal there was enormously enlarged, and there was very general transverse ulceration, exposing the circular muscular fibres, as if ulcerated from over-distension; in some parts the muscular coat also was destroyed, and slight perforation had taken place in one spot, but without extravasation of fæces; the gut was more than nine inches in circumference. The appendix was filled with mucus, which was very slightly acid; and it was adherent in the long axis of the colon. The descending colon was very much distended as far as the brim of the pelvis, where it became suddenly contracted; and this part was adherent to the uterus and to a coil of small intestine. On separation, the intestine was found to be drawn in at that part, and hardened. On opening it, the little finger could be passed, and the canal above was filled with fluid fæces; at the constriction there were vascular prominent growths, corresponding almost to the position of the longitudinal bands; this section had a yellowish color, and showed that both the muscular and mucous coat were involved. On careful microscopical examination the surface was found to present a few villous processes, and the mass consisted of abundant nuclei and many compound nucleated cells, resembling some forms of medullary cancer; above the constriction was a smooth round opening, extending through the coats of the intestine into the peritoneum, but adhesions had formed between the uterus and coils of small intestine so as to prevent extravasation. The constriction was seventeen inches from the anus; below the stricture was some dry fæcal matter. The other abdominal viscera and glands were healthy.

CASE CC.—Cancerous Disease of the Sigmoid Flexure. Ecchymosis of Stomach. Ulceration of the Ileum. Contracted Mitral Valve.—Ellen H—, æt. 53, was admitted November 7th, 1855. She was a married woman, without family, and emaciated; she had been living at Shepherd's Bush.

Seven months before admission she had had severe pain at the lower part of the abdomen, and was compelled to desist from work; the pain came on four or five times a day; the bowels were confined, but had previously been regular. The motions were then very scanty, except after injections; she had sometimes had severe vomiting, and at times offensive matter was rejected; the urine had always been abundant.

The abdomen on admission was very large and tympanitic, but it was most prominent in the umbilical region; the tongue was clean; the pulse was small and very compressible. No abdominal tumour could be felt; and there was no tenderness. On admission enemata were administered, and

purgatives, which latter aggravated the symptoms. November 24th, opium was given, gr. j every six hours. This was followed by marked improvement, the stomach became quiet, and she was able to retain food.

November 30th. She was not so well, and complained of severe pain in the stomach; the tongue was small and contracted; the bowels were open freely; enemata had been administered, and opium given.

December 19th. She was much better; the abdomen was supple, not distended; and the bowels were open; she was free from pain, and had a good appetite; she took some porter and a chop, and wine. Opium, gr. j, was continued.

January 1st. For a week she had not been so well, and was obliged to remain in bed; the abdomen became painful, and the bowels constipated.

January 8th. She was dying; the pulse was intermittent; she died 10 p.m.

January 9th. *Inspection*, 2:30 p.m., seventeen hours after death. The body was extremely emaciated; the eyes were sunken; the abdomen was greatly distended. The parietes of the abdomen were thin. On opening the peritoneal cavity, an enormously distended transverse colon was found to occupy the whole anterior region of the abdomen; from the liver it passed down to the brim of the pelvis, then ascended nearly to the scrobiculus cordis, before it formed a second smaller curve, and became the descending colon. The large intestine was distended as far as the termination of the sigmoid flexure. Along the margins of the distended coils of intestine were lines of injection, and between some of the coils were delicate flakes of lymph. At the commencement of the rectum the intestine was contracted; and a drawing in of the coats of the intestine gave the part an irregularly puckered appearance; although thus contracted, the intestine at that part was readily moveable. The whole of the colon was distended with fluid bilious fæces; at the constricted part the intestine would only admit an ordinary quill; the constriction was one inch in breadth, raised, nodular, and deeply injected; the superficial portion was soft, and of a grayish colour; this rested on firmer iron gray structure, and minute masses of yellowish fat; the muscular coat was drawn in and lost at this part; but in the colon, both above and below the stricture, it was distinct.

On careful examination of this part (Plate III. fig. 5), the surface was smooth, and presented columnar epithelium, nucleated cells, and elongated nuclei (*a*); beneath the mucous membrane, which was itself dense, changed in character and fibrous, was a considerable quantity of firm, fibrous tissue, arranged at right angles with the intestine (*b*), and leaving interspaces filled with nuclei, but without nucleoli (*c*); still deeper, muscular fibre could be detected. There was no structure of an ordinary carcinomatous character. The nuclei were different from ordinary nuclei, not having well-defined cell-wall or nucleoli. They appeared like a coagulated blastema, in course of development into a fibrous structure.

In the termination of the ileum was an ulcer affecting nearly the whole of one of Peyer's patches, and the mucous membrane was entirely destroyed but the disease was of a different character from that in the colon; the rest of the small intestine was healthy. The stomach contained some black

mucus adherent to the membrane. At the cardiac extremity was a raised, black patch, covered with white substance, but merely affecting the mucous membrane. The follicles were evident, and contained some blood slightly blackened; but at the upper part of the membrane, where the capillaries were more numerous, there was an almost uniform black colour; it appeared that before death ecchymosis had taken place from the capillaries, and that after death the blood had become changed by the action of the gastric juice. At the lesser curvature was another black patch, but without the white substance on the surface; there, too, the follicles were beautifully distinct, some being marked out by being filled with changed blood; and that which had exuded from the superficial capillaries was blackened. The white substance consisted of cells and crystals. The liver, kidneys, and spleen were healthy; the gall-bladder was distended. There was no enlargement of the lymphatic glands of the mesentery; nor of those found in the abdomen and chest. The ovaries and uterus were healthy; the os uteri was small and conical. The lungs were healthy, but were slightly emphysematous. The heart was small; the mitral was rigid, contracted, and it would admit little more than one finger. The semi-lunar ganglia appeared normal.

In this case the obstruction was diagnosed to be at the sigmoid flexure, but the great emaciation led to the belief that there was more general infiltration of the glands. This was not the case, but the distension had produced ulceration of the ileum; nutrition was much impaired, and the diseased condition of the mitral interfered with the healthy action of the heart. The opium acted well, and its use was followed by marked improvement, and by action from the bowels; the administration of purgatives increased the vomiting and prostration.

CASE CCI.—Cancer of the Liver, of the Lumbar Glands, and of the Sigmoid Flexure.—Robert W—, æt. 32, was admitted September 19th, and died October 16th. He was a patten maker, and had lived in the Borough. Four months previously he had begun to feel pain; there were symptoms of indigestion, and afterwards severe pain in the right side. He became emaciated, but the abdomen was enlarged; the liver could be felt very distinctly on the right side, and nearly reached to the crest of the ilium. The pain in the side and across the abdomen became more severe, and he gradually sank. There was no indication of disease of the sigmoid flexure observed during life.

Inspection was made twenty-seven hours after death.—The body was spare, and slightly jaundiced. The chest was healthy, with the exception of the base of the right lung, where was a large patch about three inches in diameter, white in colour, situated on the surface of the pleura, and about one-eighth of an inch in thickness; this consisted of cancer, extending through the diaphragm from the liver; there were a few tubercles in the

neighbourhood; and one of the glands of the neck was infiltrated with cancer. The lungs, bronchial glands, and heart were healthy.

Abdomen.—The peritoneum contained about three pints of serum and pus; the liver was $9\frac{1}{2}$ lbs. in weight, and towards the diaphragm had the appearance of a large abscess; the surface was irregularly contracted from the development of masses of cancer. On section, nearly the whole gland was found to be involved, with scarcely any intervening gland structure; and these cancerous masses presented nearly every stage of degeneration; some had a soft, yellow centre, others a dark green slough, and in some the centre was semi-fluid. The lumbar glands were infiltrated; and at the termination of the sigmoid flexure was a small faecal abscess; the walls of the intestine were ulcerated, broken down, and infiltrated with cancer, and some of the contents had become extravasated among the cancerous exudation.

Here there was *no marked* constipation; the cancer was medullary rather than scirrhus or epithelial; there had been some pain in the part, but no obstruction. The patient was evidently wasting from organic disease; the liver was known to be diseased, and so slight were the symptoms of disease at the sigmoid flexure, that they were scarcely noticed.

The intestine may have been the part originally diseased, but of this we must speak doubtfully; the liver was that which gave rise to the more prominent symptoms.

The diffusion of the cancer to other glands was here well marked, and presented a contrast to that of scirrhus or epithelial disease.

CASE CCII.—Cancer of the Sigmoid Flexure. Constipation. Death on the twentieth day.—Margaret S—, æt. 36, was admitted into Guy's under the care of one of my colleagues, May 29th, 1847, and died June 4th. She had been a charwoman, had enjoyed good health, and had lived temperately. At the time of admission she had been subject to constipation for two months, and for thirteen days had had no evacuation; she had taken blue pill and black draught. She was depressed and anxious; the countenance was yellow and cadaverous, and she complained of tenderness across the umbilicus. The abdomen was tympanitic; the tongue was dry and brown; there was occasional vomiting, and a scanty flow of high-colored urine. A few scybala were passed by the bowels. Colocynth, with magnesia mixture, was given, and a soap enema was administered. The symptoms continued till the 31st, when a large quantity of pale urine was passed, and the vomiting and pain became distressing. On June 4th, the day of her death, the vomiting was stercoraceous.

On inspection, the peritoneum was found universally inflamed; the intestines slightly adherent. At the termination of the sigmoid flexure was a constriction which would allow water slowly to pass, but did not admit the

flexible tube. The peritoneum appeared corrugated, and the appendices epiploicæ drawn together. On opening the canal, there was an irregular growth from the mucous membrane at the sigmoid flexure, and the mucous coat was thickened. The intestine was filled with semi-fluid fæces. The glands were not affected; and the other organs were healthy.

The symptoms in this case indicated that the seat of obstruction was in the colon. The urine was at first scanty, but afterwards became more abundant. Peritonitis came on early, and was the cause of death.

CASE CCIII.—*Cancerous Disease of the Sigmoid Flexure of the Colon. Insuperable Constipation. Death on the tenth day.* (From the Museum Records, prep. 1854²⁴.)—Mrs. G—, æt. 55, who had resided at Spitalfields, had for fourteen years, after a difficult parturition, suffered from occasional pain in the abdomen; and for several years before death had had increasing constipation of the bowels; at last the obstruction became complete, and there was no evacuation for ten days; powerful purgatives were administered.

At the commencement of the rectum, the intestine would scarcely admit a probe, and appeared to be contracted by effusion between its coats; the structure had a villous surface; other viscera were healthy. The growth had apparently been *very slow* in formation, and its cancerous character was doubtful.

CASE CCIV.—*Cancerous Ulceration of the Sigmoid Flexure of the Colon. Constipation.*—For the particulars of the following case I am indebted to my friend Dr. Gull. The preparation is in the Museum at Guy's (1854²⁵).

Mrs. H—, æt. 60, in May, 1854, had an attack of diarrhœa, and a similar attack had occurred some months previously; from that time she had been troubled with flatulence and pain in the abdomen. The diarrhœa was relieved, but the pain continued. On July 22nd she had constipation, which was not removed by the use of castor oil, rhubarb, &c. There was no vomiting, the pulse was quiet and the tongue clean. Vomiting came on, on the 24th. The examination of the rectum discovered a hard mass high up in the recto-vaginal space. Opium and ice removed the symptoms. After five days the bowels were relieved, and she then went on very well till September 20th, when the bowels again became obstructed; enemata were used, and opium was administered; croton oil was rubbed into the abdomen. Purgatives were occasionally given, but in vain; after five weeks of complete constipation, symptoms of peritonitis came on, and she died. The operation of opening the descending colon was proposed, but the patient would not consent.

In this case diarrhœa alternated with constipation, a condition which is not unfrequent in disease of the sigmoid flexure.

CASE CCV.—*Cancerous Disease of the Sigmoid Flexure. Insuperable Constipation.* (From the Museum Records, prep. 1854.)—Donnel H—, was admitted with constipation of three weeks' duration in 1827. Powerful cathartics were administered, magnesia, colocynth, scammony, calomel, &c. He was free from pain and cheerful, but he had a stercoraceous odour; the abdomen was exceedingly tympanitic, but not tender; the pulse was quick, small, and sharp; the skin was natural; the tongue was moist and clean. The urine was at first scanty, but afterwards became more abundant, and of a deep colour.

On inspection the colon was found enormously distended, the small intestine rather less so; and both contained fluid feces and gas. There were numerous eroding ulcers in the small intestine. Two inches from the termination of the colon the intestine suddenly contracted; the mucous membrane was much thickened; and an ulcer was found with broad, elevated edges of a leaden hue. The canal would only allow a goose-quill to pass. The lymphatic glands near the stricture were enlarged, and the peritoneum was of a dark colour, and presented some adhesions. Other viscera were healthy.

CASE CCVI.—*Cancerous Disease of the Sigmoid Flexure. Constipation simulating Hernia.* (From the Museum Records, No. 1853.)—Henry J—, æt. 64, was a corpulent man, who had been intemperate in his habits. On admission the bowels had been constipated for four days. For twenty years he had been subject to hernia, but it had never become strangulated, and two years before it had been returned without any unpleasant symptom. On admission, the abdomen was greatly distended, but no proof of hernia existed; he had once previously suffered from constipation, which had been relieved by fomentations. Injections, &c., were used without any relief, vomiting of coffee-ground substance came on, hiccup and death followed on the eighth day. The cæcum and colon were found to be enormously distended; in the middle of the ascending colon was some contraction, but three or four inches from the rectum was a complete and firm constriction; there was considerable peritoneal effusion, and some old adhesions had led to the partial obstruction of the ascending colon; the mouth of the hernial sac was closed; there was intense congestion of the cæcum, and submucous purulent infiltration; at the constriction all the coats were diseased; the stricture would only admit a goose-quill; the mucous membrane terminated in a red vascular edge; the mesenteric glands were not generally affected, but one of them contained a calcareous mass.

In this case, if an operation had been performed, the constriction of the ascending colon might still have led to obstruction. It presents one of the difficulties which are always hidden in the diagnosis of these affections.

CASE CCVII.—*Cancer of the Sigmoid Flexure. Obstruction. Relieved. Gradual Exhaustion.*—Richard C—, æt. 32, was admitted under Dr. Gull's

care, July 2nd, 1854, and died September 3rd. He had been troubled with symptoms of obstruction for five months, his abdomen often becoming distended, and again diminishing after escape of flatus. Various remedies were given, and with considerable success (quinine and opium, &c.) The bowels became freely acted upon, but the patient gradually wasted, and at last sank.

Inspection, twenty hours after death. The heart and lungs were healthy. The *abdomen* was enormously distended on account of the size of the large intestine; the omentum was drawn upwards. The small intestine was much enlarged; but the cæcum and colon were enormously so. Just within the hollow of the sacrum was the constriction, which could be felt as a hard lump about the size of a hen's egg. The disease occupied four inches of the canal, and consisted of epithelial cancer. The walls were much thickened, and in the cellular tissue around was hard tissue, of a scirrhus character. The interior of the gut was ulcerated, and upon it were a few vascular fringes. The mesentery contained a few hardened glands. The walls of the intestine were considerably hypertrophied. The remaining tissue was healthy.

This case was an exceedingly interesting one, showing the beneficial and marked effect produced by judicious treatment. On admission there appeared but little probability that the obstruction would be overcome; the opium which was administered with quinine, so far allayed the intestinal action, and spasmodic contraction, that fæces slowly passed the stricture, and for a time there appeared probability of recovery.

CASE CCVIII.—Colloid Cancer of the Sigmoid Flexure. Artificial Anus in the Groin. Pleuro-pneumonia.—Thomas C—, æt. 56, had had severe pain in the course of the ureter, and it was supposed that he had renal calculus. On admission it was evident that there was an abscess forming in the iliac region; this reached slowly below Poupart's ligament, and was allowed to open itself. The patient became more and more prostrate, and a few days before death troublesome diarrhœa came on.

Inspection, was made seven hours after death.—The body was rigid, and much emaciated; on the left side below Poupart's ligament, and at the crest of the ilium were two openings about a quarter of an inch in diameter, the surrounding skin being thin and red; a probe passed for several inches along the course of the crest of the ilium, and a discharge of feculent pus proceeded from the wound. The head was not examined; arcus senilis was well marked.

Chest.—There were old pleuritic adhesions on the right side, and at the base of the left pleura. The lower lobe of the left lung was consolidated, of a mottled red and grey colour, very soft, and readily breaking down, and a considerable quantity of serum was contained in it; the anterior and upper

lobes of both lungs were emphysematous in a marked degree. The bronchial glands were healthy.

Heart.—There was a white patch on the surface of the pericardium, and its cavity contained about 3ijj of slightly turbid serum. The cavities of the heart were distended with clot. The left ventricle was slightly hypertrophied. The aortic and mitral valves were atheromatous, and the endocardium opaque. The coronaries were atheromatous at their commencement, and there was commencing fatty degeneration of the heart. The ascending aorta presented rigid atheromatous plates; the weight of the heart was seven ounces.

Abdomen.—The parietes were rigid; the intestines were collapsed; two bands of omentum were adherent at the sigmoid flexure. The stomach was low down, and much distended; its mucous membrane was mammillated; the secreting cells were granular; the pylorus was healthy. The mucous membrane of the cæcum and colon were of a grey colour. The colon was contracted; at the commencement of the sigmoid flexure was a hard mass resembling scybala; on opening this it was found to consist of a soft cancerous growth. The calibre of the intestine was almost obliterated by this irregular growth from the mucous membrane, it involved the whole circumference of the gut, and would only admit the little finger at the upper margin. This growth was rounded, foliated, and extended in one part an inch up the descending colon; the lower margin was of the same kind, but more intensely congested. The breadth of this diseased portion was from one to three inches; the intermediate part was ulcerated, and a communication passed at the posterior part into an irregular sinus, behind the fascia covering the quadratus lumborum; this sinus was filled with feculent pus, and burrowed downwards along the crest of the ilium to the openings in the skin. On making a section of the growth, it was found to be soft, of a yellowish white colour, and had a striated appearance, and juice could be compressed from it; several parts presented transparent gelatinous masses of colloid cancer. The whole of the mucous and muscular coats were involved and destroyed; and the muscular tissue of the quadratus lumborum was filled with round isolated masses of colloid growth, separated by bands of muscular fibre. The surface of the growth presented under 350 diams. (Plate III, fig. 4) columnar epithelium, some cells of normal size, others much enlarged (*a* *b*) and containing single or double nuclei; some of these cells were oblong; the principal portion, however, of the growth was composed of large nuclei, about 1000th to 1500th of an inch in diameter, with distinct nucleoli, and closely packed together with very little intervening blastema (*c*); there were some large cells containing several nuclei (*d*). On the field were numerous masses resembling inflammatory granule cells (*e*). The intervening tissue consisted of delicate fibres, arranged so as to form cells (*e*); and in some parts presenting elongated cells (*f*). There was no doubt as to its cancerous character; and there were a few small infiltrated glands in the neighbourhood of the cancerous growth. On the surface of the liver, both on the right and left lobes, the peritoneum was thickened from attrition; the structure of the liver was rather fatty and coarse, its weight was 3½lbs. The vena portæ and mesenteric veins were much congested. The spleen

was soft, its corpuscles were visible; it was 7oz. in weight; the supra-renal capsules were firm, large, and apparently healthy. The kidneys were somewhat atrophied, and contained a few cysts; they were 8½oz. in weight; the genital organs were healthy.

In this case, the examination of the fæces of the discharge might have detected cancer, but no tumour could be felt; there was no marked constipation, but pain in the course of the ureter was the principal symptom.

CASE CCIX.—Cancerous Disease of the Sigmoid Flexure. Diarrhœa. Perforation. Fæcal Abscess.—Elizabeth S—, æt. 55, admitted into Guy's Hospital, March 29th, 1854, was a married woman, but she had had no children. She was much emaciated, and for three years had ceased to menstruate. On admission she had a hot and dry skin; the abdomen was tender; the pulse was sharp and frequent. She had had pain in the hypogastric region, with vomiting and purging, and the stools had contained blood. The diarrhœa became more severe, and there was increased tenderness and pain at the lower part of the abdomen, the evacuations contained inflammatory product. She died on May 16th, severe purging having continued.

On inspection the lungs and heart were found healthy. A cancerous growth was situated above the sigmoid flexure; and there was ulceration of the new growth. The calibre of the intestine was contracted, and there was thickening of the mucous and muscular coats of the descending colon. The omentum was adherent to the large intestine at that part, where a large fæcal abscess had formed, from the giving way of the descending colon above the seat of stricture. The liver was small and fatty. The kidneys were small and atrophied.

This case is one of much interest, as showing an occasional mode of fatal termination of cancerous disease of the intestine; and that after ulceration has taken place at the seat of stricture diarrhœa may come on. Here, however, the intestine had also given way, and had led to peritonitis, and the formation of fæcal abscess.

In the 'Guy's Reports,' 1850, Mr. Birkett has recorded a case of insuperable constipation, arising from stricture at the upper third of the rectum, and associated with scrotal hernia. The patient was forty-nine years of age, and for fourteen years he had had hernia. The bowels had been rather constipated. On June 13th, he could not reduce the hernia, and applied at

left side, where the tumour which had been felt during life in front of the kidney was observed. There were adhesions firmly uniting several coils of intestine together. On separating them, which could be done without tearing the intestine, a feculent cavity was found, bounded above by the transverse colon, where it joins the descending colon, and by the greater curvature of the stomach; behind, by the pancreas; below, by several coils of jejunum.

The transverse colon presented an irregular opening about three inches in circumference, the edges of the opening were thickened, stained by adherent fæces, infiltrated with cancerous product, and in some parts were half an inch in thickness. The pancreas at its lesser extremity, and some of the adjoining glands were infiltrated with cancer; the stomach, though adherent, was not affected. At the lower part of the abscess two coils of the jejunum were firmly adherent, and were perforated; one, by a transverse opening extending about half across the intestine, the edges of which were everted and much injected; the other, by a smaller opening. The mucous membrane of the jejunum generally was injected, and covered with much mucus. The stomach and remaining parts of the intestine were healthy, so also were the liver and kidneys. The uterus, ovaries, and glands were healthy.

The disease was here of a strictly local character. The examination of the growth showed that it consisted of nuclei resembling those found in cancerous disease, and the general appearance was very strikingly that of cancer; still no other part was affected. The blow which she had received at this part set up inflammatory disease, and it is probable that a cancerous action subsequently ensued; ulceration then took place, and a fecal abscess formed.

The diagnosis was difficult; the position of the tumour was that usually found in disease of the glands about the kidney, but no abnormal condition of the urine existed. The vomiting was less persistent, and the diarrhoea more severe than is usually observed in cancerous disease of the stomach; but although the colon was thus extensively diseased, constipation did not occur.

CASE CCXI.—*Carcinoma of the Rectum, of the Ovaries, and of the Peritoneum. Acute Peritonitis. Scirrhus.*—Ann S—, æt. 26, admitted March 26th, was a married woman, living at Dockhead, and her youngest child was two and a half years old. For one year she had had difficulty in the passage of the alvine discharges. She was exceedingly ill on admission, and no connected history could be obtained; the lowest part of the rectum was sacculated, and about two inches upwards a stricture was found, through which a catheter could be passed. She suffered considerable pain, but no vomiting; she gradually sank, and died April 13th.

Inspection seventeen hours after death.—The body was very much emaciated; the head was not examined. At the apices of the lungs were slight pneumonic consolidation, and a little chalky deposit. The heart was small, and without fat; a tolerably firm clot was found on the right side, and in the right pulmonary artery.

Abdomen.—The intestines were distended. The peritoneum was intensely injected, and the coils of the small intestine were matted together. The mesentery was shortened. The great omentum was contracted into a firm mass, and was nodulated; nearly the whole of the peritoneum was minutely studded with small white tubercles; these were very numerous upon the peritoneal surface of the stomach. The sigmoid flexure and the upper part of the rectum were very much distended.

On taking out the large intestine, a growth was found about three inches from the anus, having a semi-cartilaginous hardness. On its inferior surface the infiltrated mucous membrane had a double lip-like appearance, and was considerably raised. In the centre of the growth, all the coats of the intestine were destroyed, and were infiltrated with heterologous deposit. The mucous membrane had a yellowish-white appearance on section; beneath it was firmer, white fibrous product, mixed with iron-gray pigment; still lower, fat with firm tissue. The whole of the external cellular membrane was semi-cartilaginous. On careful microscopical examination, the mucous membrane was found to consist of a delicate cellular tissue of nuclear fibre, interlacing and leaving spaces filled by elongated and reniform nuclei; a few cells were observed, but the cell wall was very imperfect; the sub-mucous tissue was very beautifully composed of series of bands of fibre tissue, with intervening columns of nuclei; at the upper part these bands of fibres formed series of arches. The muscular coat of the intestine above the stricture was much hypertrophied. In the sigmoid flexure above the stricture were one or two superficial ulcers or abrasions. The descending colon was filled with solid bilious fæces, but was otherwise healthy. The cæcum and small intestine were also healthy, as to their mucous membrane. The whole of the cellular tissue about the ovaries was thickened, white, and infiltrated; both ovaries also were infiltrated with cancer, and one mass was of a yellowish colour, as if degenerating. The uterus, vagina, and bladder were healthy. The liver was fatty. The stomach and spleen were healthy. There was no infiltration of the lumbar nor of the mesenteric glands. The kidneys and supra-renal capsules were healthy.

The disease in this case began apparently in the rectum, and extended from it, by continuity of structure. It was of a scirrhus character rather than epithelial, and although the obstruction was so great as only to allow a goosequill to pass, no vomiting was produced by the constipation; the reverse would have been the case if violent drastics had been administered. The character of the pain in this instance was more severe than we find in disease of the sigmoid flexure; there was

direct pressure on the nerves of sensation, and the disease extended to the adjoining structures. The growth could be felt on rectal examination, so that there was no difficulty in the diagnosis.

CASE CCXII.—*Epithelioma of Rectum. Contraction and Obstruction. Artificial Anus in the Loins. Diseased Appendix Cæci. Fæcal Abscess.*
—Mary P—, æt. 48, was admitted into Guy's Hospital October 7th, 1859. She had suffered from constipation for three weeks, accompanied with vomiting, and great abdominal distension. She was a thin person, having an aged, haggard expression; and when brought to Guy's, she was in such a condition that life was despaired of for many hours. The abdomen was much distended, but free from pain, and there was resonance and distension in the loins. On October 8th, Mr. Bryant made an incision into the left loin, and opened the descending colon; thin fæces were abundantly discharged; the operation was performed without unusual difficulty, but the patient gradually sank, and died on October 19th, at midnight.

Inspection 20th, fifteen hours after death.—The body was very thin, and the abdomen collapsed; the peritoneum in some parts had lost its shining surface, but there was no evidence of general peritonitis. The peritoneum covering the iliac fascia on the left side was thin and green, and nearly perforated. The rectum passed directly up to the cæcum, then turned to the left along the brim of the pelvis; at the angle was a small fæcal abscess, bounded by the rectum, cæcum, and by a portion of small intestine; it contained the appendix, which was obliquely truncated near the cæcum by ulceration; there was a contracted portion of rectum about the centre of the concavity of the sacrum, having the appearance externally as if girt with a portion of string. On opening the bowel it was found to be nearly occluded; it was ulcerated on its inferior aspect; and above, it had a raised slightly vascular fringe; there was no enlargement of glands. The opening into the descending colon was well situated, it was about one inch in length, the mucous membrane had united to the muscle immediately beneath; but beyond that, near the skin, the tissues were in a state of slough; and fæces had burrowed down under the fascia, so as to occupy the whole of the iliac fossa; this fæcal abscess nearly perforated the peritoneum, and extended into the labium on the left side, where was a small opening. The bladder was distended. The liver was pressed down; a white thickened patch existed on its surface.

This patient was too prostrate to allow of reparative changes after the operation had been performed, so that fæcal extravasation took place in the loins; the disease in the rectum was of a local kind, and the operation was skilfully performed; still, not only was surgical help deferred too long, for when first admitted she was almost dying, but the disease of the appendix cæci, and

a fæcal abscess connected with it, would in themselves have led to a fatal termination.

CASE CCXIII.—*Cancer of Jejunum, and of the Mesenteric Glands. Cancerous Softening of the Spinal Cord. Paraplegia.*—Samuel S—, æt. 15, was admitted into Guy's Hospital December 17th, and died on the 31st, under Dr. Rees' care. After a fatiguing march, in a rifle corps, five weeks previous to the commencement of his illness, he began to suffer from "pins and needles" in his legs, followed by weakness, and complete paraplegia. The respiratory muscles became involved before death. A tumour was felt in the lower part of the abdomen, near the anterior and superior spinous process of the ilium on the right side; but neither history of abdominal pain, nor any distressing abdominal symptom was made out during life. The bowels acted without purgative medicine; the motions, however, were discharged involuntarily.

On inspection, the spinal membranes were found to be clear, but rather more adherent than usual; the vessels also appeared very full of blood; and at the upper part of the dorsal region the cord had a diffused ecchymosed appearance, several points of extravasated blood were also found. In the lumbar, and lower part of the dorsal region, the cord was hard, but at the upper part it became soft, and in some parts semi-diffuent; one or two spots were more prominent than others in this softened portion.

Abdomen.—The peritoneum was healthy; the omentum was spread down to the pelvis, and was adherent near the right iliac region to a hard mass, about three inches in circumference, and which consisted of an enlarged and infiltrated mesenteric gland; an enlarged coil of jejunum was adherent at the part to the omentum. The cæcum was free and empty, and was situated behind and a little to the right of the growth just mentioned. On removing the intestine and opening the distended portion opposite to its mesenteric attachment, it was found to consist of about six inches of jejunum dilated into a large sac; at the entrance into this sac the mucous membrane was infiltrated with cancerous product, so that the valvulæ conniventes were hard and prominent; a similar state also existed at the outlet of this sac; in some portions of this distended bowel the coats were very much thinned, as if about to perforate into the peritoneum. The mucous membrane of the jejunum, however, was entire. A mesenteric gland in the centre of this coil of jejunum was infiltrated with medullary cancer; it was very soft and pale in colour; other glands were infiltrated in a less degree; and one or two near the pancreas were also enlarged. The growth in the jejunum consisted of yellow cancerous matter infiltrating the coats of the intestine. In another portion of jejunum there was a white, hard mass, opposite to the mesenteric attachment, about one inch in length, and one eighth of an inch in thickness, consisting of cancerous infiltration into the submucous cellular tissue, but the mucous membrane was entire. The colon, stomach, duodenum, and pylorus, were healthy. The liver and spleen were fissured, but healthy; there were several cancerous tubercles in the kidney, and the epididymis was enlarged. The lumbar glands were healthy. The left lung was solidified at its base.

The spinal mischief led to a fatal termination, but the case is remarkable not only in the form and situation of the cancerous disease, but in the comparative absence of symptoms of abdominal mischief. No obstruction took place in the intestinal tract, and the peritoneum was not involved. If, however, life had been prolonged for a short time, the more extensive affection of the mesenteric glands and the implication of the peritoneum would soon have led to well-marked indications of abdominal disease.

CASE CCXIV.—*Cancerous Ulcer of the Colon; opening into the Duodenum. Diarrhœa. Vomiting.*—Ann S—, æt. 47, by trade a greengrocer, was a woman who had worked hard, and drank freely; her father died from phthisis, but with the exception of an attack of acute rheumatism, and of erysipelas ten years before her last illness, she had enjoyed good health. Five and a half years before admission she had free access to a fruit garden, and partook of fruit to excess, severe diarrhœa and depression followed. Eighteen months afterwards the skin became slightly jaundiced, and her medical attendant found a tumour about the size of a hen's egg, immediately above the superior spinous process of the left ilium. She had also suffered from hæmorrhoids, and from tapeworm. When she applied for admission at Guy's Hospital, November 15th, 1859, she was pale and had a careworn appearance; there was brown discoloration of the abdomen, neck, thighs, and elbows, the gums were pale, and the tongue was clean. The resonance of the chest was good, and the respiration was healthy. The pulse was 90, and compressible. The abdomen was supple and resonant, excepting in the region of the cæcum, where a rounded, uneven, and hard tumour could be felt, about the size of a turkey's egg. There was slight dulness at that part, but no pain except when pressure was made, or when a deep inspiration was taken, or after swallowing fluids; pain extended also in the course of the ilio-hypogastric nerve, reaching as far as the trochanter, and also back to the spine; on pressure, also, the pain was in the inguinal region of the opposite side. The bowels were relaxed, and mucus was passed in the motions; the urine was of light colour, and of sp. gr. 1015, and free from albumen and sugar. There was good appetite; nausea was present, but no vomiting. Eight leeches were applied, and a poultice; the spermaceti mixture given, and Dover's powder with grey powder every night. The pain was relieved by the leeches, but on the 24th it again became very severe, and the bowels were relaxed. The leeches were repeated, and calomel and opium of each gr. j given every night. 25th. The pain was less, but there was tenesmus and diarrhœa; an enema of starch with tincture of opium was administered. On the 30th, leeches were repeated to relieve the pain, and opium was given.

On December 4th, there was diarrhœa, and the swelling had increased, extending upwards; vomiting came on. Logwood mixture (G.P.) was given. On the 7th, sickness was relieved by brandy and soda-water. The symptoms partially subsided, and she left the hospital on January 31st.

She was readmitted on March 13th, 1860. Fifteen days before, at 3 p.m., she had an inclination to go to stool, but before she could reach the closet she felt something give way in the abdomen, and a profuse discharge of very fetid pus took place from the bowels. The tumour diminished in size; the bowels continued relaxed, and for one week there was blood in the evacuations; she seemed low and weak; there was no appetite, but considerable thirst; there was pain in the tumour before and after the discharge, emaciation was manifest, the mouth became sore and aphthous, and the urine contained lithic acid. When admitted, she was emaciated, and her countenance was anxious and distressed; there was a deep flush on the cheeks; the eyes were sunken and hollow, the tongue was morbidly red, glazed, and cracked, and there was aphthous ulceration at the tip; the skin was hot and dry. The abdomen was soft. There was undue prominence over the right ilium; and pain with tenderness was present at the epigastrium, as well as in the right hypochondriac and iliac regions; the tumour which had been felt so long was resonant on percussion. The hepatic dulness extended two inches below the sternum; the appetite was bad; and there were thirst and nausea; the bowels were moved ten to twelve times, and the motions consisted of very fetid brown fluid, and contained scarcely any solid matter, and no blood. The urine was scanty. The compound decoction of *krameria* was given every six hours, with brandy, milk, arrowroot, &c.

March 25th. For two days obstinate vomiting tried the patient; it increased in severity; and came on after any exertion, and after taking food; rapid prostration followed, and death.

On *inspection*, the thoracic viscera were found to be healthy. In the abdomen, the ascending colon was firmly fixed in the right loin; and it was adherent to the surface of the liver; but the liver and kidneys could be removed without interfering with the diseased part. On opening the colon there was found in the ascending part, just above the cæcum, a large carcinomatous ulcer, as large as the palm of the hand; it was circumscribed and surrounded by raised edges of morbid structure. In some parts of the diseased tissue the edges were undermined, so that bridges of tissue passed across. At the bottom of the ulcer was a large hole, through which the middle finger could easily be passed, and entered at once into the duodenum. On opening the duodenum the perforation was seen within it; the hole was as large as that in the colon, but its edges were merely fringed by the new growth. The duodenal opening was near the pylorus, and opposite to the common bile duct, so that the bile might have at once passed into the colon. The contents of the intestine, both small and large, were of a pale slate colour, showing the absence of the bile. The liver was very fatty, and it was lighter than water. The kidneys were healthy. The omentum was adherent to the right ovary, and the Fallopian tube was adherent to the ovary.

In this case there was evidence of chronic disease of the cæcum or of the colon, as shown by the gradually increasing tumour; the pain increasing at once on drinking fluids was very manifest. The enteric irritation was propagated to other

parts of the intestine, and diarrhœa was produced; this for a long time was, with the pain, a very troublesome symptom. About a month before her death, at the time that something was felt to have given way in the abdomen, the opening into the duodenum was probably suddenly made, and from that time vomiting was a more constant symptom; there was no evidence, however, by stercoraceous vomiting, that fecal matter passed from the colon into the duodenum. As to the cause of the disease of the colon, it is possible that the primary irritation of the intestine after partaking of a large quantity of fruit, might have determined the seat of the disease; this, however, is very doubtful. From the first, the treatment was in a great measure palliative rather than strictly remedial; but much may be done in these cases to diminish the sufferings of the patient and to prevent the rapid extension of the disease.

CASE CCXV.—*Ulceration of the Colon. Intestinal Obstruction from Contraction of the Transverse Colon. Cancer? Dysentery. Constipation. Diarrhœa.* (For the following case I am indebted to my friend Dr. Wilks).—David B—, æt. 72, was a rather spare man, of middle stature, and with a yellowish complexion, but he had never been abroad; his habits had been rather intemperate. He stated that his general health had been tolerably good, until within the last few years, during which he had suffered at frequent intervals from diarrhœa, with colicky pains in the abdomen, and often from painful defecation. His bowels were at times so irritable that, after swallowing only a cup of tea, it seemed at once to pass through them, almost before he could reach the closet; these symptoms became increasingly severe. Nine years previously, also, he had received a blow in the left groin, which produced a femoral hernia; it was reducible, and, with a truss, did not give him much inconvenience.

On August 11th, 1855, he complained of diarrhœa, stating that his bowels were relaxed, several times during the day; and that he suffered at times from severe pain in the abdomen; the pulse was full, 75; the tongue was clean; the appetite was good; and, excepting the symptoms just mentioned, he seemed to be in good health. He was ordered chalk mixture with aromatic confection three times a day.

On the 14th, the bowels were less relaxed, but the pain in the abdomen was greatly increased; and sulphuric ether, with tincture of opium, was given in pimento water every four hours.

On the 16th, the abdominal pain had greatly increased, it was nearly constant, but at intervals became more severe; there was no tenderness on pressure at any particular part, but the greatest amount of pain was felt about the umbilical region; the bowels had not been open for two days. A blister was ordered to be applied to the epigastrium; a powder consisting of

gr. viij of jalap, and calomel gr. ij to be taken ; and every night a grain of calomel and of opium.

On the 17th, the bowels were not open, and the pain was unrelieved; the jalap powder was repeated.

On the 18th, unable to leave his home; the pain was so much worse that he could not rest; and he was continually moving about moaning in great distress; the bowels had not been relieved by the powder, and the stomach had become irritable; the pulse was moderate and full; the tongue was clean and moist; the skin cool and natural. A senna and sulphate of magnesia draught was ordered to be taken every four hours; an enema to be administered, and a poppy fomentation applied to the abdomen.

On the 19th, he was very much worse, and had had no rest for two nights; the pain was very severe, coming on in paroxysms, at intervals of ten or fifteen minutes, and at those periods the abdomen became very tense and tympanitic; the bowels were not open; and he vomited a large quantity of dark grumous matter twice; his appetite was gone, and the stomach had become so irritable that he at once rejected everything taken; his hernia was down, but was easily reduced; there were no febrile symptoms. He was ordered the compound soap pill gr. v every six hours, a hot fomentation to the abdomen, and an enema of castor oil, and oil of turpentine.

20th. The bowels had not been open; the pain and vomiting still continued, but the paroxysms were not so frequent nor so severe, and he had slept a little. To continue the medicine.

21st. A few scybala were passed; but the pain, the vomiting, and the constipation persisted; to have an enema every six hours, and to take sulphate of magnesia \mathfrak{z} ij every two hours.

22nd. He was not so well; he had had no rest during the night; the pain had increased, and all the symptoms were more urgent, but the pulse was still natural, although the tongue was becoming rather dry. To repeat the enema every six hours, and to take colocynth pill gr. v, and croton oil $\mathfrak{m}\mathfrak{z}$ every four hours. At 11 p.m. the symptoms were more severe; the paroxysms of pain were more frequent and violent; no action had taken place from the bowels; and he had not been able to take any nourishment for several days, except a little beef tea. Ordered calomel gr. j, opium gr. $\frac{1}{2}$, every three hours, and to take every two hours an effervescing draught of soda, ammonia, and hydrocyanic acid with tartaric acid.

23rd. He was much the same; the bowels were not open; stercoraceous vomiting had come on, and the pain was equally violent and almost continuous. The skin cool and moist; the tongue was dry in the centre, and slightly furred; pulse, 70, smaller; ordered to take croton oil $\mathfrak{m}\mathfrak{j}$, and every six hours the solution of morphia.

24th. The compound soap pill was given every six hours, and the mixture was continued.

25th. On the evening of this day copious evacuation took place from the bowels; he continued the pills.

On the 26th and 27th, the diarrhœa continued; and persisted till his death, on the 29th.

Inspection, August 30th.—The body was spare, but not much wasted. The abdomen only was examined. There was recent acute peritonitis. The intestines were distended, injected, and covered with a slight exudation of lymph. The seat of stricture was at once seen to be the middle portion of the transverse colon, exactly in the median line of the body; the omentum was found abnormally adherent to this portion of the large intestine, and there was situated there a dark-coloured, hard-looking substance; this formed the constricted portion of the intestine, and both above and below it there was considerable dilatation. On removing this portion of the intestine, the disease was felt as a hard tumour situate in the substance of the organ, and producing the constriction. Although the exterior continuity of the bowel exhibited a considerable falling in at this part, its lessened calibre was not so manifest from the exterior as from the interior. When the intestine was opened, its channel was found to be so reduced in size that it would only admit a goose quill. On laying the whole of it open, the diminished passage was found to be, not only owing to the external peritoneal puckering, but to the hypertrophy of the subserous cellular tissue, and of the muscular coat, and to a raised spongy condition of the mucous membrane. When spread out, the diseased surface occupied a space not much larger than that of a five-shilling piece. The mucous membrane was red, highly vascular, and completely separated from the healthy surrounding structures, not only by its colour, but by its highly raised margin, which was considerably above the level of the adjacent mucous membrane. It had a soft, spongy, velvety appearance, and the muscular coat beneath was much hypertrophied. The microscope showed the surface of the structure to be composed of highly vascular branching villi, and therefore approaching in character the so-called villous cancer. The surface was covered over with an abundance of columnar epithelium. More deeply seated was found a delicate fibre tissue, with a number of small nucleated cells, of a shape resembling ordinary or abortive epithelium. The cæcum and ascending colon, as well as the lower part of the descending colon and the rectum, contained numerous ulcers in various stages of healing; most of them were quite healed, and presented only cicatrices. There were large irregular shaped portions of mucous membrane, of a dark blue or slate colour, which seemed to have been renewed, but causing a considerable puckering of the surface. In the cæcum and rectum the general calibre of the intestine was much altered in shape by the contraction, and also by the hypertrophy of the muscular coat, which was very considerable in the cæcum. The liver contained no cancerous disease, nor did any of the abdominal glands.

Simple acute disease of a dysenteric character took place in this patient, ulceration followed, and, at the seat of one of these ulcers a villous growth was developed, which led subsequently to constriction; the coats of the intestine at that part were hypertrophied, showing that the obstruction had been existent for some time. As to the nature of the growth, although no

strictly cancerous product was found, and although no glandular disease was present, its villous character and cellular substratum showed that it belonged to the class of growths designated as cancerous. The first symptoms were those of dysentery, as shown by the diarrhœa and the discharge of mucus; but as the disease extended to the deeper structures, and spasmodic contraction took place at the seat of the villous growth, the obstruction became complete, till vomiting even of a stercoraceous kind was set up; and it was only as the strength of the patient failed that the constriction yielded and the bowels acted; diarrhœa then came on, and continued till death. In the treatment of the patient, the increase of the symptoms after purgative medicines, and their relief after the administration of opium, were well marked.

These instances show, that with care the several forms of internal strangulation may be generally distinguished, when we have the whole of the symptoms before us; that whilst over-active and injudicious treatment increases discomfort and hastens a fatal termination, much may be done for the relief of the patient, and valuable lives thereby be prolonged. These are not the cases for do-nothing practice; the proper use of enemata, of such diet only as can be borne without injury, opium, rest, and other means to which we have referred, will mitigate suffering even where cure is impossible.

CHAPTER XV.

ON INTESTINAL WORMS.

THE presence of some species of entozoa in the alimentary canal must be considered as an indication of disordered secretion from the mucous membrane; for in most cases it is only because the germs of these entozoa have found a nidus, in which their development can take place, that human beings become subject to them; this can more especially be said of those which make their habitat on the mucous membrane of the rectum, as the minute thread-worms. As to the *strongylus*, the *trichina spiralis*, hydatids in various structures, and *filaria*, their entrance into the system arises more from locality, and sometimes from causes unconnected with the condition of the patient. The germs of the *tænia* appear almost indestructible by ordinary means.*

The forms of entozoa which infest the alimentary canal are the *tænia solium* and the *tænia* or *bothricephalus latus*, the *trichocephalus dispar*, the *ascaris vermicularis* and *lumbricoides*.

The *ascaris lumbricoides* somewhat resembles the earth-worm in external appearance; it is found in the small intestine, and sometimes clusters of them, two or three, may exist together in different portions of the whole canal, in the duodenum, jejunum, ileum, and stomach; I have seen them in the œsophagus, and in some cases have been described where they have produced fatal results by getting into the larynx; they are found also in the colon. It has been stated that they are able to destroy the coats of the intestine and thus reach the peritoneal cavity; but it is not generally believed that ulceration and perforation had taken

* 'Medico-Chir. Review,' Jan., 1857, on "Entozoa in the Human Subject."

place, and the lumbricus had used the opening thus made; the irritation they produce may, however, aggravate ulceration in a marked degree; still, some instances appear to confirm the idea that lumbrici may themselves induce ulceration and consequent perforation; such, for instance, is one recorded in the 'British Medical Journal' of 1861, by Dr. Sandwith, in which a small perforation in the middle of the duodenum led to fatal perforation. Dr. Young, in a paper in the 'Medical Gazette,' records several cases where lumbrici were evacuated through the abdominal parietes; one instance was in a child, aged seven; several worms had been discharged from the bowels, an abscess afterwards formed in the right lumbar region, and living lumbrici were evacuated; after two years the abscess healed.

In another case, a child aged fifteen had severe pain in the abdomen, and the lumbrici were found in the stools; an abscess formed on the right side of the abdomen, and lumbrici were passed through it: on inspection it was found that a faecal abscess had been formed at the commencement of the colon, into which also the jejunum opened. It is probable that in both these cases either caecal or strumous disease or direct injury had led to abscess, through which some of the lumbrici were discharged. The same paper mentions a recorded case of an infant, in whom a lumbricus was discharged from the navel; and another of a woman, in whom an artificial anus existed in the right groin, through which a lumbricus was passed. The symptoms found with this worm are a tumid, full, and doughy state of the abdomen, indicating that the secretions and muscular coats are not in their normal condition; the body is often feebly nourished; the cheerfulness is lost; the appetite is variable, sometimes craving, but uncertain; there is frequently severe colic; the breath is offensive, and there is irritation of the nose and anus. Other signs frequently are found, but they are in many instances sympathetic from the irritation of the intestine, especially when the subject is strumous, and predisposed to disease in other viscera; thus the brain often sympathises, so that the patient may be seized with convulsions, or have attacks of cholera, or epileptic fits, conditions which are entirely removed by the evacuation of the entozoa.

These worms occur more frequently in young children, but

and the female about twice as long; but they are numbered by scores.

Relief is best obtained by improving the general health of the child, by regulating its diet, by acting on the bowels by mild laxatives and alteratives, and by destroying the worm by local applications. Dr. Watson recommends the infusion of quassia, and I have often observed its efficiency; decoction of oak bark may also be used, or a solution of alum or of sulphate of iron, about gr. x to xx, in a pint of water; tincture of iron was used by Dr. Darwall in the proportion of 3ss to ʒviiij. Santonine has also been recommended.

Tricocephalus dispar.—I have seen this worm much less frequently than I expected from the descriptions of authors; having very often searched in vain for it in the post-mortem examinations made under my own superintendence at Guy's Hospital. It is about one to two inches in length, and the head is exceedingly small, resembling a hair, hence its name; it has a minute mouth, and at its larger extremity has an anal orifice; this latter extremity also presents a beautiful sheath, and in the centre of the sheath a spiculum, which is described as the penis of the male; the body is slightly curved or twisted. In the instances referred to, no sign indicated their presence during life, they were situated in or near to the cæcum; and it is in this part that observers have always found them.

Tenia solium, or *tapeworm*.—The presence of this entozoon in the intestine is an exceedingly common affection, and among the out-patients at Guy's Hospital many instances are presented. The paper of Dr. Gull, on the efficiency of the oil of male fern, in the 'Guy's Reports' of 1855, indicates the frequency of the complaint. Patients present themselves, saying that they have tapeworm, and that they have observed small portions, or several feet, with the fæces. On inquiring into the symptoms, we find the patient is generally pallid, and complains of irregular and sometimes voracious appetite; there is also pain, with sense of exhaustion at the stomach and abdomen, and often with nausea and general feeling of malaise; the tongue is clean or furred; there is some irritation of the nose and anus; the pulse is usually compressible, or it is sharp and excitable; the pain is, however, much less than with the *ascaris lumbricoides*;

dition. The *tænia solium* is that which we almost always observe in this country, but I have several times seen the *bothriocephalus*. In this the head is destitute of hooklets and suckers, but has a transverse fissure in the segments; the generative orifice is in the centre, not on the margin. The *bothr. latus* is found in Switzerland, Poland, Russia, whilst Holland and Germany have, like ourselves, the *tænia solium*.

As to the treatment of these cases, the remedies called anthelmintics have been most varied; many appear to act by destroying the worm, and then by expelling it. This is the case with the oil of turpentine given in doses of ʒss , and followed by castor oil, or other purge, if necessary; it constitutes a very effective remedy, although sometimes producing vertigo and sickness; but these symptoms are transient. Turpentine, however, is a nauseous remedy, and very trying to the patient, although almost anything will be willingly borne to be freed from such a companion as tapeworm. The turpentine was a few years ago followed by the bulky remedy, the Kousso, from Abyssinia; this was less effective, and has now, in the hands of my colleagues and myself, given place to the oil of male fern, an old remedy, but one deservedly approved; and I can bear testimony to its value. Drs. Hughes and Gull administered it in numerous cases, which are stated in the paper previously cited, and we now seldom find that it fails. In several instances I have found the patients return, after a few months, again complaining of the same disease, having passed segments; the male fern has been repeated, and with apparent success. In one instance the patient came a third time. It is true that in this case the head was not found, and without this we can never be sure that the disease is eradicated. The oil of male fern does not appear to produce any injurious effect, thus, a child, by mistake, took ʒiss . of it every night for a week; purging was the only uncomfortable symptom that followed.

It is generally given quite on an empty stomach, and I usually follow it by a dose of castor oil in about six hours. It is well not only to give the oil on an empty stomach, but to avoid taking food till it has acted, so that it is well to administer it at bed-time. I have no experience in the pomegranate root, which is stated to be of service; and none whatever in the almost ex-

ploded remedies of cowhage or metallic tin.* After the worm has been discharged, vegetable tonics, with mineral acids, and especially some of the preparations of steel, according to the age and condition of the patient, ought to be prescribed; as the steel wine, or the ammonio-tartrate, or the tincture of iron, with quassia, &c.

Dr. Gordon, in his 'Report of Diseases of the Stomach and Bowels in India,' describes a remedy for tapeworm, which he has found exceedingly effective, namely, the Kameela, or *Rottlera tinctoria*, in ʒj doses; Dr. Osborne recommends tannin; and the aromatic sulphuric acid has been used beneficially.

* 'Medical Times and Gazette,' May, 1857.

CHAPTER XVI.

PERFORATION OF INTESTINE FROM WITHOUT. ABSCESS IN ABDOMINAL PARIETES EXTENDING INTO THE INTESTINE. FÆCAL ABSCESS.

PERFORATION of the coats of the small intestine ranks in the order of frequency next to perforation of the stomach; the colon is, however, perforated more frequently than is generally supposed. These perforations of the intestinal tract divide themselves into two great classes: 1st. Those which arise from disease commencing in the intestine itself, and to which we have referred in numerous instances, as perforation of the ileum in fever and in struma; of the cæcum and its appendix; of the colon in dysentery, in cancerous disease, and in several forms of insuperable constipation. 2nd. Those in which the perforation is from without, or from the extension of disease from adjoining structures. These latter cases constitute an important and an exceedingly interesting class of diseases; and the following causes of external perforation may be enumerated:

1. From the peritoneum, as in strumous peritonitis.
2. From disease of the stomach, as ulceration and cancer extending into the transverse colon.
3. From hydatids and abscess of the liver, thus forming a means of escape into the small or large intestine.
4. From calculi in the gall-bladder, setting up ulceration of the duodenum or of the colon.
5. From abscess in the spleen.
6. From abscess in the kidney.
7. From abscess in the abdominal parietes and loins opening into the intestine.

8. From diseased ovary communicating with the cæcum, colon, or rectum.

9. From cancer of any of the abdominal organs extending into the intestine.

10. From extra-uterine fœtation.

11. From one portion of intestine opening into another, as the appendix into the rectum.

12. From blows, and external injury.

In many of these forms of disease last enumerated, various and characteristic symptoms precede the perforation of the peritoneum and of the intestine; thus, the signs of cancerous disease of the stomach arise some time before fæcal eructation or vomiting indicate extension into the colon. In hydatid disease of the liver a rounded tumour, of slow formation, is detected, having often a peculiar vibratory thrill, and without general disturbance, before the occurrence of local peritonitis takes place, and the hydatids are discharged either by the mouth or with the evacuations per rectum.

In gall-stone very severe pain in the region of the gall-bladder arises, with vomiting and often jaundice, before intense peritonitis, from rupture into the general cavity of the abdomen, and before obstruction by its impaction in the jejunum or ileum take place.

In abscess of the spleen the symptoms are more obscure, and constitute part of a general constitutional disturbance, till perhaps the discharge of pus by stool indicates that a communication has been formed with the transverse or descending colon.

In abscess of the kidney, and pyelitis, there is purulent urine; but when there is suppuration external to the tunic of the gland the symptoms are more obscure.

In ovarian and cancerous tumours tactile examination will detect them. Some of these forms of disease are more obscure than others, but when fæcal abscess is the result there is considerable uniformity in their character; severe local pain and tenderness come on, with hectic fever, and steadily increasing prostration; and when the abscess is not limited by adhesion, a rapidly fatal result occurs.

Suppuration in the *parietes* of the abdomen is frequently observed, and simulates deeply-seated mischief; and for a short

time considerable obscurity may attend it. The symptoms are generally of an acute character; considerable pain and febrile excitement precede inflammatory œdema of the skin, and while the effused products are bound down by firm fascial investments the symptoms closely resemble cæcal disease, and local peritonitis, &c.; in fact every part of the abdominal parietes presents us with disease on the surface, resembling deeper injury. In the hypochondriac regions suppuration connected with the costal cartilages and ribs simulates abscess of the liver, empyema, hydatids, diseased gall-bladder, or corresponding disease of the spleen; in the right and left iliac regions abscess in the parietes may be mistaken for affections of the cæcum and sigmoid flexure of the colon; in the lumbar regions, for renal and spinal disease; in the umbilical, for strumous and cancerous disease; and, lastly, in the hypogastric region, pelvic cellulitis, for ovarian and uterine disease.

Simple suppuration in the parietes generally tends to the surface, and the abscess is opened or it is discharged spontaneously, and in many cases recovery takes place, unless the disease be associated with pyæmia, or occur in cachectic subjects; sometimes the suppuration spreads extensively among the muscles; it extends also in depth, and gradually produces local peritonitis, or discharges itself into some of the viscera. Thus abscess about the kidney opens into the colon, that in the iliac regions into the sigmoid flexure, or cæcum.

The most fertile sources of these forms of parietal suppuration are blows and falls. I have observed them from blows, and pressure on the abdomen, from falls on the back, &c. In pyæmia and in cachectic subjects very trifling causes appear to be sufficient to lead to this disease.

Diagnosis.—The pain will generally be found to be very superficial; but in many instances, at an early stage, before any inflammatory œdema has been produced on the skin, and whilst the disease is confined beneath the fascia of the abdomen, there is much obscurity in the diagnosis; and after pus is discharged, it must be remembered that a fæcal *odour* does not necessarily imply communication with the intestine; for transfusion of gaseous contents of the intestine, when there is tolerably close

contact, may cause the contents of an abscess to have a faecal smell.

In reference to the *treatment*, this obscurity is of no great moment, for at that period local depletion, rest, warm cataplasms, are equally applicable to local peritonitis as to parietal inflammation. When suppuration has actually taken place, the sooner the pus is evacuated the less likely is it to burrow among the flat muscles and fascia of the abdomen; and even in abscesses, faecal or otherwise, extending secondarily to the parietes, unnecessary delay is sometimes made in discharging their contents. The rule is, I believe, a correct one, to open these abscesses very early.

CASE CCXVI.—Suppuration external to the Sigmoid Flexure of the Colon, opening on the Anterior Abdominal Parietes, and communicating with the Intestine.—Elizabeth R—, æt. 39, a widow, who had supported herself by dress-making, was admitted into Guy's Hospital, under my care, March, 1855. Till a fortnight before admission she had enjoyed good health, when she felt pain in the back, which extended to the shoulders and knees. The greatest pain, however, was in the course of the ilio-hypogastric nerve. These symptoms were accompanied with considerable febrile excitement.

Saline medicines with colchicum, &c., were prescribed. In a few days the pain, which had simulated rheumatism, ceased, and she appeared to gain strength under the use of decoction of bark with carbonate of soda.

On March 26th, three weeks after admission, she complained of pain in the left iliac fossa, and a firm tumour about the size of a hen's egg could be felt deeply in that part. There were no tenderness in the spine, numbness in the legs, nor other symptoms of disease of the spine. An examination per vaginam was made by my colleague, Dr. Oldham, but did not afford any evidence of disease of the ovary. The bowels were easily acted on by grey powder, followed by castor-oil, and by enemata; but this action did not affect the size of the tumour nor alleviate the symptoms. The urine was normal; and there was no indication of renal disease. The pain gradually increased in severity, but it was considerably relieved by the repeated application of leeches, by taking iodide of potassium, bichloride of mercury, and occasional doses of morphia. It was then believed that the malady consisted in disease of the sigmoid flexure of the colon, with local peritonitis.

May 10th. The pain had returned with much severity, and hectic came on. The tumour increased in size; it could be felt extending to the quadratus lumborum; and it also reached to the anterior abdominal parietes, which, at the left iliac fossa, were red, œdematous, and exceedingly tender.

19th. The bowels were acted upon three times freely, and a considerable quantity of purulent mucus was discharged. The examination of this dis-

charge could detect no cancer cells. The pain and the hectic continued; the patient became pale and exhausted; the left thigh and leg, and afterwards the right, became swollen and tender; and there was excessive pain in the course of the femoral veins. Nourishment and stimulants were administered as the patient could take them. Quinine and opium, or morphia were given.

On June 8th, the inflammatory œdema of the anterior abdominal parietes had increased. My colleague, Mr. Callaway, made an incision at this part; and more than a pint of exceedingly offensive pus was evacuated. Every means were used to sustain the patient; but the discharge continued abundant, and it had a feculent odour; at length her strength gave way. The tongue remained clean and moist; but her appetite ceased, so that she became quite unable to take food. There was no pain at the scrobiculus cordis, no vomiting, nor thirst; but emaciation with a sense of exhaustion. Bed sores formed on the sacrum; and a few days before her death, cough came on, which aggravated her distress. She gradually sank, and died June 24th.

Inspection was made twenty-four hours after death.—The body was blanched; and the lower extremities were œdematous; the pleuræ were healthy; but the posterior lobes of the lungs were in a state of red hepatization; the heart and its valves were healthy.

Abdomen.—The peritoneum was healthy, except in the left iliac region, where the omentum, and several coils of intestine, were adherent. In this region was an abscess, situated behind the peritoneum and fascia, and containing offensive, feculent pus; it extended to the anterior abdominal parietes in front, above to the diaphragm and kidney, and posteriorly nearly to the spine. Very careful examination could detect no disease of the ilium, nor vertebra, nor of the pelvic cellular tissue. The abscess communicated with the sigmoid flexure by three small openings, in close contact the one with the other; their edges were not thickened, but valvular. The small and large intestines were otherwise healthy; and the opening into the intestine was evidently secondary. The uterus, ovaries, and kidneys were normal. The stomach was of normal size; its mucous membrane was pale, and it had undergone degeneration. The liver was more than 5 lbs. in weight, and extremely fatty. The lower portions of the vena cava, and of the common iliac and external iliac veins, were filled with very firm, white, adherent fibrin; and the coats of the veins were much thickened.

The review of this case showed that the pain in the course of the ilio-hypogastric nerve arose from direct pressure upon that nerve by inflammatory effusion; that the tumour felt in the iliac fossa consisted of this effusion pushing forward the peritoneum and sigmoid flexure; that the subsequent symptoms arose from suppuration, and its extension in various directions; inwards into the colon, leading to some extravasation of fæces into the abscess and of pus into the alimentary canal; forwards,

so as to reach the anterior parietes, where it was opened upwards, to the diaphragm; and inwards, to the cava and iliac vessels, which became involved and obstructed by fibrinous material. That it did not arise from diseased bone was proved by careful examination; and it is probable that some accidental blow had led to this suppuration, with its fatal results; or, the irritation in the intestine had led to inflammation external to it, and subsequently to suppuration.

After the tumour had been felt, evidence of suppuration soon arose; and the discharge of purulent mucus showed that it had formed some connecting link with the intestine, or that there was ulceration of the coats of the intestine itself. Renal, ovarian, spinal, or parietal suppuration, or cancerous disease of the sigmoid flexure, might give rise to many of these symptoms. The absence of all indication of diseased kidney was shown by the condition of the urine. Disease of the spine was exceedingly doubtful, from the absence of tenderness and numbness and from the course of the suppuration. The position which the tumour assumed and vaginal examination showed that the ovary was not involved. The tumour appeared to arise from disease near to the sigmoid flexure, either commencing in the viscus and extending outwards, or beginning in the parietes and making its way into the intestine. It was in deciding to which of these might be the case that the principal difficulty consisted. The discharge of purulent mucus from the intestine, and the feculent character of the pus, indicated a connexion between the abscess and the intestine. Before death, I was led to believe that the disease commenced in the sigmoid flexure, and that the suppuration external to the intestine was secondary; the inspection after death showed that the reverse was the case. It was closely allied to cases of suppuration external to the rectum, but so deeply was the mischief situated that any exploratory incision would have been unjustifiable till there was more certain evidence of suppuration than was presented at the commencement of the disease.

CASE CCXVII.—*Abscess in the Loins. Feculent-smelling Discharge. Pleuropneumonia with Feculent-smelling Sputum. Recovery.*—T. H—, æt. 34, was admitted into the Clinical Ward, under my care, June, 1855. He was a man of steady and industrious habits. His health had been good till an attack

rheumatic fever two years before; and at Christmas, 1854, six months before admission, he had a very severe injury whilst at work, falling upon his head, and it was believed that the skull was fractured. He remained for some time in the hospital, under the care of Mr. Birkett, but left well, and continued so till three weeks before admission, when, in the middle of the night, he awoke with great difficulty of breathing, the respiration being accompanied with considerable distress and pain. These symptoms increased greatly in severity, and presented the signs of pleuro-pneumonia on the right side.

On admission he was exceedingly ill; his countenance was pale; his eyes were glistening; the lips and nostrils were contracted; the teeth were covered with sordes; the tongue was brown at the base and edges; and the skin was hot and clammy. In the chest there was found to be increased roundness of the right side at the base, with imperfect mobility, increased dulness on percussion, and loss of tactile vibration; and in front, below the nipple, there was a pleuritic rub. On the left side, the respiration was puerile; and at the apices, the expiratory murmur was prolonged and coarse. The position of the heart was normal; its sounds were healthy, but the precordial dulness was somewhat increased; the respiration was 26 per minute; the pulse 95, feeble and compressible; the urine was high-coloured, acid, sp. gr. 1025. His position in bed was one midway between the right side and the back, with his knees drawn up, and the head thrown forward.

The prostration and typhoid state increased till June 14th, when deep-seated fluctuation below the ribs on the right side could be detected beneath the lumbar fascia about the quadratus lumborum muscle. An exploring needle was passed, and afterwards a director, and the wound enlarged; about a pint of pus, having a strong faecal odour, was discharged, and the abscess continued to discharge freely. After the opening of the abscess the respiration became more free, and he coughed up a considerable quantity of frothy mucus, having the same odour as the pus. His strength was sustained by nourishing food and stimulants; by quinine and opium; diarrhoea was occasionally troublesome, and the offensive expectoration became exceedingly distressing.

On July 2nd, he had so much improved as to be carried out into the open air for half an hour. The offensive character of the breath and respiration gradually subsided; healthy respiration became audible nearly to the base of the lung, and he continued to gain flesh.

In October he returned to his work, and then appeared a stout hale man; but a fistulous opening remained up to that time, which occasionally discharged freely. The sinus healed in about six months, and he then appeared in sound health, May 1857.

In this case deep-seated suppuration took place near the quadratus lumborum muscle; acute pleuro-pneumonia on the right side, and the most severe constitutional symptoms followed. The pus, which was evacuated, and the mucus expectorated, were of a most offensive and feculent odour; but

character. She thought that relief would be obtained by an action of the bowels, but found that she could not pass anything, and that the sensation of forcing and pain continued. On the following day she got up, but fainted, and had a severe rigor; on the 3rd day, she noticed that her abdomen was enlarged, chiefly on the right side; this was the seat of all the pain, and became so tender that she could not bear the weight of the clothes upon the part. Her left side had been free from pain throughout. The bowels remained confined for four or five days after the commencement of her illness, and were only moved by a second dose of purgative medicine. The constipation continued till the time of admission. Miscarriage then took place.

On December 17th, she appeared anxious, and had a slightly yellow tinge of skin. There were a few bronchial râles in the chest. The heart's action was irregular and intermittent, but unattended with any bruit. Her abdomen was large, the superficial veins being distended. On the right side above the ilium there was an elastic tender swelling, with apparent fluctuation in it; in front it was ill defined, but above was separated from the liver by a resonant space. There were no symptoms to connect it with the kidney, and it did not extend into the right groin. The tongue was furred and white. The bowels were purged by medicine. The urine contained no albumen, but at times the colouring matter of bile was in abundance. The pulse was irregular, and generally intermitted every fifth beat.

December 30th. The tumour had been gradually disappearing, and for the last two days could not be detected; there was also resonance at the part; she complained much of "rheumatic" pain in both her shoulders and elbows, but less of pain in her side; the bowels had been regular till this day, when she passed three or four large evacuations, consisting chiefly of dark coagulated blood. This continued till the following day, when she sank.

Inspection was made seven hours after death. The body was slightly wasted. The lungs were healthy in structure, on the right side there were firm adhesions at the base. The pericardium contained about ʒijj of serum. The heart was a little enlarged, but there was no valvular change. There were redness and effused lymph over the peritoneum on the right side; and the cæcum was pushed a little upward and glued by recent adhesion to the posterior wall of the abdomen, and to the transverse colon. The liver was large and fatty; the gall-bladder contained two calculi, each about the size of a marble. The spleen was rather enlarged. The kidneys were large, soft, and of loose texture, and showed but little distinction between the cortical and medullary structures. Numerous small cysts were found in their substance; and the capsule readily separated. On detaching the ascending colon from its adhesions, which was done very easily, a quantity of dark clotted blood escaped, and the bowel was found to form the anterior wall of a large cavity, filled with coagulated blood. It occupied the position of a psoas abscess, and extended behind the psoas muscle, and even for a short distance below Poupart's ligament. The blood was confined by fascia, and occupied the whole of the right iliac fossa. There was a small portion of fibrin in it, but no pus could be detected; several nerves were found passing

from the spinal column through the middle of it; and the iliac vessels lay on the inner side and slightly in front of it. The artery was healthy throughout; the vein was roughened posteriorly, and formed a portion of the boundary of the cavity described. At the termination of the common iliac vein on the right side was a large irregular opening posteriorly, forming a communication between the vein and the cavity in the muscle. Above this point the vessel was healthy; below, and throughout the external and internal iliac veins, the coats appeared thickened, and the vessels were closed by firm adherent clot. Around the opening on the inner side, there were small excrescences of lymph, almost similar to those sometimes found on the valves of the heart. The lumbar portion of the spine was removed, and carefully examined, but no caries nor other disease could be found. The sacro-iliac synchondrosis was sound, and no diseased bone could be detected at any part. The small intestine was slightly injected at one or two points. The large intestines were healthy, except the ascending colon; there was some vascularity, and an injected appearance in small isolated spaces. On its posterior wall were four or five large ragged openings, varying from the size of a sixpence to a shilling. There was a brown discoloration of the bowel for some distance around the openings, and their edges were thin, having the mucous membrane entire as far as the border of the opening. The peritoneal covering was roughened and uneven, being generally wanting round the edges of the openings, so as to give them a levelled appearance on their outer aspect. The colon contained a small quantity of blood. There was thickening of the cellular tissue about the uterus on the right side; and a circumscribed abscess, the size of a small orange, was found between it and the bladder; but this was entirely distinct from the cavity in the psoas. The ovaries were small and flabby, and did not present any corpus luteum. The uterus was contracted to the size of an orange and felt soft; the os was dark and discoloured. The uterus was covered internally by a thin layer of dark clot, and to its posterior wall was firmly attached a fibrous mass of the same colour and appearance, as if a portion of the placenta had remained fixed. The structure of the walls was healthy.

Many of the earlier symptoms in this case arose from threatening miscarriage after a day of hard work; and it appeared probable that a state of pyæmia was afterwards produced, and would have probably terminated fatally, if the hæmorrhage from the divided cava had not led to comparatively sudden death. The cause of the abscess behind the colon, which extended into the cava, and afterwards into the large intestine, was not apparent; it arose either from some laceration of muscular fibre of the psoas and subsequent suppuration; or it occurred as one of the secondary deposits in pyæmia. It is exceedingly improbable that the vein was primarily lacerated; we are rather disposed to believe that ulceration

perforated its coats. The extension into the colon was evidently from without, the larger size of the ulcer on the external aspect, and the sudden discharge of blood by stool, showed this to have been the case.

CASE CCXIX.—*Abscess in the Hypogastric Abdominal Parietes, simulating Ovarian Disease. Recovery.*—Emma N—, aged about 25, a cook, was admitted under my care into Guy's Hospital, October 13th, 1856. She was a single woman, and had enjoyed good health till twelve months previously, when she had violent pain in the left side, which was said to arise from inflammation of the uterus, and she was leeches, poulticed, and blistered. Three weeks before admission a swelling was perceived in the left iliac region, which was exceedingly tender on pressure.

On admission, at the lower part of the abdomen, between the hypogastric and left iliac regions, a hard, unyielding tumour was detected, which was slightly tender and painful on pressure; it extended obliquely towards the median line of the abdomen, and appeared to be beneath the muscles; there was no redness of the skin, and the pain was only manifested on pressure. There was no febrile disturbance, and the skin was moist; rest was enjoined, and the infusion of roses with magnesia was prescribed. It appeared closely to resemble an ovarian tumour.

Ten days after admission the tumour became more painful, and on the 27th the skin slightly inflamed; leeches were applied, and cataplasma, &c.

November 2nd. The character of the tumour was now manifest, fluctuation was distinct, the pain superficial, but severe. On the 4th, the abscess was opened by my colleague, Mr. Forster, and a pint of pus was evacuated. The abscess slowly healed, and, on the 23rd, she left the hospital convalescent.

CASE CCXX.—*Suppuration external to the Right Kidney. Fibroid Thickening of the Tunic of the Kidney. Chronic Pyelitis. Obliteration of the Vena Cava. Adhesion. Thinning, and doubtful Perforation of the Ascending Colon.* Hugh M—, æt. 36, was admitted September 12th, 1855. Six months previously, he had received a blow on his back, and suffered directly afterwards from hæmaturia; this discharge of blood continued for three weeks, and severe pain in the back came on. The pain continued, and gradually a swelling formed in the loins to the right of the spine. After the blow he had several rigors, and some febrile excitement. The abscess in the loins was opened on November 26th, and from that date continued to discharge freely. He was a man of middle stature, with long thin hair, and had a haggard and cachectic appearance. The urine contained much mucus; stimulants and steel were prescribed.

November 22nd. He had gradually emaciated, but enjoyed his food; his bowels were regular, his tongue very clean; he had had swelling and pain in his right leg for two weeks. On December 19th, the abdomen and lower extremities were much swelled; the vessels on the surface were prominent, and the larger capillaries of the skin intensely injected, as if new vessels

had been formed; the skin was shining, white and tense. He was in considerable distress, and very prostrate, though not in severe pain. He had a rather troublesome cough. His strength gradually failed, and he sank.

On inspection, January 7th, the abdomen only was examined. The œdema of the lower extremities, and the congestion of the vessels had disappeared. The peritoneum contained about four quarts of clear serum; it was smooth and shining, except on the peritoneal surface of the bladder, and in the right iliac fossa, where it was opaque and covered by numerous granules, from about the size of a pin's head to a pea.

The stomach and ileum were healthy; but the lower part of the latter was of a gray colour. The whole of the cæcum and ascending colon were of a deep iron-gray, and contained much irregular pigment in the substance of the mucous membrane. The cæcum and ascending colon were so firmly adherent, and their walls so thinned posteriorly, that they gave way on removing them, even if they were not already in communication with the abscess behind them, which was probably the case. The whole of the peritoneum and sub-peritoneal tissue around the right kidney to the aorta was much thickened, semi-cartilaginous, and formed the boundaries of a sinus filled with pus; this communicated with the opening made in the loins, and extended down as low as the pelvis. The aorta was normal; the vena cava, about two inches from the liver, was completely occluded by the thickening, and by the fibrinous external deposit; its walls were irregular, puckered, and it contained a clot about the size of an ordinary probe. The right renal vein was obliterated, but the artery was normal. The left kidney was hypertrophied, the right kidney and the ureter contained opaque concrete pus, distending the calyces; the secreting structure was destroyed, and surrounded by a dense fibrous envelope. The bladder was small, thickened, and its mucous membrane irregularly granular. The dense tissue about the right kidney involved the right semilunar ganglion; it could with difficulty be dissected; the cells contained considerable quantity of pigment, and their nuclei were indistinct.

The blow on the loins produced in this instance hæmaturia; and led to abscess, and to chronic disease of the kidney. The effused tissue had become fibrous, and exceedingly dense; the ascending colon was adherent, and apparently perforated, so as to communicate with the abscess; but no fæcal discharge took place from the back, nor was there any evidence of purulent evacuation from the bowels. The superficial abdominal veins were intensely congested, evidently from obstruction of the cava.

CASE CCXXI.—*Fæcal Abscess in the Pelvis, communicating with the Ovary and Bladder, opening twice into the Rectum, and on the Groin.*—Sarah Y—, æt. 24, residing in Lambeth Road, was admitted August 22nd. Till nine weeks before admission she enjoyed good health, and at that time, on going

to breakfast, was seized with sickness, and with great pain in the abdomen; after a few days she had rigors for three or four hours, and the pain, vomiting, and purging continued for seven or eight days more; she then became more comfortable, and improved in health; one week before admission she was again attacked with vomiting, pain and purging; and these symptoms continued for three or four days. On admission her countenance was flushed, there was great pain in the abdomen, and intolerance of pressure; the countenance was anxious, and there was general tremor; the respiration was thoracic. She was somewhat emaciated, the thoracic viscera were healthy, the motions were scanty and slate coloured. She had menstruated a week before her illness came on, but during that period she always suffered from pain.

↵ Ten leeches were applied, and grey powder gr. iij, with opium gr. $\frac{1}{2}$, three times a day, were given, and low diet only allowed.

On the 31st, there was great tenderness of the abdomen; the knees were flexed, &c. Ten leeches were again applied, and a poultice; and on September 4th, 1 gr. of calomel and opium were given three times a day.

September 6th. The pain increased, and on the 8th the bowels were relaxed, and the mouth became affected by the mercury. The diarrhœa continued, and became very troublesome. Dover's powder was then given, afterwards sulphate of copper and opium; and starch and opium injection was used. The pain continued very severe till her death, and for some days the motions appeared to be of a purulent character. A few days before death a feculent abscess opened in the right groin below Poupart's ligament.

On inspection, the thoracic viscera were healthy; there were some pleuritic adhesions, but no tubercle.

Abdomen.—The general cavity of the peritoneum, except in the pelvis, appeared healthy; the viscera were collapsed; the small intestine was adherent to the brim of the pelvis, and to the cæcum.

The stomach was pale, but on microscopical examination it was found to be healthy. The small intestines were healthy; the colon, as far as the sigmoid flexure, was much contracted, and contained small scybalous grains, firmly attached to the mucus of the intestine. The viscera of the pelvis were found to be firmly united by adhesions. On taking out the uterus some superficial ulceration was found at the os uteri; the left ovary and tube were normal; but on the right side, at the position of the Fallopian tube and ovary, was a sac capable of holding about 3ij of fluid, filled with purulent and feculent secretions; the abscess communicated by an irregular opening with the first part of the rectum, and extended into an irregular abscess, containing fæces, situated between the ovary and rectum. This fæcal abscess passed downwards towards the bottom of the pelvis, and opened again into the rectum by a small circular opening about three inches above the anus; its boundaries were exceedingly irregular, burrowing beneath the pelvic fascia; it extended to the bladder, and had perforated its fundus by a circular opening; it also passed upwards to the psoas muscle, and reached the crest of the ilium; below it passed beneath Poupart's ligament, and at Scarpa's triangle formed a large ulcerative opening, about two

inches in diameter, which was covered with *faeces*. The mucous membrane of the rectum was congested and gray; its calibre at the part between the two openings was diminished. The bladder was small, and its mucous membrane was red, but the cavity did not contain any *faeces*, nor did it appear that any had passed by the urethra. The kidneys and spleen were healthy; the liver was fatty, moderate in size; its weight 2 lbs. 7 oz.

In this case, inflammation appeared to have commenced in the right ovary; suppuration and local peritonitis followed; communication then took place with the rectum; *faecal abscess* was the result; this burrowed beneath the pelvic fascia, and formed a second opening into the rectum, one into the bladder, and along the *psoas* muscle it reached the skin upon the thigh.

The diarrhoea proved to be *teneamus* with purulent discharge, and is an indication of one of the fallacies in the diagnosis of dysentery. There was some ulceration at the os uteri. The severe pain partly arose, I doubt not, from pressure on the nerves in the *psoas* muscle, namely, ilio-hypogastric, lumbar, and last dorsal nerves, &c., and from the attacks of local peritonitis. Had the diagnosis in this case been more clear, we doubt not that the mercurial and depletory measures would not have been so fully employed.

CASE CCXXII.—*Abscess External to the Rectum leading to Perforation. Considerable Fibrous Thickening, and Simulation of Cancerous Disease.*—Ann C—, æt. 55, a needlewoman, living in Bermondsey, was admitted into Guy's Hospital, March, 1857. She had been married eighteen years, and had had one child. Till ten years before admission she had enjoyed good health, when she fell and struck the sacrum; she suffered much, and was confined to her bed; from that time there had been a constant discharge from the rectum, with pain in defecation, and she had been unable to stand or walk on account of the pain. Three years later she was supposed to have inflammation of the uterus; but the catamenia had continued regularly till thirteen months before admission. For three or four months the left leg had been swollen, and for two months the abdomen had been tense and painful. She was a small delicate woman; the abdomen was tense and tympanitic. The chest was normal, and the heart feeble. There was a constant discharge of mucus and slimy secretion from the rectum; and its calibre was much contracted. The urine was albuminous. Her strength gradually became exhausted, and she died in a few weeks.

The walls of the rectum were exceedingly rigid, dense, and fibrous, and communicated with an abscess, or rather sinus on the concavity of the sacrum. This chronic inflammatory action had extended to the neighbouring parts, and led to stricture.

During life this was supposed to have been a case of cancerous disease, but the post-mortem inspection did not confirm this supposition; it showed that inflammation commenced after the fall, in the cellular tissue external to the bowel, and led to the dense fibrous constriction. This condition of rectum leads to abundant discharge of mucus, simulating diarrhoea or dysentery. A similar state of rectum was observed in a case under my own care, associated with albuminuria; there, however, limited to the walls of the rectum, and not as here produced by external inflammation. This case is another evidence of the severity of pain, when the inferior portion of the rectum is diseased, as compared with the immunity from it, in disease of the sigmoid flexure. The nerves are more easily compressed, and the outlet is also freely supplied with nerves of sensation.

CASE CCXXIII.—*Multilocular Ovarian Tumour. Perforation of Cecum. Fecal Abscess. Pneumonia. Pus in the Ovarian Veins.*—Martha L—, æt. 33, was admitted into Guy's Hospital, December 12, 1855. She was a married woman, who had resided at Poplar. She had had five children, and the youngest was fourteen months old at the period of her admission; but before the birth of her child, she had had pain occasionally, of a severe character, in the left iliac region. After parturition, swelling of the abdomen increased, but with scarcely any pain for a time, till the exertion of walking produced it severely; she stated that the tumour fell from side to side. Four months before admission great pain came on in the left iliac region, extending to the hypochondrium of the same side. The abdomen afterwards increased much in size, and the catamenia ceased. She had also suffered from diarrhoea.

On admission, she was pale, her countenance was expressive of anxiety; the mind was active. The abdomen was swollen and tense, and the skin about the umbilicus was red and inflamed; there was dulness on percussion in each iliac region, but especially on the left side, and on that side a tumour could be felt extending towards the loins. The abdomen was tympanitic at the umbilicus, and for a short distance on either side. The urine was high coloured, and contained lithates. The bowels were relaxed; there were nausea and loss of appetite. She was ordered Dover's powder three times a day, a blister to be applied to the abdomen, and milk diet. The diarrhoea became persistent, and was with difficulty checked. On December 20th, paroxysmal pain in the back came on, and the abscess at the umbilicus had broken and discharged feculent pus, with relief to the patient; opium and chloroform were given. The fecal discharge through the parietes continued till death, which was preceded by an aphthous condition of the mouth, and by violent retching and vomiting, and by aggravation of pain in the abdomen. She lingered till February 5th.

On inspection, acute disease was found at the base of the right lung ; the pleura at that part there was effusion of albuminous lymph, and considerable ecchymosis ; the lung at the same part was consolidated, red, and granular. The bronchi contained much frothy mucus. There was granular deposit on the surface of the right auricle. In the abdomen, the stomach and transverse colon were found moderately distended ; and reaching from the umbilicus to the pelvis was a tumour, composed of ovarian cysts ; it was connected with the left ovary, and filled the left iliac fossa in front of the sigmoid flexure. On cutting down the median line, an abscess was opened extending to the right, into the cæcum, the anterior surface of which was destroyed. The abscess was formed in front by the anterior abdominal parietes, which had become perforated ; to the left by the ovarian tumour to the right by the cæcum and kidney ; below, by the fundus of the uterus and by the rectum. The walls of the abscess were covered with lymph, and the cavity contained fæces. The ovarian growth was six to eight inches in diameter, composed of cysts, some capable of holding several ounces of fluid others almost microscopic ; the fluid was tenacious and gelatinous ; and the walls of the cyst were vascular ; near the cæcum one of these cysts appeared to be connected with the abscess. The large ovarian veins extending into the tumour were filled with thin pus. The right ovary was small and atrophied. The uterus was healthy. The mucous membranes of the cæcum and ileum were much congested. The stomach was pale, and presented gastric solution at its greater curvature. The liver, kidneys, spleen, and mesenteric glands were healthy.

The ovarian disease was the commencement of the fatal affection ; one of its cysts had become adherent to the cæcum, and had led to perforation ; local peritonitis and faecal abscess followed. As to the ovarian disease, it was of the ordinary multilocular character, and allied to carcinoma. The diagnosis of ovarian disease was not certain ; moveable carcinomatous tumours become developed in the omentum ; and the position of the growth was at first that often presented by diseased glands about the kidney ; it extended less into the loins, however, than is usual in the latter disease. The local peritonitis and perforation of the intestine were evident ; and the treatment most likely to afford partial relief was that adopted, namely, the administration of opiates and of nourishment to sustain the patient.

CASE CCXXIV.—*Ovarian Tumour filled with Fæces and opening into the Ileum. Pneumonic Phthisis.*—Catherine S—, æt. 47, was admitted into Guy's Hospital, December 14th, 1853, and died March 23rd, 1854. She had been a washerwoman, and had resided at Clapham. For sixteen years

the abdomen had gradually enlarged, but she had followed her occupation till five months before her death, when, during menstruation, she took cold, and suffered from pain in the abdomen. On admission there was severe pain in the whole of the abdomen, with loss of appetite, thirst, and want of sleep. The countenance was pale and anxious; the tongue was brown; the skin was hot and dry. Ovarian disease could not be detected by vaginal examination. Symptoms of pneumonic phthisis and pleurisy came on, and she died about three months after admission.

On inspection, the left pleura was found full of pus; the right lung contained a vomica, and presented indurated lung tissue around it. Peyer's glands were ulcerated. In the cæcum was a small ulcer, and another was found in the appendix, which contained fæces. The right ovary was diseased, and constituted a cyst, five inches in diameter, with thickened walls. The cyst was firmly adherent to the lower part of the ileum, and communicated with the intestine. It was filled with fæces, and was adherent to the uterus at the lower part. The left ovary was atrophied; the cervix of the uterus was elongated and thickened; the kidneys were pale; and the liver was fatty.

The commencement of the affection strongly indicated ovarian disease; suppuration took place around the ovary, and led to symptoms resembling peritonitis; pus was probably discharged by the bowel. Inflammatory disease was afterwards developed in the lung, and led to a fatal termination.

CASE CCXXV.—The following is one of a very unusual character, and the preparation is in the Guy's Museum, Nos. 2516 and 2517.

Extra-Uterine Fœtation, opening into the Sigmoid Flexure.—Elizabeth H—, aged about 20, had led an irregular life, and had had a child eighteen months previous to admission; she was not aware that she was pregnant; she had been ill for six months, but had only been confined to her bed for three or four weeks. She was in a state of extreme prostration, and had obstinate diarrhœa; the evacuations consisted of blood and pus; there was slight tenderness of the abdomen, some fulness, but no defined tumour. She died in sixteen days.

Inspection.—Abdomen.—In the pubic region there were firm adhesions, and there was a cavity bounded by the ascending colon, by the lower part of the sigmoid flexure; posteriorly, by the rectum and sacrum; anteriorly, by the parietes, by the pubes and bladder; and laterally by the pelvis. This cavity was filled with pus, and contained a decomposing fœtus about three months old. There were traces of placenta; and extending into the sigmoid flexure was an opening two to three inches in length and one in breadth. The uterus was small, and no decidua was present in it.

In the Guy's Reports of 1838, Dr. Bright records a remarkable case of abscess of the spleen, which perforated the descending colon; the diagnosis was exceedingly obscure.

The patient was a young woman, aged twenty-five, much emaciated, of peculiar sallow complexion and anxious countenance; she had great uneasiness and pain in the abdomen particularly at the scrobiculus cordis and right hypochondrium; food increased the pain; the vomiting was constant, sometimes directly after food had been taken; there was also occasional bilious vomiting; the tongue was dry and glossy, and she gradually sank. On inspection, the lungs and heart were found to be healthy; the liver was hard and granular; the lower part of the spleen was occupied by an abscess, which was firmly adherent to the transverse colon, and which had opened into it. There was also an abscess in the left ovary.

The record of these cases shows their great variety; each one presents a difference in the symptoms, and must be separately considered. Great relief may be afforded in the earlier stage by local depletion; and later, in many instances, by opening the abscesses which may have formed, and by the judicious administration of opium, tonics, and stimulants.

CHAPTER XVII.

ON PERITONITIS.

REFERENCE has frequently been made in the preceding chapters to disease of the serous investment of the intestines, the peritoneum ; and we propose now to review the general forms of peritonitis, and to notice the pathological appearances presented, the symptoms by which it is indicated, the causes which produce it, and then the treatment best calculated for its removal.

The earliest pathological appearance which the peritoneum presents, when acutely inflamed, is congestion of the capillary vessels, with a loss of the smooth and shining character of the serous membrane ; dryness then takes place, from the diminished secretion, and slight opacity follows. This congestion is especially manifest where the coils of the intestine come into contact, the one with the other ; in very acute disease, however, the whole surface is reddened. The inflammatory œdema which gives opacity to the serous membrane is an indication of the sub-peritoneal coat being involved ; the muscular coat next participates in the disease ; its contractile power is diminished, and the intestine distends, producing the tympanitis of which we shall have to speak ; and the coats of the intestine may be readily separated from each other ; effusion of lymph then takes place, and it is often found to cover over the intestine in large flakes, mixed with serum in the more depending situations ; if at a very early stage, or slight in degree, this lymph is represented by a mere stickiness and very slight adhesion between the intestines. The serum is sometimes present in considerable quantities, and in chronic peritonitis fills and distends the serous sac ;

but the conditions of the fluid effusion vary according to the character and the intensity of the disease; sometimes it consists of clear serum holding in it portions of semi-coagulated fibrin; at other times the serum may be almost of the character of pus, and between these extremes every gradation is observed. In proportion as the effusion is less organised it is corpuscular in character, and approaches proportionately to the appearance of pus. As the peritonitis subsides, these effusions may gradually become absorbed; but more generally they leave traces of their presence, the lymph becomes organized, fibres are produced, and vessels become developed in them; adhesions between portions of intestine are thus formed, either uniting them closely together, or constituting bands, which may be the cause of subsequent internal strangulation;* the lymph which covers the viscera, as the liver and spleen, often forms a thick investment, which gradually becomes firm, fibrous, and cartilaginous in density, and ossific particles are occasionally developed in it, especially in the splenic adhesions; at an early stage these investments may be stripped off from the proper tunic of the viscus, but afterwards they become inseparably united. Again, sometimes the lymph is spread generally upon the viscera, or it forms rounded patches, or it has a cribriform appearance, as if rounded portions had been removed or absorbed. Lymph deposited upon the mesentery and omentum leads to gradual contraction, to shortening, and fibroid thickening. In some cases the serous membrane has a granular appearance; these minute deposits of fibrin being more highly organized than the tubercles in strumous peritonitis. These granules must not be mistaken for an appearance sometimes found several days after death, when the serous membrane appears sprinkled with grains of sand; these are of a crystalline character, triple phosphate, and are probably formed after death. Lastly, acute peritonitis may lead to a general thickening of the serous membrane without adhesions; the membrane becomes dense, whitish, thickened, and fibrous, and this arises from the fibrinous effusion taking place in the substance of the membrane rather than on its sur-

* Bands leading to strangulation do not always have an inflammatory origin. See an interesting pamphlet on this subject, by Mr. Gay, &c.

face. In *chronic* peritonitis the same general changes are produced; lymph may be effused, become organized, and form adhesions or produce thickening of the serous membrane; or effusion of serum takes place, so as to distend the serous membrane, even without previous acute symptoms. The thickening of portions of peritoneum covering the liver which is frequently found after the pressure from stays and belts, consists of a fibroid degeneration of the serous membrane itself.

Peritonitis is either general or local; the former is often rapidly fatal; the latter is in many instances conservative in its action; and in perforation of the intestine from numerous causes, it limits or it prevents extravasation from taking place. If extravasation have occurred, it may be so localised by peritoneal adhesions that life is prolonged for weeks, or even for many months; and when acute inflammation in the abdominal glands or viscera takes place, local disease of the peritoneum in the neighbourhood is set up; thus the peritoneum covering the liver becomes involved in hepatic disease, and that in the pelvis in acute disease of the ovary.

Strumous peritonitis has already been described in speaking of strumous disease of the alimentary canal generally, as forming—1st, miliary tubercles on the visceral, parietal, and mesenteric peritoneum. 2nd. As strumous deposit on the peritoneum and in the glands, associated with inflammation, and leading to matting together of the intestines, sometimes to perforation from without, and to the formation of fæcal abscess. 3rd. As peritoneal ascites with miliary tubercles; and, 4th, with tympanitis.

Cancerous disease of the peritoneum occurs either as a primary disease of a scirrhus, medullary, and colloid form, or as a secondary one from the extension of the disease to the peritoneum from the intestine. Thus, in cancerous disease of the stomach, or of the cæcum and colon, tubercles with inflammatory lymph are often found upon the peritoneum; so also with cancerous disease of the ovaries and of the mesenteric and lumbar glands; but in advanced life especially, primary cancerous disease of the peritoneum is not unfrequent; an immense number of tubercles, some semi-transparent, others red and

congested, from the size of grains of sand to split peas, are found everywhere studding the serous surface, mixed more or less with ordinary fibrin and lymph; sometimes the deposition of pigment in these tubercles gives a black colour to them, and a form of melanosis is produced; this appearance, however, is to be distinguished from the blackness arising from effused blood. This form of cancerous disease is generally associated with considerable serous effusion, and it is more common in women than in men. Again, sometimes cancerous disease, especially involving the omentum or the mesentery; and it is generally, when so situated, either of a medullary character, constituting large masses, composed of nuclei and nucleated cells, or of scirrhous character, when the omentum and mesentery become thickened by a fibro-cancerous growth. The omentum, in these cases, often forms a dense, hard mass, extending across the abdomen at the umbilical region; it is then several lines in breadth and almost semi-cartilaginous, but still more or less transparent; and serous or fibrinous effusions may have taken place in considerable quantity into the serous sac. In some cases, both of minute cancerous tubercles on the peritoneum, and of the more general thickening to which we have just referred, it is very difficult to draw an exact line of demarcation between the simple products of ordinary inflammation and those of true cancerous disease. If a large quantity of pigment be deposited, as we sometimes find in medullary cancer of the omentum, &c., the term melanotic cancer is applied. Colloid cancer affects the peritoneum, either partially, as of the omentum, or generally, uniting almost the whole of the abdominal viscera; it presents the semi-gelatinous appearance of colloid with minute intersecting bands, and with the microscope nucleated cells are seen, separated by delicate fibres. This colloid disease of the peritoneum is either primary or it follows from colloid affection of the stomach, the cæcum, &c.

The *symptoms* of acute peritonitis are generally very characteristic, as when, for instance, the stomach and appendix cæci are perforated by ulceration, sudden intense pain comes on, the patient is "doubled up," unable to move, and lies with the legs flexed; the countenance expresses the intensity of the suffering as well as the serious nature of the disease; the distress and pain

are evident in the features, the eyes are sunken, the face is pallid, the abdomen very shortly becomes distended, tender, and tympanitic; no pressure can be borne, and even the weight of the bedclothes becomes insufferable; the pulse is small, compressible, and if reaction take place from the first sudden collapse, it becomes more hard and frequent, wiry; the bowels are generally confined, especially at the early stage of the acute disease, but sometimes towards the close of the malady diarrhoea may supervene. The urine is scanty, and if the vesical peritoneum be involved, retention often takes place. If the peritoneal surface of the stomach be implicated, vomiting is a frequent and distressing symptom, and green bilious fluid is ejected. The mind may be conscious and strong throughout, even when the powers of life are fast failing, and the pulse scarcely perceptible at the wrist.

In many cases of perforation the patient scarcely rallies from the first sudden collapse, and death takes place in five to ten hours after the onset of the disease; in other instances, however, the signs of febrile excitement are more evident, as shown by heat of skin, especially of the abdomen, by thirst, and by a frequent and hard pulse. If the disease tend to an unfavorable termination, the prostration increases, the patient is restless, the tongue dry and brown, the pulse compressible, failing, and irregular; the extremities become cold, a clammy sweat breaks out, hiccough comes on, and then death follows, the patient often remaining sensible till the close, and the subsidence of pain, as life is ceasing, occasionally gives to the superficial observer a false hope of recovery. On the contrary, when the vomiting subsides, the pain and distension lessen, the countenance becomes less haggard and dejected, the pulse soft and less frequent, but tolerably firm, and especially, when the patient has refreshing sleep, we may regard the immediate danger as less imminent. Gradually all the symptoms may disappear, and the patient completely recover, with perhaps some peritoneal adhesions and thickening. It may be that effusions take place, which are more gradually absorbed, or become very persistent; again, if faecal extravasation have occurred, repeated attacks of local peritonitis, with hectic fever, follow, or renewed general inflammation destroys the life of the patient.

Chronic Peritonitis may commence with the symptoms of acute disease; and recurrent pain, distension and effusions take place. This recurrence of symptoms is more especially the case when strumous and cancerous tubercles have been developed, and perpetuate the serous disease. In these instances the symptoms are less severe, although of the same general character; pain with distension and tympanitis, and with a peculiar haggard expression, which is very characteristic of abdominal affection; the abdomen is at the same time hot, and tender on pressure. In *strumous* peritonitis, the pain is paroxysmal, and often resembles severe colic; the bowels are often irregular, the stomach sometimes irritable, the tongue red and injected, the patient fretful, and as the intestines become matted together by adhesions, the viscera move *en masse*, and a doughy sensation is communicated on manipulation. Or these strumous and inflammatory adhesions may be local, stimulating abdominal tumours. The indications of disease are also associated with general strumous cachexia, and are often complicated with pulmonary disease. Too frequently hectic supervenes, and this is especially the case when faecal abscess has been produced, and the hope of ultimate recovery is then almost taken away. In strumous peritonitis also the pain may be very slight, whilst effusion gradually takes place to a considerable extent, as we sometimes find in children after measles, &c., or excessive tympanitis may be produced without any acute pain.

In *cancerous* peritonitis there is the same expression of distress on the countenance, with pain and heat of the abdomen, with gradual distension, and often with serous effusion; if the disease consist in cancerous growth in the omentum and mesentery, a tumour may generally be felt; if it extend to the peritoneum from the stomach, liver, intestine, ovary, &c., the earlier symptoms will be indicative of the primary malady. It must, however, be remembered, that in both strumous and cancerous disease the symptoms of acute peritonitis may be suddenly developed in the course of the chronic disease, from perforation of the intestine or otherwise, and lead to a speedy fatal termination. This acute mischief is sometimes set up by the attempt to relieve the serous effusion by paracentesis, the persistent irritation and congestion of the serous membrane

passing very readily into general and acute inflammation. In colloid disease of the peritoneum, the symptoms are often obscure—pain, with heat and more or less distension of the abdomen; diarrhoea is sometimes present, but, if the intestine become involved, other diagnostic symptoms arise.

Diagnosis.—The pain of peritonitis may not only be absent, on account of the peculiar character of the disease, but the patient may be rendered unconscious of it from cerebral oppression, or from the dyspnoea and distress of pulmonary and cardiac disease fully engrossing the sensibilities of the sufferer. Still there are painful conditions of the abdomen for which peritonitis may be mistaken.

1st. Flatulent colic. The pain and distension are in this disease sometimes very severe, the countenance may be haggard and distressed, and collapse sometimes results; but there is not the tenderness of peritoneal inflammation, the symptoms are less persistent, the pulse less affected, the collapse rarely so profound. 2nd. In hysterical affections of the abdomen, the pain is very superficial, and firm pressure can frequently be borne, whilst the patient almost shrieks before the hand has reached the surface; the countenance does not express the distress of serious organic disease, the pulse may be almost unaffected; still, in this disease, we have seen the patient bled from the arm to syncope, with the idea that acute disease existed. 3rd. The vomiting and sudden pain of perforated intestine are sometimes mistaken for gall-stone; but the latter disease is free from the acute tenderness and distension of peritonitis. 4th. Neuralgic pain from disease of the spine, of a functional or organic character, often simulates peritonitis; but here, also, there is an absence of tenderness on pressure of the abdomen, of distension and tympanitis, as well as of the general expression of peritoneal disease; the pain is situated in the course of the spinal nerves, and often extends over the crest of the ilium in the course of the last dorsal nerve, or into the groin and testicle in the course of the genito-crural nerve; there are also, generally, some indications of spinal disease in local pain of the vertebræ, with modified motion and sensibility of the lower extremities, and loss of power of the sphincter muscles. 5th. Suppuration of the abdominal parietes is at an

early stage very difficult to distinguish from peritonitis. 6th. The pain from the distension consequent on the enlargement of abdominal tumours and effusions may easily be mistaken for peritonitis, as, for instance, in aneurism, in ovarian and cancerous tumours, and in dropsies; but in these cases, as we have before said, peritoneal disease is often set up in the progress of the malady. 7th. During the course of peritonitis, the muscular fibres of the bladder sometimes fail to contract, apparently from loss of power, and the urine is retained, thus closely simulating simple retention of urine; on the contrary, we have also witnessed distension of the urinary bladder from enlarged prostate or other cause, producing pain which resembled peritonitis, and which had been sent to the hospital as a case of abdominal tumour.

In reference to the *prognosis* of peritonitis, instances of intestinal perforation are generally quickly fatal, in five to ten hours; but in some cases this issue might be prevented if perfect rest were maintained, stimulants and purgatives avoided, and opium given for by movements of the body extravasation is increased, and we have seen castor oil floating in the peritoneum after perforation had taken place. In other cases we must be guided in the prognosis by the cause of the peritonitis, whether such cause be of a remedial character or not. As to those symptoms which *immediately* indicate a favorable or unfavorable termination we have already alluded to them in describing the general characteristics of the disease.

Causes.—Although peritonitis is spoken of and treated as an idiopathic disease, we do not find that it has a local character unless excited by injury to the serous membrane, or by the direct propagation of disease; and to elucidate this part of our subject reference has been made to inspections made at Guy's Hospital during a period of twenty-five years; 3752 inspections were made, and 501 were instances of peritonitis; but we cannot find a single case thoroughly recorded in which disease could be correctly regarded as existing solely in the serous membrane.

In relation to its causes, cases of peritonitis may be divided into three classes. 1. Peritonitis produced by the extension of disease from adjoining viscera, or excited by direct injury, including cases of perforation of viscera, extravasation, violence

&c. 2. Peritonitis connected with blood changes, as when inflammation of the serous membrane occurs in the course of albuminuria, pyæmia, puerperal fever, erysipelas, &c. 3. Peritonitis caused by general nutritive changes of the system, which have been followed by acute or chronic disease of the peritoneum, such as struma, cancer, &c.; and comprising also those cases in which the circulation of the peritoneum has been so altered by continued hyperæmia (modifying its state of growth), that very slight exciting causes suffice to induce acute mischief, as occurs in peritonitis with cirrhosis, disease of the heart, &c. In reference to the instances found amongst the large number of inspections just referred to, we find—

1. Peritonitis from direct extension . . .	261
2. Peritonitis connected with blood changes .	94
3. Peritonitis connected with general or local perverted nutrition	146
	<hr/> 501

It will be evident that the first class might be regarded as disease of a local kind, and the second and third as of a general character. The *first* division includes 261 instances of peritonitis, and of these cases 102 were produced by hernia, internal and external, intussusception, bands of adhesion, and cancerous obstruction. In 19 cases the obstruction was of an internal kind, and in not a few of these death followed from rupture of the intestinal coats. 35 cases were caused by injuries or operations directly affecting the serous membrane. In some instances of severe abdominal injury, death resulted before any sign of inflammation had taken place, whilst in others the injury had been of such a character that no treatment could be more than palliative in the most trifling degree, as when the jejunum was completely divided by a vehicle passing over the body; but the Museum of Guy's Hospital contains a specimen where severe injury of the abdomen had taken place, and death had ensued in about three months, and in which a small laceration of the liver had evidently become united.

Among the operations referred to, one was for the removal of

an ovarian cyst; one a case of gastrotomy; fourteen were cases of paracentesis abdominis, which was performed in five to relieve ascites accompanying cirrhosis, in two for ascites with heart disease, and in seven to empty large ovarian cysts. In instances of ascites from heart disease, chronic bronchitis, and cirrhosis the whole of the peritoneal capillaries are in a state of continuous hyperæmia; the serous membrane becomes opaque and thickened, and a very slight fresh exciting cause is sufficient to produce acute disease. It is probable that if a larger number of instances of paracentesis abdominis had been taken, it would have been found that in ovarian disease, especially of persons advanced in life, paracentesis is much less frequently followed by severe and fatal result than in ascites following cirrhosis.

Perforation of the intestine into the peritoneal sac constitutes a most important cause of peritonitis, and we shall briefly notice the several varieties. Out of the 501 cases of peritonitis perforation occurred 56 times, namely:

- 10 from hernia;
- 9 from disease of the stomach;
- 15 from fever-ulceration of the ileum;
- 4 from strumous disease;
- 11 from disease of the cæcum and appendix;
- 1 from cancer of the vagina;
- 4 from cancer of the colon;
- 2 from ovarian adhesions.

56

This fatal lesion in fever generally supervenes about the twenty-first or twenty-second day of fever, and many instances are no doubt accelerated by muscular movements; the period however, of perforation, is subject to considerable variation occurring sometimes as early as the tenth day; in others, being postponed till the fifth or sixth week, when the patient may seem convalescent. We have before referred to the fact, that in fever, peritonitis is set up by ulceration, although the peritoneum may be entire, transudation having taken place in a sufficient degree to produce acute mischief.

In strumous disease the sudden and intense peritonitis from perforation is generally prevented by adhesions ; occasionally, however, these limitations of the mischief are incomplete ; in the ulceration of phthisis, as well as in strumous disease of the peritoneum, this sudden termination is found to occur.

Diseases of the cæcum and of the appendix are still more frequent causes of peritonitis, and in these affections perforation often takes place unexpectedly. In perforations, however, of every kind, the extravasation may be circumscribed, and local peritonitis and fæcal abscess be the result ; numerous instances of this kind are detailed in preceding chapters.

Another frequent source of peritonitis consists in the extension of disease from the bladder, uterus, and other pelvic viscera ; and in less severe degrees local peritonitis is often observed from ovarian irritation. The enumeration of the pelvic origin of forty-two cases from amongst the numbers we have before referred to will best illustrate the nature of this cause :

- 6 from ovarian disease ;
- 1 from ulcerated vagina and uterus ;
- 1 from strumous disease of the testicle and castration ;
- 1 from diseased prostate ;
- 1 from strumous pyelitis ;
- 1 from fistula in ano ;
- 1 from cancerous disease of the bladder ;
- 1 from polypus in the bladder ;
- 14 from cystitis, calculus in the bladder, stricture, &c.
- 1 from extravasation of urine ;
- 2 from sloughing perineum ;
- 1 from sloughing of the nates ;
- 10 from lithotomy ;
- 1 from diseased hip and pelvis.

Peritonitis is also caused by diseases of the liver and gall-bladder, and some of these are instances of extreme interest ; thus, abscess in the liver, hydatid disease, gall-stone, primary disease of the gall-bladder, may each induce disease of the general serous membrane. In many cases of chronic conges-

tion and enlargement of the liver, and in cirrhosis, especially of an acute kind, the peritoneum becomes implicated; and the serous membrane is found, when a fatal issue has taken place to be thickened and chronically inflamed; acute peritonitis is in these conditions very easily induced; but we have included these chronic forms of hepatic disease amongst the third class of causes of peritonitis.

In the following instance, it is probable that the peritonitis was secondary to disease of the liver, but there is much doubt as to its precise character; we believe it was one of the acute forms of cirrhosis, quickly passing on to suppurative changes, and followed by peritonitis, consequent on the direct extension of disease to the serous membrane.

CASE CCXXVI.—*Acute and Chronic Peritonitis. Suppuration in the course of the Portal Veins.*—Samuel A—, æt. 35, was admitted into Guy's Hospital February 11th, 1857, and died on the 14th. Nine days before Christmas he had been seized with rigor, pain in the chest, and vomiting, followed by tenderness of the abdomen. He was obliged to give up work; the pain continued in the abdomen, and after a time vomiting of greenish fluid came on. During the whole of the illness he had repeated rigors. He was a spare man, but not emaciated, and slightly jaundiced; the abdomen was full and tender; there was constant vomiting of bilious fluid, and the bowels were confined. The precise character of the disease was obscure till death.

With the exception of pleuritic adhesions, the thoracic viscera were healthy. The peritoneal cavity contained three or four pints of serum, and there was much lymph floating in it; the intestines were matted together, and separated from each other with difficulty. At the upper part of the abdomen, the disease was older, and the adhesions more firm; the colon, stomach, and liver were firmly adherent, and pus was confined in spaces between the viscera; a large quantity of pus was also found between the spleen and the diaphragm. Upon cutting through the lesser omentum, and separating the liver from the duodenum, thus dividing the portal vessels, a large quantity of pus escaped from Glisson's capsule, and from the portal vein. Behind the pancreas was a circumscribed abscess, holding about half a pint of pus. The pancreas was healthy, as was also the duct; so also the splenic and mesenteric veins. In the liver, there was extensive suppuration in connection with the veins; the portal veins were quite distended with pus, their walls softened, and in some parts detached from the tissue around them. In the large veins, distinct flakes of lymph adhered; and in some the lymph was found as a firm stratum lining the cavity of the vessel. The mucous membrane of the intestines was healthy.

Peritonitis is sometimes the result of acute disease of the mucous membrane of the intestine, extending to the deeper

coats, and to the peritoneum itself, leading to inflammation, or to perforation and faecal abscess. Ulceration of the colon was the cause of faecal abscess in three of the instances we have previously mentioned, and a case of simple perforation, was from this cause. In three other instances, acute dysentery, with sloughing of the coats of the intestine also produced peritonitis.

In reference to the etiology of peritonitis, the following table shows not only the large proportion of cases due to simple extension of disease to the serous membrane, but the forms of disease by which peritonitis is induced; 261 instances from 501 are thus referable to direct extension:

From hernia, of which 19 were internal obstruction	102
„ injuries or operations	35
„ perforations of the stomach, ileum, cæcum and appendix, colon, &c. (other 13 men- tioned with hernia or with cæcal disease)	43
„ and leading to faecal abscess (2 cases other- wise mentioned)	17
„ typhoid ulceration without perforation	5
„ disease or operations on the bladder and pelvic viscera, &c.	42
„ disease of liver and gall-bladder, &c.	11
„ acute disease of the colon (3 others enumerated with perforations)	3
„ diseases of cæcum or appendix (9 others pre- viously mentioned)	3
	<hr/> 261

We now turn to the *second* division, namely, peritonitis connected with a changed condition of blood, such as exists in albuminuria, in pyæmia, in puerperal peritonitis, and in erysipelas, &c.

In *albuminuria*, where the renal disease is of an acute kind, lymph may be effused, and the symptoms of peritonitis are often well marked; but in some cases, in which there is small granular kidney, with contracted liver, the peritoneum is thickened, and more chronic disease is found to exist. Albuminuria is a frequent cause of serous inflammation affecting the pleura, the pericardium, the arachnoid, sometimes the joints, and, as these cases show, the peritoneum also. It rarely, however, happens

that the peritoneum only is affected, although such is sometimes the case. The symptoms also are often masked by the distress arising from the general anasarca, and the dyspnoea from cedematous lungs, &c.

Puerperal peritonitis is often associated with suppurative either in the uterine veins, or in the pelvic cellular tissue, or the broad ligaments. But there are sometimes instances in which the peritonitis arises from a general cause, and these may perhaps be considered as pyæmic or even of a rheumatic character, or connected, as we have before said, with renal disease. Thus, peritonitis is found with pericarditis and with pleurisy; with pneumonia and dysentery, or with these conditions renal disease may be also combined. The two following cases are worthy of record in their causative relation.

CASE CCXXVII.—*Hypertrophy of the Heart. Adherent Pericarditis. Acute Pericarditis. Pleurisy and Peritonitis.*—James M—, æt. 9, was a delicate boy, who had suffered from cough, but there was no history of rheumatism; he had been in the hospital for disease of the bones of the foot and left nearly well; in one week he returned very ill, and was found to be suffering from pericarditis; there was slight pain in the shoulder, but no swelling of the joints generally. He died in three days. There was a cicatrix on the foot, showing the part from which the fourth metatarsal bone had been removed; but there was no suppuration. There was general pleurisy on both sides, recent lymph in small quantities being found; the lungs were congested, and the bronchi full of tenacious mucus. There was acute and chronic pericarditis, as shown by adhesions of recent lymph, and some parts by very firm fibrinous bands. Minute vegetations were found on the valves. In the abdomen, although the serous membrane had not lost its transparency, there were some flakes of recent lymph and a small quantity of serum.

Was this general disease of the character of rheumatism? We must regard it as arising from some general cause, and in that respect very different from idiopathic peritonitis.

CASE CCXXVIII.—*Acute Peritonitis. Pericarditis. Pleuro-pneumonia. Small Granular Kidneys.*—William B—, æt. 42, was admitted into Guy's Hospital, June 13th, 1855; he had been a labourer, and had resided in the Borough. A year previously he had had jaundice; and three days before admission had had rigors, but on the following day, although feeling ill, he went to his employment, but the next day he gave up work. On admission he was very ill, presenting the signs of pneumonia of the right lung; the

dulness rapidly increased, with bronchophony; much blood was expectorated, and he became delirious, but whilst sensible did not complain of pain. The urine was not albuminous. He died on the 20th. There was recent consolidation of the whole of the right lung; the pericardium was covered with a layer of fresh lymph; there was also acute peritonitis, the intestines being adherent by recent lymph. There was a hydatid cyst in the liver, surrounded by a dense white envelope, one eighth of an inch in thickness, and containing disintegrated membrane with opaque fluid; the gland itself was fatty.

The kidneys were granular and coarse, the tunic adherent, the secreting tubes filled with inflammatory products.

The *third* and last class comprises peritonitis of a general character, arising from *struma* and *cancer*, in which the peritoneal disease is often very insidious. We include in this division those cases in which, after a prolonged state of hyperæmia, the serous membrane becomes thickened, and a very slight cause suffices to produce acute disease.

The forms of inflammatory disease of the peritoneum in *struma* vary both in degree and in extent. Sometimes tubercles exist partially distributed, as on the peritoneal aspect of ulcers of the intestine; or they are found more generally upon the serous membrane.

These tubercles may be combined with the effusion of lymph, and with the ordinary signs of acute disease; or the changes are more chronic, and the obliteration of the serous membrane has occasionally resulted from the organization of the product poured out between the coils of intestine. Acute mischief frequently supervenes on the old disease, and lymph or pus becomes effused amongst the united intestines; and the pus sometimes finds an outlet by the perforation of the intestine. In one case the whole calibre of the bowel was divided, so that several truncated loops entered into one fæcal abscess. Firm peritoneal adhesions do not always form, and then the perforation of a strumous ulcer of the bowel leads to sudden and fatal peritonitis. A few tubercles are sometimes associated with considerable serous effusion, constituting a very intractable form of dropsy, especially when occurring in the young; in others, the intestines are distended with flatus to a great extent, without the pain of ordinary peritonitis.

These cases of strumous disease of the peritoneum are occasionally found with perfectly healthy mesenteric glands, although

this is not generally the case; in young subjects they are associated with tubercles in the pia mater, and in many others with phthisis.

Cancerous disease is frequently the cause of peritoneal effusions, both of lymph and serum; sometimes it leads to perforation into the serous membrane, but more frequently it is by formation of cancerous tubercles upon the peritoneum, or by extension from other viscera, that the peritoneum is involved. It is exceptional to find the serous membrane only affected; thus in men the liver, stomach, pancreas, and the kidney, and in women, the uterus and ovaries are also implicated; and cancerous disease of the uterus and ovaries during the earlier periods of life, lymph is often effused in considerable quantities.

The only remaining causes are those in connection with chronic disease of the liver and thoracic viscera, producing prolonged hyperæmia of the peritoneal vessels and ascites, and readily terminating in acute peritonitis; hence the danger of tapping in these instances; but acute peritonitis often occurs in cirrhosis from other very slight exciting causes.

In *cirrhosis* the peritoneal covering of the liver is very generally thickened. We do not, however, refer to this merely as a partial affection, but to those in which the serous membrane is generally affected.

In the second and third divisions of cases of peritonitis, the causes were as follows:

From Bright's disease	63
„ pyæmia, 13; erysipelas, 5; puerperal fever, 10; with pneumonia, 3 . . .	31
„ strumous disease	70
„ cancerous disease	40
„ hepatic disease	27
„ heart disease	9

240 out of 501 instances

Treatment.—The consideration of the origin of peritonitis either in its local or general source, is the best guide to proper treatment, whether it arise—1st, from extension of disease from

adjoining viscera, or from perforation and injuries; 2nd, from blood changes, such as occur in albuminuria, pyæmia, and erysipelas, &c.; 3rd, from almost imperceptible changes or deficiencies in general nutrition modifying the state of the general health, as in struma, cancer, and climacteric changes; or from the hyperæmia of the peritoneum, consequent on cirrhosis and chronic disease of the heart and lungs, when upon very slight exciting causes, acute mischief follows. In the first form, if perforation have taken place, perfect rest is exceedingly important, in diminishing extravasation, and in localising the peritoneal mischief; purgative medicines of all kinds should be avoided, and also stimulants, which are often unfortunately given at once, before a medical practitioner sees the patient, in perforation of the intestine. This injudicious attempt to relieve pain by purgatives, carminatives, and stimulants, may deprive the patient of the hope of recovery; for, as we have before said, we have seen castor oil floating in the peritoneal cavity. Food, also, should be abstained from, or only a few spoonfuls administered to relieve thirst; in more chronic forms, not arising from perforation, food of a fluid and bland kind is only admissible; and even when the more active symptoms have subsided, the return to solid forms of aliment must be very cautiously made. When there are symptoms of failing power, stimulants in small quantities may be given, but are best combined with demulcent food, as brandy with arrow-root, &c.

As regards medicinal treatment, we believe the plan recommended by Dr. Stokes and Dr. Graves to be of the greatest value, not only in cases of perforation of the intestine, but where the peritoneum is acutely inflamed from the direct extension of disease. It consists in the administration of opium in full and repeated doses; and its beneficial result arises from its favouring rest of the intestines and the localisation of the mischief, from the mitigation of suffering which it affords, whilst at the same time it alleviates nervous prostration and collapse, and facilitates reparative action. In many instances the opiate plan may be combined with local, and sometimes even with general depletion; anodyne remedies may be applied externally, or counter irritants, as cantharides and turpentine.

Local peritonitis is thus greatly relieved by local depletion external applications, as when produced by ovarian and c disease ; but blisters are of value, especially in those insta in which repeated attacks of peritonitis occur. Mercury, ei in the form of grey powder, calomel, blue pill, or as mercu inunction, is, we believe, injurious in all these cases of a direct peritonitis. It tends to prevent adhesion, it ex peristaltic action, it promotes ulcerative action, it increases depression consequent on the disease, which is often immediate cause of death, and lastly, it renders the intest contents more fluid, thereby increasing extravasation. We well aware that many instances of acute peritonitis from diseæ cæcum, from enteritis, and from ovarian disease, recover a mercury has been given ; but as far as the causes we h enumerated can be any guide, and from extensive experience these cases, we strongly deprecate its use.

Effervescent medicines generally increase the painful dist sion of the abdomen, but diaphoretics and salines are somet of value when combined with opium.

In the subsequent treatment we must not be too desirou inducing action from the bowels, and, when necessary, ger enemata are better than purgatives administered by the mou

When the more active symptoms have subsided, opium n be continued with vegetable tonics or with quinine. If fl effusions have formed, iodide of potassium and diuretics may advisable, and the abdominal glands may then be beneficia stimulated by an occasional dose of grey powder or calom Preparations of iron are not generally well borne in the æ valescence from acute peritonitis. It will often be found t as the health becomes established the fluid effusion rapidly d appears ; in other cases the repeated application of coun irritants may be required, and sometimes it is well to remc the serum by paracentesis.

In the peritonitis of *albuminuria* the best treatment consi in the relief of the general disease by diaphoretic medicin counter irritation and cupping on the loins, and by free evacu tion of the bowels ; but mercurial preparations very read affect the system, producing severe salivation, without cor sponding benefit. When effusion becomes extreme, it is bett

to attempt its removal by puncturing the thighs and by purgatives than by directly emptying the serous cavity. Hot-air baths are sometimes of great service.

In the treatment of the peritonitis of pyæmia and erysipelas the local disease is to be less regarded than the general one, nor should we attempt to cure the peritonitis of this kind by depletion and mercurial preparations. Opium and salines, with the free use of stimulants, are apparently the best remedial agents we can employ. Typhoid symptoms too frequently come on, and precede a fatal result.

In *puerperal* peritonitis the same plan of treatment may be adopted. In some cases the blood becomes affected by the absorption of pus; or pelvic phlebitis and cellulitis are followed by the peritoneal disease; and, from the beneficial effect following the internal administration of tincture of the sesquichloride of iron in erysipelas and diphtheritic disease of the throat, Dr. Heslop has recommended the same remedy in puerperal peritonitis, as being a disease closely allied in character. Those cases which we have seen recover have apparently been benefited by thoroughly washing away uterine discharges, by the free use of opium, and by stimulants; but we are quite prepared to hear further reports of the good results of the tincture of the sesquichloride. In puerperal peritonitis the use of turpentine internally has been recommended, and has been followed sometimes by a beneficial result.

In the treatment of acute peritonitis in *struma*, the same rules ought to be borne in mind as in the treatment of strumous pneumonia. Opium is of value not only in relieving the pain and the great nervous prostration so constant in disease of the abdomen, but it also facilitates the recovery of the injured structure. Warmth, anodyne applications, the use of local depletion may be used; purgatives should be avoided, and rest strictly maintained; but mercurial preparations, given so as to affect the mouth, are as injurious in this form of strumous complication as in any other, and it is not necessary to recur to mercury for an aperient remedy, nor to prevent the opiates from checking secretion.

In the more chronic forms of the disease, the means best calculated to remove the local malady are those suited for the

removal of the general state ; such as nourishment as far as can be borne, cod-liver oil, steel as the iodide, steel wine, the iodide of potassium, alkalies, &c. Occasional counter irritant may be used, and moderate pressure on the abdomen employed to promote the absorption of serous effusions ; an elastic bandage strips of plaster, as the adhesive or the belladonna plaster, may be thus applied ; in some instances in which I have used the ammoniacum plaster with mercury, the intolerable itching which was produced compelled the removal of the application. Residence at the sea-side greatly facilitates recovery in these cases. In slow strumous effusion, especially in young persons, after peritonitis, it is often extremely difficult to produce absorption, and paracentesis is sometimes advisable.

Peritonitis with *cancerous* disease is always associated with enfeebled power and diminished functional activity. Remedies such as diuretics have very little effect in promoting the absorption of fluid, and any measures which still further diminish strength appear to increase dropsical effusion. To sustain the powers of life by every available means is the best preventive against this result. If acute symptoms supervene, the opiate plan of treatment must be followed with rest and bland nutritious diet. If paracentesis be performed, temporary relief may be obtained ; but more frequently the patient very rapidly declines, and we then find that the whole of the diseased peritoneal surface has increased in vascularity, and lymph is poured out.

Peritonitis with cirrhosis is generally found in persons who have been of intemperate habits ; the arteries are often diseased, and the kidneys may be granular and atrophied. At an early stage of the disease, when the diet can be regulated, and the excretory functions of the liver, the kidneys and the skin stimulated to increased action, the symptoms may, in a great degree, be alleviated ; and when acute peritonitis is set up with cirrhosis, no class of cases are more benefited by the judicious use of the ordinary remedies for peritonitis, namely, local depletion, and mercurials with opium, on account of the stimulant effect which mercurials have on the excretory glands ; but all the good effect of mercury may be attained without this remedy being used so as to produce salivation.

If the peritonitis be of a chronic form, and associated with advanced cirrhosis, our measures will, at best, be only palliative. Some have recommended mild mercurial salivation before tapping, to prevent the supervention of acute symptoms; but we have no experience in such an application of this medicine, and believe, that if tapping be really necessary, mercurial salivation would be detrimental, and would increase the exhaustion which often follows the operation, or that the mercurial cachexia would lead to the speedy reaccumulation of the fluid. Mercurial frictions are less objectionable when used with moderation; and minute doses of blue pill, with tonics, as quinine, or with aperients, are, in many instances of chronic peritonitis from hepatic disease, of great service. Other remedies may also be tried, as diuretics, iodide of potassium, nitro-hydrochloric acid, &c., but the persistent congestion of the vena portæ interferes with their absorption and with their beneficial action. Nearly the same remarks apply to the treatment of peritonitis coming on in the course of chronic disease of the heart and of the lungs. In these cases I never recommend paracentesis, unless compelled by the urgent distress from enormous distension.

We believe that the benefit generally ascribed to mercury in the treatment of acute peritonitis is not an established fact, and the apparent good result of its use may, perhaps, be correctly attributed to the opium with which it is usually combined.

M. Beau has employed large doses of quinine in the treatment of acute peritonitis, but we have no experience in this use of quinine.

Numerous instances might be adduced to illustrate the symptoms, the pathological appearances, and the treatment of acute as well as chronic peritonitis; the fearful suddenness, and oftentimes fatal result of the former, and the insidious character of the latter, might be abundantly demonstrated; each chapter has contained some illustrations of peritoneal disease, but a few cases in which some peculiarities were observed, are all that we shall mention.

CASE CCXXIX.—*Peritonitis. Local Suppuration. Perforation of the Diaphragm.*—Robert P—, æt. 26, was admitted into Guy's Hospital, January

6th, 1858; he had been a labouring man, residing in Bermondsey; and stated that he had been well till seven weeks previously, when one night he was suddenly seized with violent pain in the abdomen, accompanied by vomiting and purging; he was, however, able to walk to an apothecary's house on the next day, but the symptoms remained, and afterwards increased in severity. He was admitted exceedingly ill, and there was much obscurity as to the precise character of the disease; he had the appearance of a patient in the later stage of typhoid fever; the diagnosis was peritoneal disease; and he died on the 9th.

The body was wasted, the abdomen tumid. The brain was not examined. The left lung was pushed up, and its base was adherent to the diaphragm; when the lung was removed, its lower surface was found to form part of an abscess situated in the abdomen, and which had perforated the diaphragm. The abscess was in the left hypochondriac region; and the surface of the spleen, the left lobe of the liver, and the diaphragm formed its walls; the surface of the lung opposed to the opening of the diaphragm was also involved. The fluid in the abscess was well-formed pus, without odour. The neighbouring organs were carefully examined, to see if any disease in them had produced the abscess, but nothing could be satisfactorily found. The stomach and its mucous membrane were healthy. The duodenum and colon were firmly united, and gave way on attempting to separate them. All the intestines were more or less united, but there was no ulceration. The kidneys were coarse; the heart and bronchial glands were healthy. We believe that the local peritoneal disease in this instance arose from some blow which the patient had received, for there was no evidence of perforation of any of the viscera.

CASE CCXXX.—*Chronic Peritonitis. Chronic Pneumonia. Struma. No Pain. Great Tympanitis.*—William F—, æt. 46, was admitted into Guy's, February 13th, 1857, and died March 26th, 1857.

He had been for several years a policeman in the Borough, and was of temperate, steady habits. He had not received any blow on the abdomen, nor was he aware of any exciting cause of the disease; he had had, however, diarrhoea some months previously. Two weeks before admission the abdomen became enlarged and tense, but without any pain; this increase of size went on till admission, when the abdomen was distended and tympanitic, but still free from pain.

He was a tall man, spare, and emaciated, the eyes glazy and sunken, the pupils contracted, and the conjunctiva injected, the pulse very small, quick,

and compressible, the tongue clean, the bowels open regularly, the urine scanty. In the abdomen slight fulness could be detected in the region of the liver; fluctuation was indistinct. The abdomen was neither hot nor tender on pressure. He was prostrate and sunken in the bed; but he could take food well, and was quite sensible. At the apices of the chest the respiration was very coarse, almost bronchial; the heart sounds were normal.

He was ordered iodide of potassium gr. ij, solution of potash \mathfrak{xxv} , tincture of henbane, \mathfrak{xxv} , in mint julap. The tympanitis in great measure subsided about a fortnight before death, but he became gradually more prostrate; still, he did not suffer from any pain; three days before death his mind wandered considerably, and only eight to ten weeks intervened between the commencement of the symptoms and the fatal issue.

Inspection about twelve hours after death. *Abdomen*.—The intestines were moderately distended; the peritoneum contained about two quarts of bloody serum; the intestines were everywhere covered over with lymph in grains somewhat resembling tubercles; there were some tolerably firm bands of adhesions, and the small intestines were united together, but not firmly. The investment of the spleen was more than a quarter of an inch in thickness, and of a whitish colour from lymph, and contained yellow opaque grains or cheesy masses from degenerating tissue. The appendix was coiled like the letter S, on the right side of the cæcum, and was adherent. The liver was fatty; the spleen semi-diffuent; the kidneys somewhat granular and atrophied. The mesenteric glands were healthy. The stomach was contracted and small, and presented scarcely any post mortem solution. In the ileum were several small passive ulcers, extending through the mucous membrane, but without any injection around them. The cæcum and colon were healthy. The appendix contained fæces surrounded by thickened mucus. *Chest*.—The pleura was adherent at both apices, but especially on the left side; the left apex was puckered, and was of an iron gray colour, consolidated, and contained some calcareous masses; some small granular masses resembling tubercles were observed. At the base of the right lobe were several lobules consolidated; and some were in a state of softening, or gray hepatization; they were situated beneath the pleura as in pyæmia. The heart was healthy, except slight atheroma of the valves. The tubercles in the peritoneum consisted of elongated fibre-cells, with a nucleus rendered very distinct by acetic acid; in some parts more rounded nucleated cells were closely aggregated, in others fibres were formed; some portions showed degenerating tissue. The structure was more highly organized than in struma, but it could not be considered cancerous.

Many facts of great interest were presented in this case. 1st. The insidious character of the peritonitis, unaccompanied by pain, but marked by tympanitis, with gradually increasing prostration. 2nd. The blood effused into the serous membrane, which apparently arose from the rupture of newly-formed capillaries. 3rd. The apparent tendency to struma, as shown by

the condition of the lungs, presenting a retrocedent state chronic pneumonia. 4th. The condition of the blood, as indicated by the pyæmic lobular pneumonia, the diffluent state of the spleen, and the effusion of blood into the peritoneum. As to the exciting or predisposing causes of disease, nothing could be found; he had not been more exposed than policemen generally are to cold or to wet; he had been a man of temperate habits, and there was no disease of the liver beyond fatty degeneration. The probable predisposing cause was the strumous diathesis. We have already referred to this case as one which might easily have been mistaken at an early stage for one of functional dyspepsia;* and other cases of a similar kind have come under our notice.

CASE CCXXXI.—*Ulceration of the Intestine. Strumous Peritonitis. Fæcal Abscess. Umbilical Discharge.*—Mary Ann E—, æt. 38, by occupation servant, was admitted under Dr. Hughes' care, into Guy's Hospital, June 18th, 1857, and died on August 4th. She had resided at Brixton, and two months previously was quite well, when she began to suffer from pain in the abdomen; this gradually increased with vomiting and distension, and at last several tumours appeared. On admission she was very ill, with symptoms of general peritonitis, tenderness, distension, &c. Three or more defined tumours could be felt; one at the umbilicus, a second in the right hypochondrium, and a third on the left side. They could be reduced by pressure, and fluctuation was believed to be felt. On July 30th, the one near the umbilicus burst at that part, and pus having a fæcal odour was discharged. She died on August 4th. The body was wasted. The thoracic viscera were healthy. There was a large fæcal abscess below the umbilicus, circumscribed by the liver and the colon, and extending into the transverse colon; upon attempting to separate the intestine, further openings were made. Numerous ulcers were found in the ileum, cæcum, and colon; as most of these ulcers had perforated the coats of the intestine; extravasation had, however, been prevented by adhesions. Between the coils of intestine close to the mesentery, were several loose earthy bodies, and similar ones were found in the mesenteric glands. The liver was fatty.

Although there was no evidence of strumous disease in the lungs of this patient, we regarded the case as one of strumous disease of the abdomen, from the condition of the mesenteric glands and the general character of the disease. It is probable that ulceration of the mucous membrane of the ileum first took place

* See p. 201.

and afterwards chronic peritonitis; the latter was marked by severe attacks of pain and vomiting; adhesions between the coils of the intestine then formed, and limited the extravasation which subsequently took place from the perforated intestine.

In strumous peritonitis, suppuration sometimes arises from the degeneration of the low organized peritoneal product, and the pus forms an opening into the intestine, namely, from the peritoneal to the mucous surfaces. In other cases, the perforation is from within; the extensive ulceration of the small and large intestine in this case rendered it probable that the latter was the course of the disease. The external opening at the umbilicus is a rare event in these cases, and if the faecal abscess thus opening had been the only one, great relief to the pain and distress might have followed; but these collections of faecal matter were numerous. The calcareous condition of some of the mesenteric glands, and loose calcareous bodies of similar kind in their neighbourhood, are interesting facts connected with the explanation of loose bodies sometimes found in the serous sac.

CASE CCXXXII.—*Peritoneal Ascites. Cancer of the Ovaries and Peritonæum. Paracenteris. Peritonitis.*—Ann H—, æt. 45, was admitted under my care, September 20th, 1858. She was a single woman, and had ceased to menstruate in December, 1857. Since that time the abdomen had begun to swell, and ascites came on; she was tapped, and dark coloured fluid was drawn off; vomiting with severe pain in the abdomen supervened, and continued till her admission.

On the 20th. There was great emaciation; the irritability of the stomach was rather less, but still very troublesome. Soda-water, creasote, hydrocyanic acid, bismuth, had been tried, with very little effect. The abdomen again became very large, and fluctuation was distinct; some sleep was obtained by the daily administration of opium.

On October 6th, she was very prostrate, and suffered much from the distension of the abdomen, and from flatulence, &c. The countenance was haggard and wasted; she was partially relieved by turpentine injection.

On the 7th, she had very severe pain in the abdomen, which was much distended and tympanitic; fluctuation could be felt at the most depending part. She gradually sank.

On inspection, the peritonæum was found studded with cancerous tubercles; the omentum was contracted, and contained much deposit; the stomach was irregularly contracted into two pouches; the pylorus was healthy; but an inch from the pylorus its coats were invaded by cancerous disease; the mucous membrane in several parts was intensely congested. The intestines were distended with flatus, and there were indications of peritonitis at the

lines of contact of the coils of intestine; some serous effusion was observed. The tubercular deposit on the peritoneum consisted of abundant nuclei. There was a medullary tubercle in the kidney; the glands were otherwise healthy; but there were no tubercles in the liver. The pleura presented numerous tubercles; and there was a considerable quantity of serous effusion in both serous cavities. The heart was normal. The spleen was enlarged. Both ovaries with the uterus were united into one mass by deposit of a fibro-medullary character; and numerous cysts were observed containing secondary cysts. The os uteri was healthy.

The acute peritoneal mischief was set up by the paracentesis constant pain, and almost daily vomiting, with gradually increasing exhaustion followed, till death ensued. The onset of the disease was insidious, and probably arose in the ovaries. Cancerous tubercles were then developed on the peritoneum, and an effusion took place. This effusion may be without pain, excepting that arising from distension; at length the large size of the abdomen, the painful distension, the inability to take food from the pressure on the stomach, the impeded respiration, the sleepless nights, and the general distress, cause the patient to long for relief; but there is always the danger of more acute inflammatory disease being set up, as was the case in this patient before she applied for admission into Guy's Hospital.

CASE CCXXXIII.—*Carcinoma of the Peritoneum, with Effusion. Paracentesis.*—Mary T—, æt. 57, admitted into Guy's Hospital April 25th, 1857 and died on August 28th.

She stated that about two months before admission, the abdomen began to swell, and gradually increased. On June 13th, paracentesis abdominis was performed, and again on July 9th; at each operation more than five gallons of fluid being drawn off. No signs of peritonitis followed, but she gradually sank.

On inspection, the lungs and pleura were found to be healthy; there was recent lymph effused on the anterior aspect of the pericardium, upon the ventricles; and a small quantity posteriorly. In the abdomen there were several pints of fluid; numerous white cancerous nodules were scattered over the serous membrane, especially on the intestines. The membrane also was thickened by chronic action, in a more marked degree on the anterior parietes. The peritoneum covering the liver was opaque, and in some places tubercular. The omentum was drawn up and formed a solid mass (cancerous). The lymphatic glands were only slightly affected, but there were some large masses between the diaphragm and the liver. The surface of the diaphragm itself was affected. The tunic of the spleen opaque. The liver, spleen, supra-renal capsules, and kidneys, were healthy. The right ovary consisted of a mass of new growths and cysts, about the size of the

fist; the former were made up of a spongy-looking fungus, composed of fibro-cellular structure, and the latter were filled with fluid. There were some smooth cavities containing a soft sebaceous matter, and amongst these were numerous hairs.

In this case the disease also probably commenced in the ovaries; there was absence of pain, but gradual effusion; the ascites was, however, greatly relieved by paracentesis, no acute peritonitis followed, but as too frequently happens, the fluid soon re-collected, and again required to be drawn off; but at each time the patient was less able to rally from the effect of the operation.

CASE CXXXIV.—*Chronic Peritonitis. Renal Disease. Spurious Cysts in the Peritoneum.*—Elizabeth S—, æt. 44, a married woman, was admitted, December 7, 1859, under Dr. Barlow's care, and died February 11th, 1860. She had been confined eight months before admission; and swelling of the legs was followed by effusion into the abdomen. The urine was albuminuric. The abdomen became distended; cough distressed her; prostration supervened; and death followed.

Inspection.—Stout person, affected with general anasarca. There was effusion into the left pleura; and the left lung on that side was much compressed. There was evidence of bronchitis, and the bronchial tubes were much congested. *Heart.*—The muscular fibre pale; the right and left ventricles were both much dilated. *Abdomen.*—The intestines distended; on opening the serous membrane an appearance very closely resembling ovarian dropsy presented itself. The supposed cyst on the right side consisted of peritoneal effusion, shut off by fibrinous adhesions. This serous collection extended from the right iliac region as high as the liver, and between the liver and diaphragm; the fibrous cyst could in great measure be separated from the peritoneum, and contained many pints of slightly milky serum. Another cyst was found between the colon and the stomach. The mesentery was slightly contracted, and all the intestines were matted together. The kidneys contained cysts and some deposit. The liver was soft and congested. The ovaries were atrophied.

Effusion into the peritoneum; and serous inflammation is of frequent occurrence in renal disease, and such was the cause of the ascites in this case. It is, however, especially recorded, on account of the very peculiar appearance that was presented on opening the abdomen; there was the precise resemblance of a patient affected with ovarian dropsy, for the lymph effused upon the peritoneum had formed a cyst, and shut off the serous exudation. The lymph with its contained serum could be separated

from the liver and intestine, but the ovaries themselves were healthy.

CASE CCXXXV.—*Colloid Cancer of the Peritoneum and Ascending Colon*—Harriet K—, æt. 32, a hawker residing in Kent Street, was admitted in Guy's Hospital March 27th, 1861, under the care of Dr. Rees. Her general health was good, and her habits of life temperate; but she had an anæmic appearance. Her employment exposed her to wet, cold, and fatigue and she had evidently suffered from causes of depression, having had no proper food, and the care of a family of five children. There was no history of hereditary disease.

Six months previous to her admission, she was seized with a sharp pain in the lower part of the abdomen, and the pain was gradually increased in pressure; from that time the bowels became relaxed, but the diarrhoea had been more severe for three months, ten to twelve motions being passed; the motions were of a dark colour, and sometimes were of white glairy matter, like white of egg. Small masses of indurated feces were also sometimes passed. She complained of a sense as of great weight at the lower part of the abdomen when standing. For a short time she suffered from dyspnoea; but the cough was slight; the expectoration was greenish and tenacious. There had been no catamenial discharge for eighteen months. She suffered from vomiting; the pulse, 120, was small and soft; the tongue was dry in the centre, but the tip and the edges were clean and moist. The diarrhoea continued after admission, but it was partially checked by treatment; she sank a week after admission.

Inspection was made twenty-four hours after death. The thoracic viscera were natural. The whole of the peritoneum was covered with colloid cancer which extended from the diaphragm to the pelvis; no viscera, however, were involved except the ascending colon, and it was doubtful whether this was primary or secondary disease. On opening the abdomen, the omentum was seen spread over the surface of the intestine, and reached nearly to the pubes. The mesentery was converted into a solid mass of colloid, an inch in thickness, and of considerable density. It formed a tolerably uniform mass except at the edges, where the disease existed in the form of distinct nodules; the omentum was slightly adherent. On raising the omentum, adhesion existed with the ascending colon, and on opening this portion of the intestine the walls were found to be involved, and in part destroyed by the disease. The cæcum itself was free, but above it the walls of the colon were converted into a solid tumour for about five inches. In other parts the disease was confined to the peritoneum, but in the ascending colon it had penetrated the coats and had partially destroyed them; thus, on section, the muscular coat could be seen traversing the centre of the colloid growth. The mesentery and its glands and the lumbar glands were free from disease; but the whole of the diaphragmatic peritoneum was covered, so also the peritoneal surface of the bladder, liver, and spleen. The substance of the liver and spleen were healthy, so also the kidneys, uterus, and ovaries. The left iliac vein was distended with an ante-mortem clot. The microscope showed colloid structure in the peritoneal growth.

This was a well-marked instance of colloid disease of the peritoneum originating in similar disease of the colon. A glairy discharge from the bowels was present, but I have not observed this symptom in other instances of colloid disease which have come under my notice. Pain in the region of the colon was the first indication of disease, and was followed by diarrhoea of a very persistent character, by vomiting, and gradual prostration. The patient was only thirty-two years of age, but she had had five children. There was nothing, however, during the course of the complaint to enable us to diagnose its precise character.

LOOSE BODIES are occasionally found in the peritoneal cavity, oval in form, and in size from a bean to a pigeon's egg. They do not appear to produce any symptom, and they have been known to pass into hernial sacs. They are variously constituted. 1st. Some arise from growths connected with the intestine, and, whilst some may be quite free, others are found pendant by long thread-like bands. 2nd. The separation of appendicæ epiploicæ apparently constitutes others, as in a case recorded by Mr. Shaw, and as shown by the arrangement of its layers. 3rd. The separation of calcareous and degenerate glands may give rise also to these loose bodies. This mode of formation appeared probable in Case CCXXXI. 4th. The coagulation of fibrinous masses or of blood leads perhaps to their formation; but this source is of very doubtful occurrence, although in spurious melanosis dark portions of effused blood may seem to be in an almost separated condition.


In the several chapters of this work allusions have been made to growths arising from morbid processes connected with the intestine, constituting *tumours*; and we are guided to a correct appreciation of their character partly by their position, but especially by the disturbance of the function of that part of the intestine which is diseased, whether it be the stomach, duodenum, jejunum, ileum, cæcum, or colon. So also enlargements of the liver afford diagnostic signs of hepatic disturbance, and a tumour connected with the liver is modified by respiratory movements; the dulness on percussion is greater than that produced by growths connected with the intestine, unless, as in rare cases, the colon overlies the margin of the liver. In pancreatic tumours diseases of the stomach are more closely simulated; in aneur-

ismal tumours the expansive thrill is more uniform than enlarged glands pressing upon the aorta, and the pain is of peculiarly severe in aneurism. In enlargement of the spleen the growth is situated deeply in the left hypochondriac region it is uniformly dull, and generally presents on its anterior margin a deep notch; we have, however, known this notch be compounded with cancerous enlargement of the lumbar gland and even with the solid portion of a large ovarian cyst; all diseases of the spleen are accompanied with anæmia of a peculiar character.

The glands about the kidney, and even a large kidney itself or a moveable one, may obscure the diagnosis of omentum, mesenteric, and intestinal tumour; the renal disease is situated more deeply in the loin than is found to be the case with tumours of the ascending or descending colon; the former has deeper dullness on percussion, and as enlargement takes place anteriorly the growth extends on the side of the spine, and is felt near the lateral part of the umbilical region. The state of the urine also affords important diagnostic indications. Omentum and mesenteric growths are also more free in their movements, and when the patient inclines forwards on the hands and knees, may often be felt against the abdominal parietes. Ovarian tumours are felt at their early stages in the iliac fossa; but the evidence of cæcal disease and of disease of the sigmoid flexure are wanting, and vaginal examination in many cases clears up the diagnosis; still there is sometimes difficulty in distinguishing great ascites with contraction of the mesentery from very large ovarian cysts or form tumours. Again, spasmodic contraction of portions of the straight and of the oblique muscles of the abdominal parietes, we find in states of spinal irritation, adds another fallacy to the diagnosis of abdominal diseases; they constitute the so-called "phantom tumours," but after repeated examinations they will be found to change their position, and sometimes they will entirely disappear under the hand. Although in most instances morbid growths affecting the abdominal glands and the several portions of the alimentary canal may be readily distinguished the one from the other, still there are cases of peculiar difficulty and complexity; every instance must be individually considered in all its relations, and careful and accurate

observation will, in most cases, unravel the true character of the disease.

In each Section of our Work we have sought to show the leading characteristics of diseased conditions as manifested in the various portions of the alimentary canal; and to do this have given the instances themselves, as facts upon which each one may form their own opinion, rather than depend entirely upon the deductions we have drawn from them. Such general conclusions in most chapters have preceded the cases upon which they are founded; and we leave them before our readers with the hope that they will serve further to elucidate the general symptoms, pathology, and treatment of diseases of the alimentary canal.



INDEX TO THE ILLUSTRATIVE CASES.

CASE

- I.—Diffused inflammation of the throat.
- II.—Diffused inflammation of the throat. Ulceration of the pharynx.
- III.—Hydrophobia.
- IV.—Dysphagia. Mania.
- V.—Poisoning by sulphuric acid—death on the 11th day.
- VI.—Poisoning by soap lees.
- VII.—Poisoning by sulphuric acid.
- VIII.—Poisoning by nitric acid—secondary effects.
- IX.—Poisoning by nitric acid—recovery from the primary effects.
- X.—Diseased cartilages of the trachea. Ulceration of the œsophagus.
- XI.—Ulceration of the œsophagus. Perforation of the trachea.
- XII.—Cancer of the œsophagus. Sloughing pneumonia; the pneumogastric nerve involved.
- XIII.—Epithelial cancer of the œsophagus, communicating with the trachea. Pneumonia. Granular kidneys.
- XIV.—Cancer of the œsophagus, of the cervical glands, and of the thyroid body. Gangrene of the lung.
- XV.—Epithelial cancer of the œsophagus, pancreas, liver, and stomach. The pneumogastric nerves involved. Granular kidneys. Chronic pleuro-pneumonia, with cancer. Fibrous tumour in the uterus. Cancer of the supra-renal capsules, and semilunar ganglion.
- XVI.—Epithelial cancerous tumour in the pharynx, closing the entrance into the œsophagus. Effusion of false membrane in the larynx and trachea. Acute bronchitis.
- XVII.—Carcinoma of the œsophagus, communicating with the trachea. Cancer of the lung and of the kidney.
- XVIII.—Cancer of the œsophagus—death from exhaustion.
- XIX.—Cancer of the œsophagus. Gangrene of the lung.
- XX.—Cancer of the œsophagus, with pleurisy and contracted kidneys.
- XXI.—Cancer of the œsophagus. Gangrene of the lung. Communication with the trachea.
- XXII.—Cancer of the œsophagus. Extension into the lung. Gangrene.

CASE

- XXIII.—Carcinoma of the œsophagus. Pleuro-pneumonia from direct extension of disease.
- XXIV.—Cancer of the œsophagus. Pneumonia. The pneumogastric nerve involved.
- XXV.—Cancer of the œsophagus. Pneumogastric nerves involved. Pneumonia.
- XXVI.—Carcinoma of the throat. Strumous pneumonia.
- XXVII.—Cancer of the œsophagus. The left pneumogastric involved. Pneumonia.
- XXVIII.—Cancer of the œsophagus. Communication with left bronchus. The pneumogastric involved. Old vomica in the lung. Extension of disease through the diaphragm.
- XXIX.—Cancer of the œsophagus. Pneumogastric nerves truncated. Sloughing extending into the lung and through the diaphragm.
- XXX.—Cancer of the œsophagus. Chronic pneumonia. Vomica. Acute pneumonia.
- XXXI.—Cancer of the œsophagus. Artificial opening made into the stomach.
- XXXII.—Cancer of the œsophagus. Sloughing. Perforation of the aorta. Sudden and fatal hæmorrhage.
- XXXIII.—Aneurism of the aorta, and sloughing of the œsophagus.
- XXXIV.—Aneurism of the ascending aorta, rupturing into the pericardium. Communication of the œsophagus with the left bronchus.
- XXXV.—Aneurism. Pressure on the œsophagus and on the left bronchus. Difficulty in deglutition. Sudden death.
- XXXVI.—Rupture of the œsophagus.
- XXXVII.—Strumous disease of the third rib. Phthisis. Ulceration of the intestine. Strumous disease of the kidney, ureter, bladder, prostate gland, vesiculæ seminales, spermatic cord and testis. Fatty degeneration of the stomach and liver, &c.
- XXXVIII.—Thinning of the mucous membrane, with the appearance of minute cysts (probably post-mortem) from the stomach of a man who died five weeks after taking a strong solution of chloride of zinc.
- XXXIX.—Poisoning by brandy.
- XL.—Poisoning by sulphuric acid—death on the eleventh day. Destruction of the mucous membrane of the lesser curvature and pyloric extremity of the stomach. Acute inflammation of the colon and small intestines.
- XLI.—Poisoning by chloride of zinc—Burnett's disinfecting fluid. Death in fourteen weeks.
- XLII.—Poisoning by corrosive sublimate—death on the sixth day.

CASE

- XLIII.—Bronzed discoloration of the skin. Violent vomiting. Excessive prostration.
- XLIV.—Syphilis. Diphtheritic inflammation of the stomach. Diseased kidneys. Necrosis of the bones of the nose.
- XLV.—Suppuration in the coats of the stomach.
- XLVI.—Superficial ulceration of the stomach. Diseased supra-renal capsules.
- XLVII.—Superficial ulceration of the stomach and duodenum. Phthisis. Ulceration of the ileum, cæcum, colon, and rectum. Great intemperance.
- XLVIII.—Chorea. Vegetations on the mitral. Ulceration of the stomach.
- XLIX.—Chronic catarrh, and superficial ulceration of the stomach with intense congestion. Disease of the heart. Small degenerated kidneys,
- L.—Catarrh and superficial ulceration of the stomach; cystic disease of the ovary.
- LI.—Follicular ulceration of the mucous membrane of the stomach with renal anasarca, and diseased heart.
- LII.—Follicular inflammation of the stomach. Burn on the leg. Amputation. Abscess in the lungs and spleen.
- LIII.—Chronic ulcer of the stomach. Phthisis. Branches of the pneumogastric nerve involved.
- LIV.—Chronic ulceration of the stomach, involving the pneumogastric nerve. Atrophy of the left lobe of the liver. Death from exhaustion.
- LV.—Chronic ulceration of the stomach, with painters' colic. Perforation.
- LVI.—Perforating ulcer of the stomach, with a second small chronic ulcer in the same organ.
- LVII.—Chronic perforating ulcer of the stomach.
- LVIII.—Chronic ulceration. Death from perforation.
- LIX.—Chronic ulcer of the stomach. Perforation.
- LX.—Perforating ulcer of the stomach.
- LXI.—Chronic ulcer of the stomach. Perforation, excepting through the peritoneum. Fatal peritonitis.
- LXII.—Perforation of the stomach. Local suppuration in the peritoneum. Pleuro-pneumonia.
- LXIII.—Chronic ulceration of the stomach, extending to the diaphragm, and simulating pneumothorax.
- LXIV.—Fæcal abscess connected with the stomach, the lung, the spleen, and the transverse colon.
- LXV.—Chronic ulcer of the stomach. Peritoneal abscess. Perforation of the diaphragm. Empyema.

CASE

- LXVI.—Chronic ulceration of the stomach. Perforation. A sinus extending into the left lung. Gangrene. Empyema.
- LXVII.—Chronic ulceration of the stomach. Fatal hæmorrhage. Perforation of the splenic, and of the pancreatic arteries.
- LXVIII.—Ulceration of the stomach. Fatal hæmorrhage.
- LXIX.—Chronic bronchitis. Emphysema. Chronic ulcer of the stomach.
- LXX.—Chronic ulceration with villous growth. Stomach exceedingly contracted. Simulating cancer.
- LXXI.—Chronic ulcer of the stomach relieved.
- LXXII.—Ulceration (cancerous?) of the stomach. Relieved.
- LXXIII.—Ulceration of the stomach. Sloughing. Paraplegia. Softening of the spinal cord. Disease of the vertebrae.
- LXXIV.—Mottled kidney. Anasarca. Pneumonia. Sloughing of the mucous membrane of the stomach.
- LXXV.—Thickened pylorus. Cicatrix of the mucous membrane. Hypertrophy. Ulceration of the cæcum and colon. Fatty degeneration of the heart.
- LXXVI.—Diseased pylorus. Phthisis.
- LXXVII.—Scirrhus pylorus. Carcinomatous tubercles in the liver, spleen, and kidney, and on the diaphragm.
- LXXVIII.—Medullary cancer of the stomach, having a villous character.
- LXXIX.—Villous cancer of the stomach? Cirrhosis. Ascites.
- LXXX.—Colloid cancer of the stomach, and of the colon.
- LXXXI.—Colloid cancer of the stomach, omentum, peritoneum, and rectum.
- LXXXII.—Medullary cancer of the stomach, of the liver, and of the lungs.
- LXXXIII.—Chronic ulceration of the stomach. Cancer.
- LXXXIV.—Cancer of the stomach; communication with the colon. Ulceration of the cæcum and ileum. Chronic phthisis.
- LXXXV.—Cancer of the stomach.
- LXXXVI.—Struma. Cancer of the stomach.
- LXXXVII.—Cancer of the stomach. Disease of the supra-renal capsule.
- LXXXVIII.—Cancer at the pylorus. Simulating disease of the œsophagus.
- LXXXIX.—Cancer of the pylorus. Hydatid disease of the cellular tissue of the bladder.
- XC.—Pyrosis.
- XCI.—Dyspepsia. Pyrosis.
- XCII.—Infantile dyspepsia. Sudden collapse.

CASE

- XCIII.—Hæmatemesis from cancer of the liver.
- XCIV.—Hæmatemesis from portal congestion.
- XCV.—Hæmatemesis after great intemperance.
- XCVI.—Hæmatemesis from ulceration of the stomach.
- XCVII.—Hæmatemesis from ulceration of the stomach.
- XCVIII.—Hæmatemesis. Vicarious menstruation. Aggravated hysteria, simulating fever.
- XCIX.—Hæmatemesis. Vicarious menstruation.
- C.—Hæmatemesis. Typhus fever.
- CI.—Acute jaundice. Violent vomiting. Recovery.
- CII.—Old peritoneal adhesions. Twist of the sigmoid flexure. Acute peritonitis. Death on the sixth day.
- CIII.—Violent vomiting. Gall-stone. Pleuro-pneumonia.
- CIV.—Anæmia. Struma. Vomiting and diarrhœa every other day.
- CV.—Chlorosis. Vomiting. Tubercles in the brain. Fatty degeneration of the heart.
- CVI.—Vomiting—cerebral? Diseased ear.
- CVII.—Inflammation of the bronchi, and of the bile ducts—biliary hepatitis. Acute congestion of the duodenum.
- CVIII.—Ulceration of the duodenum. Perforation.
- CIX.—Chronic ulcer in the duodenum. Carcinoma of the liver. Jaundice. Granular kidneys. Obliteration of the bile duct.
- CX.—Strumous disease of the peritoneum. Perforation of the duodenum, and of the cæcum.
- CXI.—Cancer of the duodenum.
- CXII.—Chronic peritonitis. Acute peritonitis. Tubercular deposit on the serous membrane, and in the glands. Constriction of the duodenum, and great dilatation of its first portion. Small ulcer in the duodenum. Large chronic ulcer in the colon.
- CXIII.—Obstruction from biliary calculus in the upper part of the jejunum, thirty inches from the pylorus.
- CXIV.—Disease of the pancreas. Suppuration and gangrene. Pressure on the duodenum.
- CXV.—Perforation of the duodenum, by solution of the gastric juice.
- CXVI.—Gastro-enteritis—recovery.
- CXVII.—Acute enteritis.
- CXVIII.—Enteritis simulating mechanical obstruction.
- CXIX.—Sloughing ileum. Peritonitis. Large and fatty kidneys. Degeneration of the left lobe of the liver. Lobular pneumonia. Small and fatty heart.
- CXX.—Slight strumous disease of the mesenteric glands. Diarrhœa. Pneumonia.

CASE

- CXXI.—Strumous peritonitis.
- CXXII.—Strumous peritonitis. Fæcal abscess. Artificial anus.
- CXXIII.—Strumous disease of the mesenteric glands. Obstruction of the lacteals. Ulceration of the large and small intestine. Dysentery. Phthisis.
- CXXIV.—Ulcerated colon. Phthisis. No cough.
- CXXV.—Phthisis. Ulceration of the rectum and sigmoid flexure. Hæmorrhage from the bowels. Ulceration of appendix cæci.
- CXXVI.—Phthisis. Inflamed hæmorrhoids. Sciatica. Extensive ulceration of the larynx and intestine.
- CXXVII.—Unusually free cæcum.
- CXXVIII.—Unusually free cæcum. Fatty degeneration of organs.
- CXXIX.—Cæcum inverted. Phthisis. Local empyema. Large white kidneys.
- CXXX.—Intestinal obstruction of the ascending colon. Cæcum twisted to the left side into the left iliac and left hypochondriac regions. Death on the twentieth day.
- CXXXI.—Displaced cæcum. Perforation. Peritonitis. Paraplegia.
- CXXXII.—Twisted cæcum. Obstruction. Peritonitis.
- CXXXIII.—Cæcal distension and inflammation.
- CXXXIV.—Cæcitis.
- CXXXV.—Cæcal distension and inflammation.
- CXXXVI.—Cæcal distension and inflammation.
- CXXXVII.—Cæcal distension and inflammation.
- CXXXVIII.—Cæcal distension with inflammation.
- CXXXIX.—Cæcal inflammation, simulating hip-joint disease.
- CXL.—Cæcitis? Local peritonitis.
- CXLI.—Cæcal disease.
- CXLII.—Strumous inflammation of the cæcum.
- CXLIII.—Perforation of the cæcum. Abscess extending to the groins. Phthisis.
- CXLIV.—Bronchitis. Phthisis. Diseased appendix cæci.
- CXLV.—Phthisis. Ulceration of the larynx and of the ileum. Concretion in the appendix.
- CXLVI.—Pyæmia. Necrosed humerus. Cæcal disease.
- CXLVII.—Perforation of appendix cæci. Laminated concretions. Fatal peritonitis.
- CXLVIII.—Phthisis. Ulcerated appendix cæci opening into the ileum.
- CXLIX.—Disease of the cæcum, following a blow. Perforation of the appendix. Suppuration. General peritonitis. —Almost complete secondary perforation of the cæcum.

CASE

- CL.—Local peritonitis. Perforation of appendix cæci. Strangulation of the ileum by the appendix.
- CLI.—Ulceration of the appendix cæci. Perforation. Fæcal abscess. Secondary perforation of the ileum.
- CLII.—Fæcal concretion. Perforated appendix.
- CLIII.—Concretion in the appendix cæci. Perforation.
- CLIV.—Fæcal concretion in the appendix. Perforation. Peritonitis.
- CLV.—Ulceration of the appendix, with disease of the kidney.
- CLVI.—Cancer of the cæcum. Abscess in the groin.
- CLVII.—Carcinoma of the cæcum, of the omentum, and of the lumbar glands. Fæcal abscess.
- CLVIII.—Colloid cancer of the cæcum. Jaundice.
- CLIX.—Appendix cæci in the inguinal canal.
- CLX.—Appendix cæci adherent with the omentum at the internal abdominal ring. Supra-renal capsular disease.
- CLXI.—Inanition. Diarrhœa.
- CLXII.—Chronic diarrhœa. Hysteria. Great relief from tincture of iron.
- CLXIII.—Inflammation of the colon. Aphtha or muguet of the pharynx.
- CLXIV.—Diphtherite of the colon. Dysentery. Chorea.
- CLXV.—Inflammation of the colon. Hernia.
- CLXVI.—Dysentery. Ulceration of the small intestines. Perforation. Fæcal abscess. Peritonitis.
- CLXVII.—Ulceration of large intestine. Perforation. Submucous suppuration. Pus in the portal vein, and inflammatory patches in the liver.
- CLXVIII.—Dysentery. Perforation of colon.
- CLXIX.—Chronic bronchitic phthisis. Cirrhotic and lardaceous liver. Contracted abscess of liver. Chronic dysentery, and chronic peritonitis.
- CLXX.—Chronic dysentery. Hepatic abscess. Pyæmia. Abscess in the brain and lung.
- CLXXI.—Dysentery. Abscess of the liver. Perforation of the diaphragm. Empyema.
- CLXXII.—Cicatrization and contraction of the rectum, and sigmoid flexure after dysentery.
- CLXXIII.—Chronic ulceration of the intestine. Dysentery. Cicatrization. Contraction. Perforation. Abscess near the crest of the ilium.
- CLXXIV.—Dysentery. Pneumonia. Hydrencephaloid disease.
- CLXXV.—Diphtherite of the cæcum and colon. Bronchitis. Pneumonia. Cirrhosis.

CASE

- CCXX.—Suppuration external to the right kidney. Fibroid thickening of the tunic of the kidney. Chronic pyelitis. Obliteration of the vena cava. Adhesion. Thinning and doubtful perforation of the ascending colon.
- CCXXI.—Fæcal abscess in the pelvis, communicating with the ovary and bladder; opening twice into the rectum, and once on the groin.
- CCXXII.—Abscess external to the rectum, leading to perforation. Considerable fibrous thickening, and simulation of cancerous disease.
- CCXXIII.—Multilocular ovarian tumour. Perforation of cæca. Fæcal abscess. Pneumonia. Pus in the ovarian veins.
- CCXXIV.—Ovarian tumour filled with fæces, and opening into the ileum. Pneumonic phthisis.
- CCXXV.—Extra-uterine foetation opening into the sigmoid flexure.
- CCXXVI.—Acute and chronic peritonitis. Suppuration in the coats of the portal veins.
- CCXXVII.—Hypertrophy of the heart. Adherent pericarditis. Acute pericarditis. Pleurisy and peritonitis.
- CCXXVIII.—Acute peritonitis. Pericarditis. Pleuro-pneumonia. Small granular kidneys.
- CCXXIX.—Peritonitis. Local suppuration. Perforation of the diaphragm.
- CCXXX.—Chronic Peritonitis. Chronic pneumonia. Struma. Pain. Great tympanitis.
- CCXXXI.—Ulceration of the intestine. Strumous peritonitis. Fæcal abscess. Umbilical discharge.
- CCXXXII.—Peritoneal ascites. Cancer of the ovaries and peritoneum. Paracentesis. Peritonitis.
- CCXXXIII.—Carcinoma of the peritoneum with effusion. Paracentesis.
- CCXXXIV.—Chronic peritonitis. Renal disease. Spurious cysts of the peritoneum.
- CCXXXV.—Colloid cancer of the peritoneum, and ascending colon.

CASE

- CC.—Cancerous disease of the sigmoid flexure. Ecchymosis of the stomach. Ulceration of the ileum. Contracted mitral valve.
- CCI.—Cancer of the liver, of the lumbar glands, and of the sigmoid flexure.
- CCII.—Cancer of the sigmoid flexure. Constipation. Death on the twentieth day.
- CCIII.—Cancerous disease of the sigmoid flexure of the colon. Insuperable constipation. Death on the tenth day.
- CCIV.—Cancerous ulceration of the sigmoid flexure of the colon. Constipation.
- CCV.—Cancerous disease of the sigmoid flexure. Insuperable constipation.
- CCVI.—Cancerous disease of the sigmoid flexure. Constipation simulating hernia.
- CCVII.—Cancer of the sigmoid flexure. Obstruction relieved. Gradual exhaustion.
- CCVIII.—Colloid cancer of the sigmoid flexure. Artificial anus in the groin. Pleuro-pneumonia.
- CCIX.—Cancerous disease of the sigmoid flexure. Diarrhœa. Perforation. Fæcal abscess.
- CCX.—Cancerous disease of the transverse colon. Fæcal abscess.
- CCXI.—Carcinoma of the rectum, of the ovaries, and of the peritoneum. Acute peritonitis. Scirrhus.
- CCXII.—Epithelioma of rectum. Contraction and obstruction. Artificial anus in the loins. Diseased appendix cæci. Fæcal abscess.
- CCXIII.—Cancer of the jejunum, and of the mesenteric gland. Cancerous softening of the spinal cord. Paraplegia.
- CCXIV.—Cancerous ulcer of the colon; opening into the duodenum. Diarrhœa. Vomiting.
- CCXV.—Ulceration of the colon. Intestinal obstruction from contraction of the transverse colon. Cancer? Dysentery. Constipation. Diarrhœa.
- CCXVI.—Suppuration external to the sigmoid flexure of the colon. Opening on the abdominal parietes, and communicating with the intestine.
- CCXVII.—Abscess in the loins. Feculent smelling discharge. Pleuro-pneumonia with feculent smelling sputum. Recovery.
- CCXVIII.—Miscarriage. Pyæmia. Abscess between the uterus and bladder. Abscess in the loins opening into the ascending colon and into the iliac vein.
- CCXIX.—Abscess in the hypogastric abdominal parietes. Simulating ovarian disease. Recovery.

CASE

- CCXX.—Suppuration external to the right kidney. Fibroid thickening of the tunic of the kidney. Chronic pyelitis. Obliteration of the vena cava. Adhesion. Thinning and doubtful perforation of the ascending colon.
- CCXXI.—Fæcal abscess in the pelvis, communicating with the ovary and bladder; opening twice into the rectum, and also on the groin.
- CCXXII.—Abscess external to the rectum, leading to perforation. Considerable fibrous thickening, and simulation of cancerous disease.
- CCXXIII.—Multilocular ovarian tumour. Perforation of cæcum. Fæcal abscess. Pneumonia. Pus in the ovarian veins.
- CCXXIV.—Ovarian tumour filled with fæces, and opening into the ileum. Pneumonic phthisis.
- CCXXV.—Extra-uterine foetation opening into the sigmoid flexure.
- CCXXVI.—Acute and chronic peritonitis. Suppuration in the course of the portal veins.
- CCXXVII.—Hypertrophy of the heart. Adherent pericardium. Acute pericarditis. Pleurisy and peritonitis.
- CCXXVIII.—Acute peritonitis. Pericarditis. Pleuro-pneumonia. Small granular kidneys.
- CCXXIX.—Peritonitis. Local suppuration. Perforation of the diaphragm.
- CCXXX.—Chronic Peritonitis. Chronic pneumonia. Struma. No pain. Great tympanitis.
- CCXXXI.—Ulceration of the intestine. Strumous peritonitis. Fæcal abscess. Umbilical discharge.
- CCXXXII.—Peritoneal ascites. Cancer of the ovaries and peritoneum. Paracentesis. Peritonitis.
- CCXXXIII.—Carcinoma of the peritoneum with effusion. Paracentesis.
- CCXXXIV.—Chronic peritonitis. Renal disease. Spurious cysts in the peritoneum.
- CCXXXV.—Colloid cancer of the peritoneum, and ascending colon.

INDEX.

	PAGE		PAGE
ABERCROMBIE, spasmodic stricture of œsophagus . . .	19	BALDING, Dr., hæmatemesis in ulceration of œsophagus, case of . . .	37
„ inflammation of œsophagus . . .	23	Ballard, Dr., on pepsine . . .	190
„ ulceration of stomach . . .	128	Baly, Dr., on dysentery . . .	381
„ ileus . . .	460	Bands, intestinal, causing obstruction . . .	459, 465, 477
Abdominal walls, abnormal state causing constipation . . .	444	Barlow, Dr., ischuria renalis . . .	234, 279
„ suppuration in . . .	530	„ internal strangulation . . .	459, 463, 464
Abscess connected with œsophagus . . .	34	Barker, Dr., on dilatation of œsophagus . . .	79
(See Suppuration—Fæcal Abscess).		Bell, Sir C., œsophageal pouch . . .	79
Addison, Dr., supra-renal capsular disease . . .	104	Bernard, post-mortem solution . . .	92
„ absence of pain in pericarditis . . .	220	Bird, Dr. G., on chlorate of potash . . .	24
Albuminuria, dyspepsia in . . .	196, 198	Birkett, Mr. J., villous growth from colon . . .	470
„ as cause of peritonitis . . .	559, 564	„ hernia, with diseased rectum . . .	511
Alderson, Dr., cancer of œsophagus . . .	38	Bodies, loose, in peritoneum . . .	575
Anæmia, producing dyspepsia . . .	187	Boiling water, dysphagia from . . .	25
Andral, post-mortem solution . . .	91	Bougies œsophageal . . .	20, 44
Aneurism, sudden death—sympathetic nerve . . .	6	Brain, disease of, gastric solution . . .	95
„ producing dysphagia . . .	74, 77	„ cause of vomiting . . .	235, 237
„ sloughing œsophagus . . .	75	Brandy, poisoning by . . .	94
„ causing hæmatemesis . . .	208	Bright, Dr., fatty evacuations . . .	266
„ producing pain at stomach . . .	224	Brinton, Dr., on ulcer of stomach . . .	119, 123, 166
„ producing vomiting . . .	228	„ on duodenum . . .	244
Aorta injured by cancer of œsophagus . . .	38, 73, 74	„ on cæcal regurgitation . . .	316
Appendix cæci, diseases of . . .	313	„ on intestinal obstruction . . .	467
„ increase of length . . .	321	Bristowe, Dr., ulcerated œsophagus . . .	37
„ atrophy . . .	321	Bronchus, communication of, with cancer . . .	58
„ dilatation . . .	321	Brown, Dr. Blakeley, obstr. duodenum . . .	266
„ perforation of . . .	324, 345	Budd, Dr., hepatic abscess . . .	11, 386
„ concretions . . .	322	„ post-mortem solution . . .	91, 92
Ascaris lumbricoides . . .	522	„ dyspepsia . . .	193
„ vermicularis, or oxyuris . . .	524	„ gastric fermentation . . .	205
Ashton, Mr. on disease of rectum . . .	312	CANCER of œsophagus . . .	36, 45
Atrophy of mucous membrane . . .	84, 87, 304	„ of stomach . . .	162, 177—180

	PAGE		PAGE
Dysentery pus in vena portæ	405	Fæcal abscess from ovarian disease	540
" hepatic abscess 407, 409, 410	410	" from diseased rectum	542
" contraction of colon	412	" from strumous disease	543
" with cancer of colon	518	" and with umbilical discharge	301, 570
Dyspepsia, causes of	185	Fermentation of food	205
" from deficiency of gastric juice	186	Fever, hæmorrhage in	211
" from excess of gastric juice	192	Fistula, gastric	64, 73
" from morbid condition of gastric juice	193	Fletcher, Mr., œsophageal bougies	44
" from impeded movements	199	Fleming, Dr., local pharyngeal sup-puration	16
" from changed vascular supply	201	Foreign bodies in intestine	458
" from state of nervous system	201	Forster, Mr. J. C., opening stomach	64
" in phthisis	202	Fraser, Dr., green vomit	4, 209
" duodenal	248	Fright, producing dysphagia	19
Dysphagia, causes of	13	GAIRDNER, Dr., fæcal vomiting	165, 471
" from diphtheria	14	Galvanism, use of	132, 454
" from paralysis of soft palate	15	Gall-stones, causing obstruction	258, 261
" from suppuration in the pharynx	16	Ganglia, semi-lunar	40
" from fright	19	Gangrene of lung, with cancer of œsophagus	46, 48, 53, 54
" laryngeal	18	Garrod, Dr., dyspepsia	196
" from paralysis of pharynx	22	Gastric juice, deficiency	116
" in mania	22	" excess	192
" from poisons	25, 30	" irregular secretion of	193
" from ulceration of œsophagus	32	Gay, Mr. J., insuperable constipation	452
" from syphilitic ulceration	36	Glands, enlargement of, in cancer of œsophagus	38
" from cancer	36	" causing duodenal obstruction	258
" from polypus	74	Goodsir, Prof., sarcina ventriculi	126
" from aneurism 74, 75, 76, 77	77	Goodfellow, Dr., renal vomiting	198
" from pericarditis	78	Gorham, Mr., discharge of blood in intussusception	278, 464
EMPHYSEMA of mucous membrane of stomach	85	Gout, dyspepsia in	196
Empyema from ulcer of stomach perforating the diaphragm	140, 142	Gray, Mr. H., cancer of œsophagus	40
" from abscess of the liver	410	Graves, Dr., opium in perforated stomach	154
Enteritis	268, 272	" vomiting in disease of brain	235, 237
" simulating obstruction	273	Gull, Dr., dysphagia in diphtheria	14
" diagnosis of	278	HARLEY, Dr., intestinal concretions	457
" treatment of	281	Harty, Dr., on dysentery	391
FÆCES, character of	370	Hawkins, Mr. Cæsar, on opening colon	477
Fæcal abscess, with perforation of stomach	139	Headland, Dr., on action of medicines	7
" with dysentery	404	Hilton, Mr., on opening the colon	476
" with diseased cæcum 315, 352, 543	543	Hæmorrhage from cancer of œsophagus	37
" with cancer of cæcum	358	" from perforation of aorta	73
" with cancer of colon	511, 514	" from subclavian	38
" from suppuration in parietes of the abdomen	532, 536	" from œsophageal ecchymosis	81
		" from ulceration of stomach	120, 143, 144

	PAGE		PAGE
Hæmorrhage from stomach, causes of symptoms, treatment	206, 210, 212, 218	Larynx, phthisis	1
Holleston, Mr., diseased trachea . . .	33	Laryngeal spasm in aneurism . . .	7
Hour-glass contraction of stomach . .	121	Law, Dr., ulcer of stomach	11
Hunt, Mr., chlorate of potash	24	Lead colic	43
Hunter, John, post mortem solution	91, 92	Lees, Dr., on stomach	123, 12
Hutchinson, Mr. Jon., dyspepsia in phthisis	6	Leudet, M., on cæcal disease . . .	32
" internal strangulation	477	Liver, disease of, as cause of vomiting	23
Hydatid disease of bladder	182	Ludlow, Mr., œsophageal pouch . .	7
" causing obstruction	258, 266	Lung, injury of, from perforation of diaphragm	139, 142, 56
Hydrophobia, case of	21	" disease of, as cause of vomiting	24
Hypertrophy of mucous membrane of stomach	89		
ILEO-COLITIS	268	MAINGAULT on diphtheria	1
Inflation of intestine in intussusception	477	Mammillation of stomach	8
Inflammation of throat	17	Mania, causing dysphagia	2
" of œsophagus	23, 24	Marsh, Sir H., on regurgitative disease	202, 226, 24
" of stomach	93, 102, 103	" on jaundice	24
" of cæcum	318	Maunsell, on diseases of children .	23
" of colon	381	Mayo, Dr., spasmodic stricture of œsophagus	1
" of peritoneum	547	" cast of œsophagus	2
" strumous, of peritoneum	295, 561	" œsophageal pouches	7
Intestine, obstruction of	230, 459	" rupture of œsophagus	8
Intestinal bands	465	Mayne, Dr., on dysentery	38
" rotation	314, 465	Melena	36
Intussusception, symptoms, &c. . . .	468	Menstruation, vicarious	208, 210
" ileum and cæcum	492	Mesenteric glands, strumous diseases of	285, 288, 294
" into colon	492	Mineral acids, poisoning by . . .	9
" cæcum discharged	494	Mineral waters	19
" of ileum	497	Monro, Dr., disease of œsophagus .	31
" of colon	498	Moore, Mr. C. H., internal obstruction	46
JAUNDICE, vomiting in	229	Morehead on dysentery	38
Jenner, Dr., hyposulphite of soda . .	152, 206	Muco-enteritis	268, 275, 27
Jepson, Mr., diseased trachea	83	Murchison, Dr., fecal vomiting . .	122
Jones, Dr. Handfield, atrophy of stomach	83		165, 471
" pyrosis	194		
KENNEDY, Dr., cancer of stomach . .	164	OBSTRUCTION, duodenal	25
" hæmorrhage in typhus	211	" intestinal	45
Kidney, disease as cause of vomiting	234	Odling, Dr., analysis of food . .	20
Kingsford, Dr., dysphagia	14	Esophagus, diseases of	1
King, Mr. C., case of intussusception .	494	" spasmodic stricture	19, 2
King, Mr. Wilkinson, post-mortem solution	80, 91	" inflammation of	23, 2
LARYNX, disease of, causing dysphagia	18	" local suppuration	3
" diseased cartilages	18	" organic obstruction	2
		" ulceration of	32, 3
		" cancer of	36, 7
		" polypus	7
		" tumours—aneurism	7
		" pouches	7
		" warty growths	8
		" post-mortem solution	8
		" rupture of	8
		" regurgitation of food	23
		Omental perforation, causing internal strangulation	48

	PAGE		PAGE
Opening the stomach, operation of	36	Pharynx, connections of	13
the colon	477	„ local suppuration	16
Osborne, Dr., ulceration of sto-		„ ulceration of	17
mach	124, 222	„ cancer of	51
„ character of fæces	370	Philip, Dr. W., pneumogastric in	
Ovarian tumour, fæcal abscess	543, 544	digestion	6
Oxalic acid, poisoning by	94	„ on dyspepsia	200
		Phillips, Mr. B., intussusception	495
PAIN, absence of	3	Phthisis dysphagia	18
„ in inflammation of		with œsophageal cancer	42, 56, 63
stomach	93, 94, 220	atrophy of mucous mem-	
„ in poisoning	100, 102	brane of stomach	87
„ sometimes in peritonitis	552, 568	with ulcer of stomach	129
„ from destruction		fibroid disease of pylorus	160
of pneumogastric		„ with cancer of stomach	177
nerve	223	„ dyspepsia	202
„ in ulceration of the stomach		„ disease of intestine	302
	124, 221	„ with fistula	305
„ dyspepsia	202	„ with albuminuria	305
„ strangulated intestine	461	„ with dysentery	307
„ as a sign of disease	218	„ with sciatica and hæmor-	
Pancreatic disease	258, 264	rhoids	310
Paralysis of muscles of deglutition	22	Pneumonia, with cancer of œso-	
Parietes, abdominal suppuration in	530	phagus	41
Paracentesis	567, 571, 572	„ with dysentery	413
Pavy, Dr., post-mortem solution	92	„ with peritonitis	560
Pericardium, perforation of, by ulceration of œsophagus	35	Pneumogastric, influence of	6, 157, 223
„ by aneurism	76	„ involved in cancer	
Pericarditis producing dysphagia	78	of œsophagus	41
Peritoneum, pain in affections of	4	„ involved in chronic	
„ inflammation of	547, 554	ulcer of stomach	129, 131
„ adhesions causing obstruction	258	Podophyllin	455
„ strumous disease of	295	Poisons, dysphagia from	25
	549, 552	„ symptoms of	26
„ cancer of	549, 552	„ chloride of zinc	88, 98
„ loose bodies in	575	„ nitric acid	100
Perforation of trachea	33	„ brandy	94
„ pericardium	35	„ sulphuric acid	93
„ œsophagus	80	„ corrosive sublimate	100
„ stomach, 92; from ulceration, 132 to 137; causing local suppuration, 137; and simulating pneumothorax	139	Polypus, œsophageal	78
„ duodenum, 250; from cancer of colon	516	„ in stomach	162
„ colon in dysentery, 404, 405, 406, 412		„ rectal	420
„ ileum in fever	427	Pouch, œsophageal	78
„ from strangulation	465, 496	Prout, Dr., dyspepsia	196
„ from extra uterine fixation	545	Puerperal peritonitis	559, 565
„ of intestine from without	529	Pyæmia, causing peritonitis	559, 565
„ as cause of peritonitis	556	Purgatives	454
		Pylorus, fibroid degeneration	158
		„ cancer of	181, 182
		Pyrosis	153, 193, 194
		RECTUM, discharge of mucus	419
		„ state of, in int. strang.	461
		Remedies, classification of	7
		„ modified by nervous system	9
		Robin, merismopedæia ventriculi	205
		Roper, Mr., cæcal regurgitation	316

	PAGE		PAGE
SCIATICA from hemorrhoids . . .	310	Suppuration in course of portal veins .	54
Sigmoid flexure, cancer of . . .	471	" in parietes of abdomen .	53
Shaw, Mr., loose bodies in peritoneum . . .	575	Supra-renal capsule, cancer of . . .	4
Soap lees, poisoning by . . .	28	" disease, with subacute	
Spine, disease of, as cause of vomiting . . .	235	inflam. of stomach 104, 11	
Strickland, Mr., on constipation . .	452	" with cancer of stomach 11	
Stimulants, effects of, on mucous membrane . . .	186	" as cause of vomiting . .	24
Stokes, Dr., on pericarditis . . .	78	Sympathetic nerve, symptoms of abdominal disease . . .	1
" on ulcer of stomach . . .	154	Syphilis disease, laryngeal cartilages .	1
" on duodenal jaundice . . .	249	" causing ulc. of œsophagus .	3
Stomach, artificial fistula . . .	64	TAYLOR, Dr., on poisons . . .	4
" organic diseases . . .	83	Thompson, Dr. Th., gums in phthisis .	26
" atrophy of m. membrane .	85	" charcoal injection 36	
" emphysema of . . .	85	Throat, diffused inflammation of . .	1
" hypertrophy of . . .	89	Tenia solium . . .	52
" post-mortem solution . . .	99	Trachea, diseased cartilages . . .	3
" inflammation of . . .	98, 102	" perforation of, by cancer 47, 5	
" subacute inflam. of . . .	103	Travers, Mr., cancer of œsophagus .	3
" catarrh . . .	107	" perforation of duodenum 25	
" diphtheritic inflam. of . .	109	Tricocephalus dispar . . .	321, 52
" suppuration in coats of . .	110	Tumours, causing dysphagia . . .	7
" superficial ulc. of . . .	111, 113	" intestinal . . .	57
" follicular ulceration 117, 119		Turnbull, Dr., varieties of fermentative action . . .	8, 20
" perforating . . .	120, 123	Tympanitis in int. strangulation .	46
" involving pneumogastric nerve .	129, 131	" in peritonitis 547, 549, 552, 56	
" term. of ulceration . . .	128	Typhoid disease of intestine 422, 426, 427, 55	
" chronic ulcer, with perf. .	132	Typhus, hæmatemesis . . .	20
" chronic ulcer, with local suppuration . . .	137, 140	ULCERATION of œsophagus . . .	3
" simulat. pneumothorax 139		" stomach . . .	111, 117, 12
" ulc., with fecal abscess 139		" duodenum . . .	25
" ulceration, with perforation of diaphragm . . .	140, 142	" ileum in phthisis . . .	30
" hæmorrhage . . .	143, 208	" cæcum . . .	31
" causes and treatment 149		" colon, dysentery . . .	38
" sloughing of m. memb. 156		Umbilical disch., fecal abscess 301, 57	
" fibroid disease . . .	158	Urine, state of, in int. strangulation .	46
" cancer, 162, 163, 166—168		Uterus, diseases of, as cause of vomiting . . .	23
" functional diseases of . .	185	VERTEBRA, disease of, producing dysphagia . . .	1
" pain as sign of disease . .	218	Vomiting in supra-renal caps. dis. .	10
" vomiting . . .	224	" in ulc. of stomach 125, 15	
Strangulated intestine . . .	459	" as sign of disease . . .	22
Stricture of œsophagus from abscess 34		" in internal strangulation 46	
" " " cancer 36		WEST, Mr. W., syphilitic dysphagia .	3
" " " tumours 74		" Dr., diarrhoea . . .	26
" intestine . . .	459	Wilks, Dr., cases of poisoning . . .	2
Strumous disease of stomach . . .	87	" burns in relation to duodenum . . .	24
" alimentary canal . . .	284	Worms, intestinal . . .	52
" peritonitis, 295, 549, 552, 561, 565		Worthington, Mr. W. C., œsophageal pouch . . .	7
Sulphuric acid, poisoning by 27, 29, 94		Zinc, chloride of, poisoning by 88, 9	
Suppuration in the pharynx . . .	16		
" with strict. of œsophagus 34			
" in coats of stomach . . .	110		
" in dysentery 405, 407, 409, 410			

Post 8vo, cloth, 3s. 6d.,

ON THE INJURIOUS EFFECTS OF MERCURY IN THE TREATMENT OF DISEASE.

By S. O. HABERSHON, M.D. LOND.,


FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS; SENIOR ASSISTANT-
PHYSICIAN AND LECTURER ON MATERIA MEDICA AT
GUY'S HOSPITAL.

"We would recommend practitioners to study this little work, and so test by their own experience, the conclusions of the author."—*British Medical Journal*.

"Dr. Habershon has treated his subject very methodically, very fully, and yet briefly. To every class of medical men this short treatise will be most interesting and suggestive."—*Medical Times and Gazette*.

"The abuse of mercury in the treatment of disease is well and forcibly expounded by Dr. Habershon, in a little book that we strenuously recommend to the perusal and consideration of all believers in that drug. The subject is of too large a character fairly to be discussed in the brief space we can afford to it; but, at the same time, it is of too much importance not to render it imperative upon us to draw the attention of our readers to Dr. Habershon's valuable little book at the earliest opportunity afforded us, and in terms of strong recommendation."—*British and Foreign Medico-Chirurgical Review*.

JOHN CHURCHILL, NEW BURLINGTON STREET.



1 1512

London, New Burlington Street,
April, 1862.

MR. CHURCHILL'S
Publications,
IN
MEDICINE AND OTHER BRANCHES
OF
NATURAL SCIENCE.



"It would be unjust to conclude this notice without saying a few words in favour of Mr. Churchill, from whom the profession is receiving, it may be truly said, the most beautiful series of Illustrated Medical Works which has ever been published."—*Lancet*.

"All the publications of Mr. Churchill are prepared with so much taste and neatness, that it is superfluous to speak of them in terms of commendation."—*Edinburgh Medical and Surgical Journal*.

"No one is more distinguished for the elegance and *recherché* style of his publications than Mr. Churchill."—*Provincial Medical Journal*.

"Mr. Churchill's publications are very handsomely got up: the engravings are remarkably well executed."—*Dublin Medical Press*.

"The typography, illustrations, and getting up are, in all Mr. Churchill's publications, most beautiful."—*Monthly Journal of Medical Science*.

"Mr. Churchill's illustrated works are among the best that emanate from the Medical Press."—*Medical Times*.

"We have before called the attention of both students and practitioners to the great advantage which Mr. Churchill has conferred on the profession, in the issue, at such a moderate cost, of works so highly creditable in point of artistic execution and scientific merit."—*Dublin Quarterly Journal*.

Mr. CHURCHILL is the Publisher of the following Periodicals, offering to Authors a wide extent of Literary Announcement, and a Medium of Advertisement, addressed to all Classes of the Profession. COMMUNICATIONS, Books for REVIEW, addressed to the respective Editors, are received and duly forwarded by Mr. Churchill.

**THE BRITISH AND FOREIGN MEDICO-CHIRURGICAL REVIEW,
AND
QUARTERLY JOURNAL OF PRACTICAL MEDICINE AND SURGERY.**
Price Six Shillings. Nos. I. to LVIII.

**THE QUARTERLY JOURNAL OF MICROSCOPICAL
SCIENCE.**

Edited by EDWIN LANKESTER, M.D., F.R.S., F.L.S., and GEORGE BUSK, F.R.C.S.E.,
F.R.S., F.L.S. Price 4s. Nos. I. to VI. *New Series.*

*. * A few Nos. of the Old Series are out of print; the others may be obtained.

THE JOURNAL OF MENTAL SCIENCE.

Published by authority of the Association of Medical Officers of Asylums and Hospitals for the Insane.

Edited by JOHN CHARLES BUCKNILL, M.D. Published Quarterly, price Half-a-Crown.
New Series. Nos. I. to V.

ARCHIVES OF MEDICINE:

A Record of Practical Observations and Anatomical and Chemical Researches, connected with the Investigation and Treatment of Disease. Edited by LIONEL S. BEALE, M.B., F.R.S.
Published Quarterly, price 2s. 6d. from No. IX.; Nos. I. to VIII., 3s. 6d.

**THE ROYAL LONDON OPHTHALMIC HOSPITAL REPORTS, AND JOURNAL OF
OPHTHALMIC MEDICINE AND SURGERY.**

Published Quarterly, price 2s. Nos. I. to XVII.

THE MEDICAL TIMES AND GAZETTE.

Published Weekly, price Sixpence, or Stamped, Sevenpence.

Annual Subscription, £1. 6s., or Stamped, £1. 10s. 4d., and regularly forwarded to all parts of the Kingdom.

The MEDICAL TIMES AND GAZETTE is favoured with an amount of Literary and Scientific support which enables it to reflect fully the progress of Medical Science, and insure for it a character, an influence, and a circulation possessed at the present time by no Medical Periodical.

**THE HALF-YEARLY ABSTRACT OF THE
MEDICAL SCIENCES.**

Being a Digest of the Contents of the principal British and Continental Medical Works together with a Critical Report of the Progress of Medicine and the Collateral Sciences. Edited by W. H. RANKING, M.D., Cantab., and C. B. RADCLIFFE, M.D., Lond. Post 8vo cloth, 6s. 6d. Vols. I. to XXXIV.

THE PHARMACEUTICAL JOURNAL.

New Series. Published Monthly, price One Shilling.

*. * Vols. I. to XX., bound in cloth, price 12s. 6d. each.

THE BRITISH JOURNAL OF DENTAL SCIENCE.

Published Monthly, price One Shilling. Nos. I. to LXIX.

THE DUBLIN MEDICAL PRESS.

Published Weekly, Stamped, price Sixpence, free to any part of the Empire.

THE MEDICAL DIRECTORY FOR THE UNITED KINGDOM

Published Annually. 8vo. cloth, 10s. 6d.

A CLASSIFIED INDEX TO MR. CHURCHILL'S CATALOGUE.

ANATOMY.

Anatomical Remembrancer	PAGE 3
Beale on Liver	5
Flower on Nerves	11
Hassall's Micros. Anatomy	14
Heale's Anatomy of the Lungs	15
Holden's Human Osteology	15
Do. on Dissections	15
Jones' and Stevking's Pathological Anatomy	17
Maclean's Surgical Anatomy	19
Sibson's Medical Anatomy	25
Waters' Anatomy of Lung	29
Wheeler's Anatomy for Artists	30
Wilson's Anatomy	30

CHEMISTRY.

Abel & Bloxam's Handbook	3
Bowman's Practical Chemistry	7
Do. Medical do.	7
Fownes' Manual of Chemistry	12
Do. Actonian Prize	12
Do. Qualitative Analysis	12
Fresenius' Chemical Analysis	12
Galloway's First Step	12
Do. Analysis	12
Do. Tables	12
Griffiths' Four Seasons	13
Horsley's Chem. Philosophy	16
Jones.—Mulder on Wine	17
Plattner on Blowpipe	22
Speer's Pathol. Chemistry	26

CLIMATE.

Barker on Worthing	4
Bennet on Mentone	6
Dalrymple on the Climate of Egypt	10
Francis on Change of Climate	12
Hall on Torquay	14
Haviland on Climate	14
Lee on Climate	18
McClelland on Bengal	19
McNicoll on Southport	19
Martin on Tropical Climates	20
Moore's Diseases of India	20
Scotchesby-Jackson's Climatology	24
Taylor on Pau	27

DEFORMITIES, &c.

Bigg on Deformities	6
Bishop on Deformities	6
Do. Articulate Sounds	6
Brodhurst on Spine	7
Do. on Clubfoot	7
Godfrey on Spine	13
Hare on Spine	14
Hugman on Hip Joint	16
Inman on Myalgia	16
Tamplyn on Spine	27

DISEASES of the URINARY and GENERATIVE ORGANS, and SYPHILIS.

Acton on Urinary Organs	PAGE 7
Do. on Reproductive Organs	8
Coots on Syphilis	10
Coulson on Bladder	10
Do. on Lithotomy	10
Gant on Bladder	12
Judd on Syphilis	17
Milton on Gonorrhoea	20
Parker on Syphilis	21
Todd on Urinary Organs	28
Wilson on Syphilis	31

DISEASES OF WOMEN AND CHILDREN.

Ballard on Infants and Mothers	4
Barker on Children	4
Bennet on Uterus	6
Do. on Uterine Pathology	6
Bird on Children	7
Blake on the Skin in Children	7
Eyre's Practical Remarks	11
Hood on Scarlet Fever	16
Kiwisch (ed. by Clay) on Ovaries	9
Lee's Ovarian & Uterine Diseases	18
Lee on Diseases of Uterus	18
Do. on Speculum	18
Seymour on Ovaria	25
Smith on Leucorrhoea	26
Tilt on Diseases of Women	28
Do. on Change of Life	28
Underwood on Children	29
West on Women	30

HYGIENE.

Armstrong on Naval Hygiene	4
Beale's Laws of Health	5
Do. Health and Disease	5
Bennet on Nutrition	6
Carter on Training	8
Chavasse's Advice to a Mother	9
Do. Advice to a Wife	9
Granville on Vichy	13
Hartwig on Sea Bathing	14
Do. Physical Education	14
Hufeland's Art	16
Lee's Watering Places of England	18
Do. Rhenish Watering Places	18
Parkin on Disease	21
Pickford on Hygiene	22
Robertson on Diet	24
Routh on Infant Feeding	24
Rumsey's State Medicine	24
Wells' Seamen's Medicine Chest	29
Wife's Domain	30
Wilson on Healthy Skin	31
Do. on Mineral Waters	31
Do. on Turkish Bath	31

MATERIA MEDICA and PHARMACY.

Bateman's Magnacopia	PAGE 5
Beasley's Formulary	5
Do. Receipt Book	5
Do. Book of Prescriptions	5
Pereira's Selecta e Prescriptis	21
Pharmacopoeia Londinensis	22
Prescriber's Pharmacopoeia	22
Royle's Materia Medica	24
Steggall's First Lines for Chemists	26
Stowe's Toxicological Chart	26
Taylor on Poisons	27
Wittstein's Pharmacy	31

MEDICINE.

Adams on Rheumatic Gout	3
Addison on Cella	3
Anderson on Fever	3
Barclay on Medical Diagnosis	4
Barlow's Practice of Medicine	4
Basham on Dropsy	4
Beale on Urine	5
Bird's Urinary Deposits	6
Brinton on Stomach	7
Do. on Ulcer of do.	7
Budd on the Liver	8
Do. on Stomach	8
Camplin on Diabetes	8
Chambers on Digestion	8
Davey's Ganglionic	10
Dobell's Germs and Vestiges of Disease	11
Eyre on Stomach	11
French on Cholera	12
Fuller on Rheumatism	12
Gairdner on Gout	12
Gibb on Throat	13
Granville on Sudden Death	13
Gully's Simple Treatment	13
Habershon on Stomach	13
Do. on Mercury	13
Hall on Apnoea	13
Hall's Observations	13
Harrison on Lead in Water	14
Hassall on Urine	14
Headland on Medicines	14
Hooper's Physician's Vade-Mecum	16
Inman's New Theory	13
James on Laryngoscope	17
Jones' Animal Chemistry	17
Marcel on Chronic Alcoholism	19
Parkes on Urine	21
Peacock on Influenza	21
Richardson's Aesclepiad	23
Roberts on Palsy	24
Robertson on Gout	24
Savory's Compendium	24
Temple on Cough	25
Seymour on Dropsy	25
Shaw's Remembrancer	25
Smee on Debility	25

CLASSIFIED INDEX.

MEDICINE—continued.

Steggall's Medical Manual ..	26
Steggall's Gregory's Conspectus ..	26
Do. Celsus ..	26
Thomas' Practice of Physic ..	27
Thudichum on Urine ..	28
Todd's Clinical Lectures ..	28
Wells on Gout ..	29
What to Observe ..	19
Whitehead on Transmission ..	30
Williams' Principles ..	30
Wright on Headaches ..	31

MICROSCOPE.

Beale on Microscope in Medicine ..	5
Do. How to Work ..	5
Carpenter on Microscope ..	8
Schacht on do. ..	24

MISCELLANEOUS.

Acton on Prostitution ..	8
Bascome on Epidemics ..	4
Bryce on Sebastopol ..	8
Cooley's Cyclopædia ..	9
Forbes' Nature and Art in Disease ..	12
Guy's Hospital Reports ..	13
Haycock's Veterinary ..	14
Lane's Hydropathy ..	18
Lee on Homœop. and Hydrop. ..	18
Marcel on Food ..	19
Maury on Recruits ..	20
Part's Case Book ..	21

NERVOUS DISEASES AND INDIGESTION.

Birch on Constipation ..	6
Carter on Hysteria ..	8
Downing on Neuralgia ..	11
Hunt on Heartburn ..	16
Leared on Imperfect Digestion ..	18
Lobb on Nervous Affections ..	19
Radcliffe on Epilepsy ..	23
Reynolds on the Brain ..	23
Do. on Epilepsy ..	23
Rowe on Nervous Diseases ..	24
Steveking on Epilepsy ..	25
Turnbull on Stomach ..	28

OBSTETRICS.

Barnes on Placenta Prævia ..	4
Davis on Parturition ..	11
Lee's Clinical Midwifery ..	18
Mackenzie on Phlegmasia Do- lens ..	19
Pretty's Aids during Labour ..	22
Priestley on Gravid Uterus ..	22
Ramsbotham's Obstetrics ..	23
Do. Midwifery ..	23
Sinclair & Johnston's Midwifery ..	25
Smellie's Obstetric Plates ..	25
Smith's Manual of Obstetrics ..	26
Swayne's Aphorisms ..	27
Waller's Midwifery ..	29

OPHTHALMOLOGY.

Cooper on Injuries of Eye ..	9
Do. on Near Sight ..	9
Dalrymple on Eye ..	10
Dixon on the Eye ..	11
Hogg on Ophthalmoscope ..	15
Holthouse on Strabismus ..	15
Do. on Impaired Vision ..	15
Hulke on the Ophthalmoscope ..	16
Jacob on Eye-ball ..	16
Jones' Ophthalmic Medicine ..	17
Do. Defects of Sight ..	17
Do. Eye and Ear ..	17
Nunneley on the Organs of Vision ..	21
Walton on the Eye ..	29
Wilde on Malformations of Eye ..	30

PHYSIOLOGY.

Beale on Tissues ..	5
Carpenter's Human ..	8
Do. Comparative ..	8
Do. Manual ..	8
Heale on Vital Causes ..	15
O'Reilly on the Nervous System ..	21
Richardson on Coagulation ..	23
Virchow's (ed. by Chance) Cel- lular Pathology ..	8

PSYCHOLOGY.

Arlidge on the State of Lunacy ..	4
Austin on Paralysis ..	4
Bucknill and Tuke's Psycholo- gical Medicine ..	8
Conolly on Asylums ..	9
Davey on Nature of Insanity ..	10
Dunn's Physiological Psycho- logy ..	11
Hood on Criminal Lunatics ..	16
Millingen on Treatment of In- sane ..	20
Noble on Mind ..	20
Williams (J.) on Insanity ..	30
Williams (J. H.) Unsoundness of Mind ..	30

PULMONARY AND CHEST DISEASES, &c.

Addison on Healthy and Dis- eased Structure ..	3
Allison on Pulmonary Consump- tion ..	3
Billing on Lungs and Heart ..	6
Blakiston on the Chest ..	7
Bright on the Chest ..	7
Cotton on Consumption ..	10
Do. on Stethoscope ..	10
Davies on Lungs and Heart ..	11
Dobell on the Chest ..	11
Fenwick on Consumption ..	11
Laennec on Auscultation ..	18
Markham on Heart ..	20
Richardson on Consumption ..	23
Salter on Asthma ..	24
Skoda on Auscultation ..	20
Thompson on Consumption ..	27
Timms on Consumption ..	28
Turnbull on Consumption ..	28
Weber on Auscultation ..	29

SCIENCE.

Baxter on Organic Polarity ..	4
Bentley's Manual of Botany ..	6
Bird's Natural Philosophy ..	6
Craig on Electric Tension ..	10
Hardwich's Photography ..	14
Hinds' Harmonies ..	15
Jones on Vision ..	17
Do. on Body, Sense, and Mind ..	17
Mayne's Lexicon ..	20
Pratt's Genealogy of Creation ..	23
Price's Photographic Manipula- tion ..	22
Rainey on Shells ..	23
Reynold's Animal Electricity ..	23
Taylor's Medical Jurisprudence ..	21
Vestiges of Creation ..	28
Sequel to ditto ..	21
Unger's Botanical Letters ..	22

SURGERY.

Adams on Reparation of Tendons ..	1
Do. Subcutaneous Surgery ..	1
Anderson on the Skin ..	1
Ashton on Rectum ..	1
Barwell on Diseases of Joints ..	1
Bigg on Artificial Limbs ..	1
Brodburst on Anchylosis ..	1
Bryant on Diseases of Joints ..	1
Chapman on Ulcers ..	1
Do. Varicose Veins ..	1
Cooper (Sir A.) on Testis ..	1
Do. (S.) Surg. Dictionary ..	1
Curling on Rectum ..	1
Do. on Testis ..	1
Druitt's Surgery ..	1
Fergusson's Surgery ..	1
Gray on the Teeth ..	1
Heath's Minor Surgery and Bandaging ..	1
Higginbottom on Nitrate of Silver ..	1
Hodgson on Prostate ..	1
Holt on Stricture ..	1
James on Hernia ..	1
Jordan's Clinical Surgery ..	1
Lawrence on Ruptures ..	1
Liston's Surgery ..	1
Macleod's Surgery of the Crinæa ..	1
Macleae on Fractures ..	1
Maumder's Operative Surgery ..	2
Nunneley on Erysipelas ..	2
Pemberton on Melanosis ..	2
Pirrie's Surgery ..	2
Price on Scrofula ..	2
Smith on Stricture ..	2
Do. on Hemorrhoids ..	2
Snow on Chloroform ..	2
Steggall's Surgical Manual ..	2
Teale on Amputation ..	2
Thompson on Stricture ..	2
Do. on Prostate ..	2
Tomes' Dental Surgery ..	2
Toynbee on Ear ..	2
Wade on Stricture ..	2
Watson on the Larynx ..	2
Webb's Surgeon's Ready Rules ..	2
Williamson on Gunshot Injuries ..	3
Wilson on the Skin ..	3
Do. Portraits of Skin Diseases ..	3
Yearley on Deafness ..	3
Do. on Throat ..	3

MR. F. A. ABEL, F.C.S., & MR. C. L. BLOXAM.
**HANDBOOK OF CHEMISTRY: THEORETICAL, PRACTICAL,
 AND TECHNICAL.** Second Edition. 8vo. cloth, 15s.

MR. ACTON, M.R.C.S.

I.
**A PRACTICAL TREATISE ON DISEASES OF THE URINARY
 AND GENERATIVE ORGANS IN BOTH SEXES.** Third Edition. 8vo. cloth,
 £1. 1s. With Plates, £1. 11s. 6d. The Plates alone, limp cloth, 10s. 6d.

II.
**THE FUNCTIONS AND DISORDERS OF THE REPRODUC-
 TIVE ORGANS IN YOUTH, IN ADULT AGE, AND IN ADVANCED
 LIFE.** Considered in their Physiological, Social, and Psychological Relations. Third
 Edition. 8vo. cloth, 10s. 6d.

III.
PROSTITUTION: Considered in its Moral, Social, and Sanitary Bearings,
 with a View to its Amelioration and Regulation. 8vo. cloth, 10s. 6d.

DR. ADAMS, A.M.
**A TREATISE ON RHEUMATIC GOUT; OR, CHRONIC
 RHEUMATIC ARTHRITIS.** 8vo. cloth, with a Quarto Atlas of Plates, 21s.

MR. WILLIAM ADAMS, F.R.C.S.

I.
**ON THE REPARATIVE PROCESS IN HUMAN TENDONS
 AFTER SUBCUTANEOUS DIVISION FOR THE CURE OF DEFORMITIES.**
 With Plates. 8vo. cloth, 6s.

II.
**SKETCH OF THE PRINCIPLES AND PRACTICE OF
 SUBCUTANEOUS SURGERY.** 8vo. cloth, 2s. 6d.

DR. WILLIAM ADDISON, F.R.S.

I.
CELL THERAPEUTICS. 8vo. cloth, 4s.
 II.
**ON HEALTHY AND DISEASED STRUCTURE, AND THE TRUE
 PRINCIPLES OF TREATMENT FOR THE CURE OF DISEASE, ESPECIALLY CONSUMPTION
 AND SCROFULA,** founded on MICROSCOPICAL ANALYSIS. 8vo. cloth, 12s.

DR. SOMERVILLE SCOTT ALISON, M.D. EDIN., F.R.C.P.
**THE PHYSICAL EXAMINATION OF THE CHEST IN PUL-
 MONARY CONSUMPTION, AND ITS INTERCURRENT DISEASES.** With
 Engravings. 8vo. cloth, 12s.

**THE ANATOMICAL REMEMBRANCER; OR, COMPLETE
 POCKET ANATOMIST.** Fifth Edition, carefully Revised. 32mo. cloth, 3s. 6d.

DR. ANDREW ANDERSON, M.D.
TEN LECTURES INTRODUCTORY TO THE STUDY OF FEVER.
 Post 8vo. cloth, 5s.

DR. MCCALL ANDERSON, M.D.
PARASITIC AFFECTIONS OF THE SKIN. With Engravings.
 8vo. cloth, 5s.

MR. BATEMAN.

MAGNACOPIA: A Practical Library of Profitable Knowledge, communicating the general Minutiae of Chemical and Pharmaceutic Routine, together with the generality of Secret Forms of Preparations; including Concentrated Solutions of Camphor and Copaiba in Water, Mineral Succedaneum, Marmoratum, Silicia, Terro-Metallicum, Pharmaceutic Condensations, Prismatic Crystallization, Crystallized Aromatic Salt of Vinegar, Spa Waters; newly-invented Writing Fluids; Etching on Steel or Iron; with an extensive Variety of *et cætera*. Third Edition. 18mo. 6s.

MR. LIONEL J. BEALE, M.R.C.S.

THE LAWS OF HEALTH IN THEIR RELATIONS TO MIND AND BODY. A Series of Letters from an Old Practitioner to a Patient. Post 8vo. cloth, 7s. 6d.

HEALTH AND DISEASE, IN CONNECTION WITH THE GENERAL PRINCIPLES OF HYGIENE. Fcap. 8vo., 2s. 6d.

DR. BEALE, F.R.S.

ON URINE, URINARY DEPOSITS, AND CALCULI: their Microscopical and Chemical Examination; the Anatomy of the Kidney, and General Remarks on the Treatment of certain Urinary Diseases. Numerous Engravings. Post 8vo. cloth, 8s. 6d.

HOW TO WORK WITH THE MICROSCOPE. Illustrated Edition. Crown 8vo. cloth, 5s. 6d.

THE MICROSCOPE, IN ITS APPLICATION TO PRACTICAL MEDICINE. With a Coloured Plate, and 270 Woodcuts. Second Edition. 8vo. cloth, 14s.

ON THE ANATOMY OF THE LIVER. Illustrated with 66 Photographs of the Author's Drawings. 8vo. cloth, 6s. 6d.

ILLUSTRATIONS OF THE SALTS OF URINE, URINARY DEPOSITS, and CALCULI. 37 Plates, containing upwards of 170 Figures copied from Nature, with descriptive Letterpress. 8vo. cloth, 9s. 6d.

ON THE SIMPLE TISSUES OF THE HUMAN BODY. With Plates, 8vo. cloth, 7s. 6d.

MR. BEASLEY.

THE BOOK OF PRESCRIPTIONS; containing 3000 Prescriptions. Collected from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Second Edition. 18mo. cloth, 6s.

THE DRUGGIST'S GENERAL RECEIPT-BOOK; comprising a copious Veterinary Formulary and Table of Veterinary Materia Medica; Patent and Proprietary Medicines, Druggists' Nostrums, &c.; Perfumery, Skin Cosmetics, Hair Cosmetics, and Teeth Cosmetics; Beverages, Dietetic Articles, and Condiments; Trade Chemicals, Miscellaneous Preparations and Compounds used in the Arts, &c.; with useful Memoranda and Tables. Fifth Edition. 18mo. cloth, 6s.

THE POCKET FORMULARY AND SYNOPSIS OF THE BRITISH AND FOREIGN PHARMACOPŒIAS; comprising standard and approved Formulæ for the Preparations and Compounds employed in Medical Practice. Seventh Edition, corrected and enlarged. 18mo. cloth, 6s.

DR. HENRY BENNET.

I.
A PRACTICAL TREATISE ON INFLAMMATION
OTHER DISEASES OF THE UTERUS. Fourth Edition, revised, with Ad.
8vo. cloth, 16s.

II.
A REVIEW OF THE PRESENT STATE OF UTERINE
PATHOLOGY. 8vo. cloth, 4s.

III.
NUTRITION IN HEALTH AND DISEASE. Post 8vo. cloth, 4s.

IV.
MENTONE AND THE RIVIERA AS A WINTER CLIMATE.
Post 8vo. cloth, 3s. 6d.

PROFESSOR BENTLEY, F.L.S.

A MANUAL OF BOTANY. With nearly 1,200 Engravings on
Feap. 8vo. cloth, 12s. 6d.

MR. HENRY HEATHER BIGG.

I.
THE MECHANICAL APPLIANCES NECESSARY FOR
TREATMENT OF DEFORMITIES. Post 8vo. cloth, 4s.

II.
ARTIFICIAL LIMBS; THEIR CONSTRUCTION AND APPLICATION.
With Engravings on Wood. 8vo. cloth, 3s.

DR. BILLING, F.R.S.

ON DISEASES OF THE LUNGS AND HEART. 8vo. cloth, 4s.

DR. S. B. BIRCH, M.D.

CONSTIPATED BOWELS: the Various Causes and the Rational
of Cure. Post 8vo. cloth, 2s. 6d.

DR. GOLDING BIRD, F.R.S.

I.
URINARY DEPOSITS; THEIR DIAGNOSIS, PATHOLOGY
AND THERAPEUTICAL INDICATIONS. With Engravings on Wood.
Edition. Post 8vo. cloth, 10s. 6d.

II.
ELEMENTS OF NATURAL PHILOSOPHY; being an Experimental
Introduction to the Study of the Physical Sciences. Illustrated with numerous
ings on Wood. Fifth Edition. By GOLDING BIRD, M.D., F.R.S., and C.
BROOKES, M.B. Cantab., F.R.S. Feap. 8vo. cloth, 12s. 6d.

MR. BISHOP, F.R.S.

I.
ON DEFORMITIES OF THE HUMAN BODY, their Pathology
and Treatment. With Engravings on Wood. 8vo. cloth, 10s.

II.
ON ARTICULATE SOUNDS, AND ON THE CAUSES
CURE OF IMPEDIMENTS OF SPEECH. 8vo. cloth, 4s.

MR. P. HINCKES BIRD, F.R.C.S.

PRACTICAL TREATISE ON THE DISEASES OF CHILDREN AND INFANTS AT THE BREAST. Translated from the French of M. BOUCHUT, with Notes and Additions. 8vo. cloth. 20s.

MR. ROBERT HOWARTH BLAKE, M.R.C.S.L.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN IN CHILDREN. From the French of CAILLAULT. With Notes. Post 8vo. cloth, 8s. 6d.

DR. BLAKISTON, F.R.S.

PRACTICAL OBSERVATIONS ON CERTAIN DISEASES OF THE CHEST; and on the Principles of Auscultation. 8vo. cloth, 12s.

MR. JOHN E. BOWMAN.

I.

PRACTICAL CHEMISTRY, including Analysis. With numerous Illustrations on Wood. Fourth Edition. Foolsap 8vo. cloth, 6s. 6d.

II.

MEDICAL CHEMISTRY; with Illustrations on Wood. Third Edition. Fcap. 8vo. cloth, 6s. 6d.

DR. JAMES BRIGHT.

ON DISEASES OF THE HEART, LUNGS, & AIR PASSAGES; with a Review of the several Climates recommended in these Affections. Third Edition. Post 8vo. cloth, 9s.

DR. BRINTON.

I.

THE DISEASES OF THE STOMACH, with an Introduction on its Anatomy and Physiology; being Lectures delivered at St. Thomas's Hospital. Post 8vo. cloth, 10s. 6d.

II.

THE SYMPTOMS, PATHOLOGY, AND TREATMENT OF ULCER OF THE STOMACH. Post 8vo. cloth, 5s.

MR. BERNARD E. BRODHURST, F.R.C.S.

I.

ON LATERAL CURVATURE OF THE SPINE: its Pathology and Treatment. Post 8vo. cloth, with Plates, 3s.

II.

ON THE NATURE AND TREATMENT OF CLUBFOOT AND ANALOGOUS DISTORTIONS involving the TIBIO-TARSAL ARTICULATION. With Engravings on Wood. 8vo. cloth, 4s. 6d.

III.

PRACTICAL OBSERVATIONS ON THE DISEASES OF THE JOINTS INVOLVING ANCHYLOSIS, and on the TREATMENT for the RESTORATION of MOTION. Third Edition, much enlarged, 8vo. cloth, 4s. 6d.

MR. THOMAS BRYANT, F.R.C.S.

ON THE DISEASES AND INJURIES OF THE JOINTS. CLINICAL AND PATHOLOGICAL OBSERVATIONS. Post 8vo. cloth, 7s. 6d.

MR. CHURCHILL'S PUBLICATIONS. 9

MR. H. T. CHAPMAN, F.R.C.S.

I.
**THE TREATMENT OF OBSTINATE ULCERS AND CUTA-
NEOUS ERUPTIONS OF THE LEG WITHOUT CONFINEMENT.** Third
Edition. Post 8vo. cloth, 3s. 6d.

II.
VARICOSE VEINS: their Nature, Consequences, and Treatment, Pallia-
tive and Curative. Post 8vo. cloth, 3s. 6d.

MR. PYE HENRY CHAVASSE, F.R.C.S.

I.
**ADVICE TO A MOTHER ON THE MANAGEMENT OF
HER OFFSPRING.** Sixth Edition. Foolscep 8vo., 2s. 6d.

II.
**ADVICE TO A WIFE ON THE MANAGEMENT OF HER
OWN HEALTH.** With an Introductory Chapter, especially addressed to a Young
Wife. Fourth Edition. Fcap. 8vo., 2s. 6d.

MR. JOHN CLAY, M.R.C.S.

KIWISCH ON DISEASES OF THE OVARIES: Translated, by
permission, from the last German Edition of his Clinical Lectures on the Special Patho-
logy and Treatment of the Diseases of Women. With Notes, and an Appendix on the
Operation of Ovariectomy. Royal 12mo. cloth, 16s.

DR. CONOLLY.

**THE CONSTRUCTION AND GOVERNMENT OF LUNATIC
ASYLUMS AND HOSPITALS FOR THE INSANE.** With Plans. Post 8vo.
cloth, 6s.

MR. COOLEY.

COMPREHENSIVE SUPPLEMENT TO THE PHARMACOPŒIAS.

**THE CYCLOPŒDIA OF PRACTICAL RECEIPTS, AND COL-
LATERAL INFORMATION IN THE ARTS, PROFESSIONS, MANU-
FACTURES, AND TRADES, INCLUDING MEDICINE, PHARMACY, AND
DOMESTIC ECONOMY;** designed as a Compendious Book of Reference for the
Manufacturer, Tradesman, Amateur, and Heads of Families. Third and greatly
enlarged Edition, 8vo. cloth, 26s.

SIR ASTLEY COOPER, BART., F.R.S.

ON THE STRUCTURE AND DISEASES OF THE TESTIS.
With 24 Plates. Second Edition. Royal 4to., 20s.

MR. W. WHITE COOPER.

I.
ON WOUNDS AND INJURIES OF THE EYE. Illustrated by
17 Coloured Figures and 41 Woodcuts. 8vo. cloth, 12s.

II.
**ON NEAR SIGHT, AGED SIGHT, IMPAIRED VISION,
AND THE MEANS OF ASSISTING SIGHT.** With 31 Illustrations on Wood.
Second Edition. Fcap. 8vo. cloth, 7s. 6d.

MR. COOPER.

A DICTIONARY OF PRACTICAL SURGERY AND ENCYCLOPÆDIA OF SURGICAL SCIENCE. New Edition, brought down to the present time. By SAMUEL A. LANE, F.R.C.S., assisted by various eminent Surgeons. Vol. 8vo. cloth, £1. 5s.

MR. HOLMES COOTE, F.R.C.S.

A REPORT ON SOME IMPORTANT POINTS IN THE TREATMENT OF SYPHILIS. 8vo. cloth, 5s.

DR. COTTON.

I.
ON CONSUMPTION: Its Nature, Symptoms, and Treatment. which Essay was awarded the Fothergillian Gold Medal of the Medical Society London. Second Edition. 8vo. cloth, 8s.

II.
PHTHISIS AND THE STETHOSCOPE; OR, THE PHYSICAL SIGNS OF CONSUMPTION. Second Edition. Foolscap 8vo. cloth, 3s.

MR. COULSON.

I.
ON DISEASES OF THE BLADDER AND PROSTATE GLAND. The Fifth Edition, revised and enlarged. 8vo. cloth, 10s. 6d.

II.
ON LITHOTRITY AND LITHOTOMY; with Engravings on Wood. 8vo. cloth, 8s.

MR. WILLIAM CRAIG, L.F.P.S. GLASGOW.

ON THE INFLUENCE OF VARIATIONS OF ELECTRIC TENSION AS THE REMOTE CAUSE OF EPIDEMIC AND OTHER DISEASES. 8vo. cloth, 10s.

MR. CURLING, F.R.S.

I.
OBSERVATIONS ON DISEASES OF THE RECTUM. Second Edition. 8vo. cloth, 5s.

II.
A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM. Second Edition, with Additions. 8vo. cloth, 14s.

DR. DALRYMPLE, M.D. LOND., F.R.C.S.

METEOROLOGICAL AND MEDICAL OBSERVATIONS ON THE CLIMATE OF EGYPT, with Practical Hints for Invalid Travellers. Post 8vo. cloth, 4s.

MR. JOHN DALRYMPLE, F.R.S., F.R.C.S.

PATHOLOGY OF THE HUMAN EYE. Complete in Nine Fasciculi imperial 4to., 20s. each; half-bound morocco, gilt tops, 9l. 15s.

DR. DAVEY.

I.
THE GANGLIONIC NERVOUS SYSTEM: its Structure, Functions, and Diseases. 8vo. cloth, 9s.

II.
ON THE NATURE AND PROXIMATE CAUSE OF INSANITY. Post 8vo. cloth, 3s.

DR. HERBERT DAVIES.

ON THE PHYSICAL DIAGNOSIS OF DISEASES OF THE LUNGS AND HEART. Second Edition. Post 8vo. cloth, 8s.

DR. HALL DAVIS.

ILLUSTRATIONS OF DIFFICULT PARTURITION. Post 8vo. cloth, 6s. 6d.

MR. DIXON.

A GUIDE TO THE PRACTICAL STUDY OF DISEASES OF THE EYE. Second Edition. Post 8vo. cloth, 9s.

DR. DOBELL.

I.
DEMONSTRATIONS OF DISEASES IN THE CHEST, AND THEIR PHYSICAL DIAGNOSIS. With Coloured Plates. 8vo. cloth, 12s. 6d.

II.
LECTURES ON THE GERMS AND VESTIGES OF DISEASE, and on the Prevention of the Invasion and Fatality of Disease by Periodical Examinations. 8vo. cloth, 6s. 6d.

DR. TOOGOOD DOWNING.

NEURALGIA: its various Forms, Pathology, and Treatment. THE JACKSONIAN PRIZE ESSAY FOR 1850. 8vo. cloth, 10s. 6d.

DR. DRUITT, F.R.C.S.

THE SURGEON'S VADE-MECUM; with numerous Engravings on Wood. Eighth Edition. Foolscep 8vo. cloth, 12s. 6d.

MR. DUNN, F.R.C.S.

AN ESSAY ON PHYSIOLOGICAL PSYCHOLOGY. 8vo. cloth, 4s.

SIR JAMES EYRE, M.D.

I.
THE STOMACH AND ITS DIFFICULTIES. Fifth Edition. Fcap. 8vo. cloth, 2s. 6d.

II.
PRACTICAL REMARKS ON SOME EXHAUSTING DISEASES. Second Edition. Post 8vo. cloth, 4s. 6d.

DR. FENWICK.

ON SCROFULA AND CONSUMPTION. Clergyman's Sore Throat, Catarrh, Croup, Bronchitis, Asthma. Fcap. 8vo., 2s. 6d.

MR. FERGUSSON, F.R.S.

A SYSTEM OF PRACTICAL SURGERY; with numerous Illustrations on Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. FLOWER, F.R.C.S.

DIAGRAMS OF THE NERVES OF THE HUMAN BODY, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. Folio, containing Six Plates, 14s.

SIR JOHN FORBES, M.D., D.C.L. (OXON.), F.R.S.
NATURE AND ART IN THE CURE OF DISEASE.
 Edition. Post 8vo. cloth, 6s.

MR. FOWNES, PH.D., F.R.S.

A MANUAL OF CHEMISTRY; with numerous Illustrations of
 Eighth Edition. Fcap. 8vo. cloth, 12s. 6d.

Edited by H. BENCE JONES, M.D., F.R.S., and A. W. HOFMANN, PH.D., F.

**CHEMISTRY, AS EXEMPLIFYING THE WISDOM
 BENEFICENCE OF GOD.** Second Edition. Fcap. 8vo. cloth, 4s. 6d.

INTRODUCTION TO QUALITATIVE ANALYSIS. Post 8vo.

DR. D. J. T. FRANCIS.

CHANGE OF CLIMATE; considered as a Remedy in Dyspepsia, nervous, and other Chronic Affections; with an Account of the most Eligible Residence for Invalids in Spain, Portugal, Algeria, &c., at different Seasons of and an Appendix on the Mineral Springs of the Pyrenees, Vichy, and Aix
 Post 8vo. cloth, 2s. 6d.

MR. J. G. FRENCH, F.R.C.S.

THE NATURE OF CHOLERA INVESTIGATED.
 Edition. 8vo. cloth, 2s.

G. REMIGIUS FRESENIUS.

**ELEMENTARY INSTRUCTION IN CHEMICAL ANALYSIS
 AS PRACTISED IN THE LABORATORY OF GIESSEN.** Edited
 BRUNN, F.R.S.

QUALITATIVE Fifth Edition. 8vo. cloth, 9s.

QUANTITATIVE Third Edition. 8vo. cloth, 16s.

DR. FULLER.

ON RHEUMATISM, RHEUMATIC GOUT, AND SCLETTIC FEVER. their Pathology, Symptoms, and Treatment. Third Edition. 8vo. cloth, 12s.

DR. GARDNER.

ON GOUT: its History, its Causes, and its Cure. Fourth Edition
 8vo. cloth, 2s. 6d.

MR. GALLOWAY.

THE FIRST STEP IN CHEMISTRY. Third Edition. Fcap. 8vo. cloth, 2s.

A MANUAL OF QUALITATIVE ANALYSIS. Third
 Edition. 8vo. cloth, 2s.

CHEMICAL TABLES. On Five Large Sheets, for School and
 Revision. Several Editions. 2s. 6d.

MR. F. J. GANT.

THE INEVITABLE BLAPPER: its Causes and Curative Treatment.
 8vo. cloth, 2s. 6d.

DR. GIBB, M.R.C.P.

ON DISEASES OF THE THROAT, EPIGLOTTIS, AND
WINDPIPE. Post 8vo. cloth, 5s.

MRS. GODFREY.

ON THE NATURE, PREVENTION, TREATMENT, AND CURE
OF SPINAL CURVATURES and DEFORMITIES of the CHEST and LIMBS,
without ARTIFICIAL SUPPORTS or any MECHANICAL APPLIANCES.
Third Edition, Revised and Enlarged. 8vo. cloth, 5s.

DR. GRANVILLE, F.R.S.

THE MINERAL SPRINGS OF VICHY: their Efficacy in the
Treatment of Gout, Indigestion, Gravel, &c. 8vo. cloth, 5s.

II.
ON SUDDEN DEATH. Post 8vo., 2s. 6d.

MR. GRAY, M.R.C.S.

PRESERVATION OF THE TEETH indispensable to Comfort and
Appearance, Health, and Longevity. 18mo. cloth, 3s.

MR. GRIFFITHS.

CHEMISTRY OF THE FOUR SEASONS—Spring, Summer,
Autumn, Winter. Illustrated with Engravings on Wood. Second Edition. Foolscep
8vo. cloth, 7s. 6d.

DR. GULLY.

THE SIMPLE TREATMENT OF DISEASE; deduced from the
Methods of Expectancy and Revulsion. 18mo. cloth, 4s.

DR. GUY.

HOOVER'S PHYSICIAN'S VADE-MECUM; OR, MANUAL OF
THE PRINCIPLES AND PRACTICE OF PHYSIC. New Edition, considerably
enlarged, and rewritten. Foolscep 8vo. cloth, 12s. 6d.

GUY'S HOSPITAL REPORTS. Third Series. Vols. I. to VII., 8vo.,
7s. 6d. each.

DR. HABERSHON, F.R.C.P.

OBSERVATIONS ON DISEASES OF THE ALIMENTARY
CANAL, OESOPHAGUS, STOMACH, CÆCUM, and INTESTINES. 8vo. cloth,
10s. 6d.

II.
ON THE INJURIOUS EFFECTS OF MERCURY IN THE
TREATMENT OF DISEASE. Post 8vo. cloth, 3s. 6d.

DR. MARSHALL HALL, F.R.S.

PRONE AND POSTURAL RESPIRATION IN DROWNING
AND OTHER FORMS OF APNŒA OR SUSPENDED RESPIRATION.
Post 8vo. cloth. 5s.

II.
PRACTICAL OBSERVATIONS AND SUGGESTIONS IN MEDI-
CINE. Second Series. Post 8vo. cloth, 8s. 6d.

DR. HEALE.

I.
A TREATISE ON THE PHYSIOLOGICAL ANATOMY OF
THE LUNGS. With Engravings. 8vo. cloth, 8s.

II.
A TREATISE ON VITAL CAUSES. 8vo. cloth, 9s.

MR. CHRISTOPHER HEATH, F.R.C.S.

A MANUAL OF MINOR SURGERY AND BANDAGING, FOR
THE USE OF HOUSE-SURGEONS, DRESSERS, AND JUNIOR PRAC-
TITIONERS. With Illustrations. Fcap. 8vo. cloth, 5s.

MR. HIGGINBOTTOM, F.R.S., F.R.C.S.E.

I.
AN ESSAY ON THE USE OF THE NITRATE OF SILVER
IN THE CURE OF INFLAMMATION, WOUNDS, AND ULCERS. Second
Edition. Price 5s.

II.
ADDITIONAL OBSERVATIONS ON THE NITRATE OF SIL-
VER; with full Directions for its Use as a Therapeutic Agent. 8vo., 2s. 6d.

DR. HINDS.

THE HARMONIES OF PHYSICAL SCIENCE IN RELATION
TO THE HIGHER SENTIMENTS; with Observations on Medical Studies, and on
the Moral and Scientific Relations of Medical Life. Post 8vo., cloth, 4s.

DR. DECIMUS HODGSON.

THE PROSTATE GLAND, AND ITS ENLARGEMENT IN
OLD AGE. With 12 Plates. Royal 8vo., cloth, 6s.

MR. JABEZ HOGG.

THE OPHTHALMOSCOPE: an Essay on its value in the Exploration
of Internal Eye Diseases. Second Edition. Cloth, 3s. 6d.

MR. LUTHER HOLDEN, F.R.C.S.

I.
HUMAN OSTEOLOGY: with Plates, showing the Attachments of the
Muscles. Third Edition. 8vo. cloth, 16s.

II.
A MANUAL OF THE DISSECTION OF THE HUMAN BODY.
With Engravings on Wood. Second Edition. 8vo. cloth, 16s.

MR. BARNARD HOLT, F.R.C.S.

ON THE IMMEDIATE TREATMENT OF STRICTURE OF
THE URETHRA. 8vo. cloth, 3s.

MR. C. HOLTHOUSE.

I.
ON SQUINTING, PARALYTIC AFFECTIONS OF THE EYE,
and CERTAIN FORMS OF IMPAIRED VISION. Fcap. 8vo. cloth, 4s. 6d.

II.
LECTURES ON STRABISMUS, delivered at the Westminster Hospital.
8vo. cloth, 4s.

DR. W. CHARLES HOOD.
SUGGESTIONS FOR THE FUTURE PROVISION OF CR
NAL LUNATICS. 8vo. cloth, 5s. 6d.

MR. P. HOOD.
THE SUCCESSFUL TREATMENT OF SCARLET FEVER
 also, OBSERVATIONS ON THE PATHOLOGY AND TREATMENT
 CROWING INSPIRATIONS OF INFANTS. Post 8vo. cloth, 5s.

MR. JOHN HORSLEY.
A CATECHISM OF CHEMICAL PHILOSOPHY; being a Fair
 Exposition of the Principles of Chemistry and Physica. With Engravings on
 Designed for the Use of Schools and Private Teachers. Post 8vo. cloth, 6s. 6d.

DR. HUFELAND.
THE ART OF PROLONGING LIFE. Second Edition. E
 by ERASMUS WILSON, F.R.S. Foolscep 8vo., 2s. 6d.

MR. W. OURTIS HUGMAN, F.R.C.S.
ON HIP-JOINT DISEASE; with reference especially to Treat
 by Mechanical Means for the Relief of Contraction and Deformity of the Affected
 8vo. cloth, 3s. 6d.

MR. HULKE, F.R.C.S.
A PRACTICAL TREATISE ON THE USE OF T
OPHTHALMOSCOPE. Being the Jacksonian Prize Essay for 1859. Royal
 cloth, 8s.

DR. HENRY HUNT.
ON HEARTBURN AND INDIGESTION. 8vo. cloth, 5s.

DR. INMAN, M.R.C.P.
ON MYALGIA: ITS NATURE, CAUSES, AND TREATMENT
 being a Treatise on Painful and other Affections of the Muscular System. 3s.
 Edition. 8vo. cloth, 9s.
FOUNDATION FOR A NEW THEORY AND PRACT
OF MEDICINE. Second Edition. Crown 8vo. cloth, 10s.

DR. ARTHUR JACOB, F.R.O.S.
A TREATISE ON THE INFLAMMATIONS OF THE EYE-BA
 Foolscep 8vo. cloth, 5s.

MR. J. H. JAMES, F.R.C.S.
PRACTICAL OBSERVATIONS ON THE OPERATIONS I
STRANGULATED HERNIA. 8vo. cloth, 5s.

DR. PROSSER JAMES, M.D.

SORE-THROAT: ITS NATURE, VARIETIES, AND TREATMENT; including the Use of the LARYNGOSCOPE as an Aid to Diagnosis. Post 8vo. cloth, 4s. 6d.

DR. BENICE JONES, F.R.S.

I.

MULDER ON WINE. Foolsap 8vo. cloth, 6s.

II.

ON ANIMAL CHEMISTRY, in its relation to STOMACH and RENAL DISEASES. 8vo. cloth, 6s.

DR. HANDFIELD JONES, F.R.S., & DR. EDWARD H. SIEVEKING.

A MANUAL OF PATHOLOGICAL ANATOMY. Illustrated with numerous Engravings on Wood. Foolsap 8vo. cloth, 12s. 6d.

MR. WHARTON JONES, F.R.S.

I.

A MANUAL OF THE PRINCIPLES AND PRACTICE OF OPHTHALMIC MEDICINE AND SURGERY; illustrated with Engravings, plain and coloured. Second Edition. Foolsap 8vo. cloth, 12s. 6d.

II.

THE WISDOM AND BENEFICENCE OF THE ALMIGHTY, AS DISPLAYED IN THE SENSE OF VISION; being the Actonian Prize Essay for 1851. With Illustrations on Steel and Wood. Foolsap 8vo. cloth, 4s. 6d.

III.

DEFECTS OF SIGHT: their Nature, Causes, Prevention, and General Management. Fcap. 8vo. 2s. 6d.

IV.

A CATECHISM OF THE MEDICINE AND SURGERY OF THE EYE AND EAR. For the Clinical Use of Hospital Students. Fcap. 8vo. 2s. 6d.

V.

A CATECHISM OF THE PHYSIOLOGY AND PHILOSOPHY OF BODY, SENSE, AND MIND. For Use in Schools and Colleges. Fcap. 8vo., 2s. 6d.

MR. FURNEAUX JORDAN, M.R.C.S.

AN INTRODUCTION TO CLINICAL SURGERY; WITH A Method of Investigating and Reporting Surgical Cases. Fcap. 8vo. cloth, 5s.

MR. JUDD.

A PRACTICAL TREATISE ON URETHRITIS AND SYPHILIS; including Observations on the Power of the Menstruous Fluid, and of the Discharge from Leucorrhœa and Sores to produce Urethritis: with a variety of Examples, Experiments, Remedies, and Cures. 8vo. cloth, £1. 5s.

DR. LAENNEC.

A MANUAL OF AUSCULTATION AND PERCUSSION. Translated and Edited by J. B. SHARPE, M.R.C.S. 3s.

DR. LANE, M.A.

HYDROPATHY; OR, HYGIENIC MEDICINE. An Explana Essay. Second Edition. Post 8vo. cloth, 5s.

MR. LAWRENCE, F.R.S.

A TREATISE ON RUPTURES. The Fifth Edition, considerably enlarged. 8vo. cloth, 16s.

DR. LEARED, M.R.O.P.

IMPERFECT DIGESTION: ITS CAUSES AND TREATMENT. Second Edition. Foolscap 8vo. cloth, 3s. 6d.

DR. EDWIN LEE.

I.

THE EFFECT OF CLIMATE ON TUBERCULOUS DISEASES, with Notices of the chief Foreign Places of Winter Resort. Small 8vo. cloth, 4s. 6c.

II.

THE WATERING PLACES OF ENGLAND, CONSIDERED with Reference to their Medical Topography. Fourth Edition. Foolscap 8vo. cloth, 7s. 6d.

III.

THE BATHS OF RHENISH GERMANY. Post 8vo. cloth, 3s.

IV.

HOMŒOPATHY AND HYDROPATHY IMPARTIALLY APPRECIATED. With Notes illustrative of the Influence of the Mind over the Body. Fourth Edition. Post 8vo. cloth, 3s. 6d.

DR. ROBERT LEE, F.R.S.

I.

A TREATISE ON THE SPECULUM; with Three Hundred Cases. 8vo. cloth, 4s. 6d.

II.

CLINICAL REPORTS OF OVARIAN AND UTERINE DISEASES, with Commentaries. Foolscap 8vo. cloth, 6s. 6d.

III.

CLINICAL MIDWIFERY: comprising the Histories of 545 Cases of Difficult, Preternatural, and Complicated Labour, with Commentaries. Second Edition. Foolscap 8vo. cloth, 5s.

IV.

PRACTICAL OBSERVATIONS ON DISEASES OF THE UTERUS. With coloured Plates. Two Parts. Imperial 4to., 7s. 6d. each Part.

MR. LISTON, F.R.S.
PRACTICAL SURGERY. Fourth Edition. 8vo. cloth, 22s.

MR. H. W. LOBB, L.S.A., M.R.O.S.E.
ON SOME OF THE MORE OBSCURE FORMS OF NERVOUS AFFECTIONS, THEIR PATHOLOGY AND TREATMENT. With an Introduction on the Physiology of Digestion and Assimilation, and the Generation and Distribution of Nerve Force. Based upon Original Microscopical Observations. With Engravings. 8vo. cloth, 10s. 6d.

LONDON MEDICAL SOCIETY OF OBSERVATION.
WHAT TO OBSERVE AT THE BED-SIDE, AND AFTER DEATH. Published by Authority. Second Edition. Foolscep 8vo. cloth, 4s. 6d.

DR. MACKENZIE, M.D., M.R.O.P.
THE PATHOLOGY AND TREATMENT OF PHLEGMASIA DOLENS, as deduced from Clinical and Physiological Researches. Being the Lettsomian Lectures on Midwifery, delivered before the Medical Society of London during the Session 1861-62. 8vo. cloth, 6s.

MR. M'CLELLAND, F.L.S., F.G.S.
SKETCH OF THE MEDICAL TOPOGRAPHY, OR CLIMATE AND SOILS, OF BENGAL AND THE N. W. PROVINCES. Post 8vo. cloth, 4s. 6d.

DR. GEORGE H. B. MACLEOD, F.R.C.S. (EDIN.)
NOTES ON THE SURGERY OF THE CRIMEAN WAR; with REMARKS on GUN-SHOT WOUNDS. 8vo. cloth, 10s. 6d.

MR. JOSEPH MACLISE, F.R.C.S.
I.
SURGICAL ANATOMY. A Series of Dissections, illustrating the Principal Regions of the Human Body.
The Second Edition, imperial folio, cloth, £3. 12s.; half-morocco, £4. 4s.

II.
ON DISLOCATIONS AND FRACTURES. This Work is Uniform with the Author's "Surgical Anatomy;" each Fasciculus contains Four beautifully executed Lithographic Drawings. Imperial folio, cloth, £2. 10s.; half-morocco, £2. 17s.

DR. MONICOLL, M.R.C.P.
A HAND-BOOK FOR SOUTHPORT, MEDICAL & GENERAL; with Copious Notices of the Natural History of the District. Second Edition. Post 8vo. cloth, 3s. 6d.

DR. MARCET, F.R.S.
I.
ON THE COMPOSITION OF FOOD, AND HOW IT IS ADULTERATED; with Practical Directions for its Analysis. 8vo. cloth, 6s. 6d.

II.
ON CHRONIC ALCOHOLIC INTOXICATION, OR ALCOHOLIC STIMULANTS IN CONNEXION WITH THE NERVOUS SYSTEM. Foolscep 8vo. cloth, 3s. 6d.

MR. NUNNELEY, F.R.O.S.E.

I.
ON THE ORGANS OF VISION: THEIR ANATOMY AND PHYSIOLOGY. With Plates, 8vo. cloth, 15s.

II.
A TREATISE ON THE NATURE, CAUSES, AND TREATMENT OF ERYSIPELAS. 8vo. cloth, 10s. 6d.

DR. O'REILLY.

THE PLACENTA, THE ORGANIC NERVOUS SYSTEM, THE BLOOD, THE OXYGEN, AND THE ANIMAL NERVOUS SYSTEM, PHYSIOLOGICALLY EXAMINED. With Engravings. 8vo. cloth, 5s.

MR. LANGSTON PARKER.

THE MODERN TREATMENT OF SYPHILITIC DISEASES, both Primary and Secondary; comprising the Treatment of Constitutional and Confirmed Syphilis, by a safe and successful Method. Fourth Edition, 8vo. cloth, 10s.

DR. PARKES, F.R.C.P.

THE URINE: ITS COMPOSITION IN HEALTH AND DISEASE, AND UNDER THE ACTION OF REMEDIES. 8vo. cloth, 12s.

DR. PARKIN.

THE CAUSATION AND PREVENTION OF DISEASE.
8vo. cloth, 5s.

MR. JAMES PART, F.R.C.S.

THE MEDICAL AND SURGICAL POCKET CASE BOOK, for the Registration of important Cases in Private Practice, and to assist the Student of Hospital Practice. Second Edition. 3s. 6d.

DR. THOMAS B. PEACOCK, M.D.

ON THE INFLUENZA, OR EPIDEMIC CATARRHIAL FEVER OF 1847-8. 8vo. cloth, 5s. 6d.

MR. OLIVER PEMBERTON, M.R.O.S.

OBSERVATIONS ON THE HISTORY, PATHOLOGY, AND TREATMENT OF CANCEROUS DISEASES. Part I.—MELANOSIS. With coloured Plates. Royal 8vo. cloth, 4s. 6d.

DR. PEREIRA, F.R.S.

SELECTA E PRÆSCRIPTIS: with a Key, containing the Prescriptions in an Unabbreviated Form, and a Literal Translation. Thirteenth Edition. 24mo. cloth, 5s.

MR. P. C. PRICE, F.R.C.S.E.

SCROFULOUS DISEASES OF THE EXTERNAL LYMPHATIC GLANDS: their Nature, Variety, and Treatment; with Remarks on the Management of Scrofulous Ulcerations, Scars, and Cicatrices. Post 8vo. cloth, 3s. 6d.

DR. RADCLIFFE, F.R.C.P. LOND.

ON EPILEPTIC AND OTHER CONVULSIVE AFFECTIONS OF THE NERVOUS SYSTEM. Third Edition. Post 8vo. cloth, 7s. 6d.

MR. RAINEY.

ON THE MODE OF FORMATION OF SHELLS OF ANIMALS, OF BONE, AND OF SEVERAL OTHER STRUCTURES, by a Process of Molecular Coalescence, Demonstrable in certain Artificially-formed Products. Fcap. 8vo. cloth, 4s. 6d.

DR. F. H. RAMSBOTHAM.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY. Illustrated with One Hundred and Twenty Plates on Steel and Wood; forming one thick handsome volume. Fourth Edition. 8vo. cloth, 22s.

DR. RAMSBOTHAM.

PRACTICAL OBSERVATIONS ON MIDWIFERY, with a Selection of Cases. Second Edition. 8vo. cloth, 12s.

DR. DU BOIS REYMOND.

ANIMAL ELECTRICITY; Edited by H. BENGE JONES, M.D., F.R.S. With Fifty Engravings on Wood. Foolscep 8vo. cloth, 6s.

DR. REYNOLDS, M.D., LOND.

I.
EPILEPSY: ITS SYMPTOMS, TREATMENT, AND RELATION TO OTHER CHRONIC CONVULSIVE DISEASES. 8vo. cloth, 10s.

II.
THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND THEIR APPENDAGES. 8vo. cloth, 8s.

DR. B. W. RICHARDSON.

I.
ON THE CAUSE OF THE COAGULATION OF THE BLOOD. Being the ASTLEY COOPER PRIZE ESSAY for 1856. With a Practical Appendix. 8vo. cloth, 16s.

II.
THE HYGIENIC TREATMENT OF PULMONARY CONSUMPTION. 8vo. cloth, 5s. 6d.

III.
THE ASCLEPIAD. Vol. I., Clinical Essays. 8vo. cloth, 6s. 6d.

MR. WILLIAM ROBERTS.

AN ESSAY ON WASTING PAISY; being a Systematic Treatise on the Disease hitherto described as **ATROPHIE MUSCULAIRE** PROGRESS With Four Plates. 8vo. cloth, 7s. 6d.

DR. W. H. ROBERTSON.

THE NATURE AND TREATMENT OF GOUT.
8vo. cloth, 10s. 6d.

A TREATISE ON DIET AND REGIMEN.

Fourth Edition. 2 vols. post 8vo. cloth, 12s.

DR. ROUTH.

INFANT FEEDING, AND ITS INFLUENCES ON LIFE
Or, the Causes and Prevention of Infant Mortality. Fcap. 8vo. cloth, 5s.

DR. ROWE.

NERVOUS DISEASES, LIVER AND STOMACH COMPLAINTS, LOW SPIRITS, INDIGESTION, GOUT, ASTHMA, AND ORDERS PRODUCED BY TROPICAL CLIMATES. With Cases. Sixth Edition. Fcap. 8vo. 2s. 6d.

DR. ROYLE, F.R.S., AND DR. HEADLAND, M.D.

A MANUAL OF MATERIA MEDICA AND THERAPEUTICS
With numerous Engravings on Wood. Third Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. RUMSEY, F.R.C.S.

ESSAYS ON STATE MEDICINE. 8vo. cloth, 10s. 6d.

DR. SALTER, F.R.S.

ON ASTHMA: its Pathology, Causes, Consequences, and Treatment. 8vo. cloth, 10s.

MR. SAVORY.

A COMPENDIUM OF DOMESTIC MEDICINE, AND CONNECTION TO THE MEDICINE CHEST; comprising Plain Directions for the Employment of Medicines, with their Properties and Doses, and Brief Descriptions of Symptoms and Treatment of Diseases, and of the Disorders incidental to Infants and Children, with a Selection of the most efficacious Prescriptions. Intended as a Guide of Easy Reference for Clergymen, and for Families residing at a Distance from Medical Assistance. Fifth Edition. 12mo. cloth, 5s.

DR. SCHACHT.

THE MICROSCOPE, AND ITS APPLICATION TO VEGETABLE ANATOMY AND PHYSIOLOGY. Edited by FREDERICK CURREY, M.A. 8vo. cloth, 6s.

DR. SCORESBY-JACKSON, M.D., F.R.S.E.

MEDICAL CLIMATOLOGY; or, a Topographical and Meteorological Description of the Localities resorted to in Winter and Summer by Invalids of various classes both at Home and Abroad. With an Isothermal Chart. Post 8vo. cloth,

DR. SEMPLE.

ON COUGH: its Causes, Varieties, and Treatment. With some practical Remarks on the Use of the Stethoscope as an aid to Diagnosis. Post 8vo. cloth, 4s. 6d.

DR. SEYMOUR.

ILLUSTRATIONS OF SOME OF THE PRINCIPAL DISEASES OF THE OVARIA: their Symptoms and Treatment; to which are prefixed Observations on the Structure and Functions of those parts in the Human Being and in Animals. With 14 folio plates, 12s.

THE NATURE AND TREATMENT OF DROPSY; considered especially in reference to the Diseases of the Internal Organs of the Body, which most commonly produce it. 8vo. 5s.

MR. SHAW, M.R.C.S.

THE MEDICAL REMEMBRANCER; OR, BOOK OF EMERGENCIES: in which are concisely pointed out the Immediate Remedies to be adopted in the First Moments of Danger from Poisoning, Apoplexy, Burns, and other Accidents; with the Tests for the Principal Poisons, and other useful Information. Fourth Edition. Edited, with Additions, by JONATHAN HUTCHINSON, M.R.C.S. 32mo. cloth, 2s. 6d.

DR. SIBSON, F.R.S.

MEDICAL ANATOMY. With coloured Plates. Imperial folio. Fasciculi I. to VI. 5s. each.

DR. E. H. SIEVEKING.

ON EPILEPSY AND EPILEPTIFORM SEIZURES: their Causes, Pathology, and Treatment. Second Edition. Post 8vo. cloth, 10s. 6d.

MR. SINCLAIR AND DR. JOHNSTON.

PRACTICAL MIDWIFERY: Comprising an Account of 13,748 Deliveries, which occurred in the Dublin Lying-in Hospital, during a period of Seven Years. 8vo. cloth, 15s.

MR. ALFRED SMEE, F.R.S.

GENERAL DEBILITY AND DEFECTIVE NUTRITION; their Causes, Consequences, and Treatment. Fcap. 8vo. cloth, 3s. 6d.

DR. SMELLIE.

OBSTETRIC PLATES: being a Selection from the more Important and Practical Illustrations contained in the Original Work. With Anatomical and Practical Directions. 8vo. cloth, 5s.

MR. HENRY SMITH, F.R.C.S.

ON STRICTURE OF THE URETHRA. 8vo. cloth, 7s. 6d.

HÆMORRHOIDS AND PROLAPSUS OF THE RECTUM: Their Treatment by the Application of Nitric Acid. Second Edition. Fcap. 8vo. cloth, 3s.

DR. SWAYNE.

OBSTETRIC APHORISMS FOR THE USE OF STUDENTS
COMMENCING MIDWIFERY PRACTICE. With Engravings on Wood. Second
Edition. Fcap. 8vo. cloth, 3s. 6d.

MR. TAMPLIN, F.R.C.S.E.

LATERAL CURVATURE OF THE SPINE: its Causes, Nature, and
Treatment. 8vo. cloth, 4s.

DR. ALEXANDER TAYLOR, F.R.S.E.

THE CLIMATE OF PAU; with a Description of the Watering Places
of the Pyrenees, and of the Virtues of their respective Mineral Sources in Disease. Third
Edition. Post 8vo. cloth, 7s.

DR. ALFRED S. TAYLOR, F.R.S.

I.

A MANUAL OF MEDICAL JURISPRUDENCE. Seventh Edition.
Fcap. 8vo. cloth, 12s. 6d.

II.

ON POISONS, in relation to MEDICAL JURISPRUDENCE AND
MEDICINE. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

MR. TEALE.

**ON AMPUTATION BY A LONG AND A SHORT RECTAN-
GULAR FLAP.** With Engravings on Wood. 8vo. cloth, 5s.

DR. THEOPHILUS THOMPSON, F.R.S.

I.

CLINICAL LECTURES ON PULMONARY CONSUMPTION.
With Plates. 8vo. cloth, 7s. 6d.

II.

LETTSONIAN LECTURES ON PULMONARY CONSUMPTION;
with Remarks on Microscopical Indications, and on Cocoa-nut Oil. Post 8vo., 2s. 6d.

DR. THOMAS.

THE MODERN PRACTICE OF PHYSIC; exhibiting the Symp-
toms, Causes, Morbid Appearances, and Treatment of the Diseases of all Climates.
Eleventh Edition. Revised by ALGERNON FRAMPTON, M.D. 2 vols. 8vo. cloth, 28s.

MR. HENRY THOMPSON, F.R.O.S.

I.

STRICTURE OF THE URETHRA; its Pathology and Treatment.
The Jacksonian Prize Essay for 1852. With Plates. Second Edition. 8vo. cloth, 10s.

II.

THE DISEASES OF THE PROSTATE; their Pathology and Treat-
ment. Comprising a Dissertation "On the Healthy and Morbid Anatomy of the Prostate
Gland;" being the Jacksonian Prize Essay for 1860. With Plates. Second Edition.
8vo. cloth, 10s.

DR. UNDERWOOD.

TREATISE ON THE DISEASES OF CHILDREN. Tenth Edition,
with Additions and Corrections by HENRY DAVIES, M.D. 8vo. cloth, 15s.

DR. UNGER.

BOTANICAL LETTERS. Translated by Dr. B. PAUL. Numerous
Woodcuts. Post 8vo., 2s. 6d.

MR. WADE, F.R.C.S.

**STRICTURE OF THE URETHRA, ITS COMPLICATIONS
AND EFFECTS;** a Practical Treatise on the Nature and Treatment of those
Affections. Fourth Edition. 8vo. cloth, 7s. 6d.

DR. WALLER.

ELEMENTS OF PRACTICAL MIDWIFERY; or, Companion to
the Lying-in Room. Fourth Edition, with Plates. Fcap. cloth, 4s. 6d.

MR. HAYNES WALTON, F.R.C.S.

SURGICAL DISEASES OF THE EYE. With Engravings on
Wood. Second Edition. 8vo. cloth, 14s.

DR. WATERS, M.R.C.P.

THE ANATOMY OF THE HUMAN LUNG. The Prize Essay
to which the Fothergillian Gold Medal was awarded by the Medical Society of London.
Post 8vo. cloth, 6s. 6d.

DR. EBEN. WATSON, A.M.

**ON THE TOPICAL MEDICATION OF THE LARYNX IN
CERTAIN DISEASES OF THE RESPIRATORY AND VOCAL ORGANS.**
8vo. cloth, 5s.

DR. ALLAN WEBB, F.R.C.S.L.

**THE SURGEON'S READY RULES FOR OPERATIONS IN
SURGERY.** Royal 8vo. cloth, 10s. 6d.

DR. WEBER.

**A CLINICAL HAND-BOOK OF AUSCULTATION AND PER-
CUSSION.** Translated by JOHN COCKLE, M.D. 5s.

MR. T. SPENCER WELLS, F.R.C.S.

**PRACTICAL OBSERVATIONS ON GOUT AND ITS COMPLI-
CATIONS,** and on the Treatment of Joints Stiffened by Gouty Deposits. Foolsap 8vo.
cloth, 5s.

II.

**SCALE OF MEDICINES WITH WHICH MERCHANT VES-
SELS ARE TO BE FURNISHED,** by command of the Privy Council for Trade;
With Observations on the Means of Preserving the Health of Seamen, &c. &c.
Seventh Thousand. Fcap. 8vo. cloth, 3s. 6d.

MR. ERASMUS WILSON, F.R.S. (*continued*).

II.

DISEASES OF THE SKIN: A Practical and Theoretical Treatise on the DIAGNOSIS, PATHOLOGY, and TREATMENT OF CUTANEOUS DISEASES. Fourth Edition. 8vo. cloth, 16s.

THE SAME WORK; illustrated with finely executed Engravings on Steel, accurately coloured. 8vo. cloth, 34s.

III.

HEALTHY SKIN: A Treatise on the Management of the Skin and Hair in relation to Health. Sixth Edition. Foolsap 8vo. 2s. 6d.

IV.

PORTRAITS OF DISEASES OF THE SKIN. Folio. Fasciculi I. to XII., completing the Work. 20s. each.

V.

ON SYPHILIS, CONSTITUTIONAL AND HEREDITARY; AND ON SYPHILITIC ERUPTIONS. With Four Coloured Plates. 8vo. cloth, 16s.

VI.

A THREE WEEKS' SCAMPER THROUGH THE SPAS OF GERMANY AND BELGIUM, with an Appendix on the Nature and Uses of Mineral Waters. Post 8vo. cloth, 6s. 6d.

VII.

THE EASTERN OR TURKISH BATH: its History, Revival in Britain, and Application to the Purposes of Health. Foolsap 8vo., 2s.

DR. G. C. WITTSTEIN.

PRACTICAL PHARMACEUTICAL CHEMISTRY: An Explanation of Chemical and Pharmaceutical Processes, with the Methods of Testing the Purity of the Preparations, deduced from Original Experiments. Translated from the Second German Edition, by STEPHEN DARBY. 18mo. cloth, 6s.

DR. HENRY G. WRIGHT.

HEADACHES; their Causes and their Cure. Third Edition. Fcap. 8vo. 2s. 6d.

MR. YEARSLEY.

I.

DEAFNESS PRACTICALLY ILLUSTRATED; being an Exposition of Original Views as to the Causes and Treatment of Diseases of the Ear. Fifth Edition. Foolsap 8vo., 2s. 6d.

II.

ON THE ENLARGED TONSIL AND ELONGATED UVULA, and other Morbid Conditions of the Throat. Seventh Edition. 8vo. cloth, 5s.



BOUND BY
EDMOND & REMNANT
LONDON

